

Historic, Archive Document

Do not assume content reflects current scientific knowledge, policies, or practices.

Reserve
aSB951
.5
.B48

AD-83 Bookplate
(1-68)

NATIONAL

**A
G
R
I
C
U
L
T
U
R
A
L**




LIBRARY

COOPERATIVE
IMPACT ASSESSMENT
REPORT

THE BIOLOGIC AND ECONOMIC
ASSESSMENT OF

MEVINPHOS



United States
Department of
Agriculture


In cooperation with
State Agricultural
Experiment Stations

Technical Bulletin
Number XXXX

Cooperative Extension
Service

Other State Agencies

U.S. Environmental
Protection Agency



**THE BIOLOGIC AND ECONOMIC
ASSESSMENT OF**

MEVINPHOS

A report of the Mevinphos Assessment Team
to the Special Review of Mevinphos

Submitted to the Environmental Protection
Agency on August 15, 1989

United States
Department of
Agriculture

In cooperation with
State Agricultural
Experiment Stations

Cooperative Extension
Service

Other State Agencies

U.S. Environmental
Protection Agency

Technical Bulletin
Number XXXX

PREFACE

This report is a joint project of the U.S. Department of Agriculture, the State Land-Grant Universities, and the U.S. Environmental Protection Agency. This report is prepared by a team of scientists from these organizations to provide sound, current scientific information on the benefits of, and exposure to, mevinphos.

The report is a scientific presentation to be used in connection with other data as a portion of the total body of knowledge in a final benefit/risk assessment under the Review process in connection with the Federal Insecticide, Fungicide, and Rodenticide Act as Revised in October 1988.

Sincere appreciation is extended to the Assessment Team members and to all others who gave so generously in the development of information and in the preparation of the report.

Mevinphos Assessment Team

Clyde S. Gorsuch	Entomologist	Clemson University Clemson, South Carolina
Richard A. Goyer	Entomologist	Louisiana State University Baton Rouge, Louisiana
Armon J. Keaster	Entomologist	University of Missouri Columbia, Missouri
George G. Kennedy	Entomologist	N.C. State University Raleigh, North Carolina
Ken Kido	Entomologist	University of California Riverside, California
John M. Love	Economist	Cornell University Ithaca, New York
Eugene E. Nelson	Entomologist	Colorado State University Grand Junction, Colorado
Bobby C. Pass	Team Leader Entomologist	University of Kentucky Lexington, Kentucky
Robert L. Pienkowski	Entomologist	VPI and SU Blacksburg, Virginia
Stanley D. Pilcher	Entomologist	Colorado State University Akron, Colorado
Roscoe Randell	Entomologist	University of Illinois Champaign, Illinois

Kenneth A. Sorensen Entomologist

N.C. State University
Raleigh, North Carolina

Gerald B. White Economist

Cornell University
Ithaca, New York

ACKNOWLEDGMENTS

Appreciation is extended to the following people for their assistance in providing information on the uses of mevinphos, acreage treated, production costs, comparative efficiency of mevinphos and available alternative insecticides, the losses associated with inadequate control of the various insect pests, and other related information.

Steve L. Brown	Auburn University Auburn, Alabama
Wayne Vandre	University of Alaska Anchorage, Alaska
Gary Cramer	University of Arizona Tucson, Arizona
Bill F. Jones	University of Arkansas Fayetteville, Arkansas
Harold G. Alford	University of California Davis, California
Bert Bohmont Whitney Cranshaw	Colorado State University Fort Collins, Colorado
Candace L. Bartholomew	University of Connecticut West Hartford, Connecticut
Mark R. Graustein	University of Delaware Newark, Delaware
William B. Easley	University of the District of Columbia Washington, D. C.
O. Norman Nesheim F. A. Johnson W. F. Martinkovic	University of Florida Gainesville, Florida
David B. Adams	University of Georgia Tifton, Georgia
R. Muniappan	University of Guam Mangilao, Guam
Larry Sandvol	University of Idaho Aberdeen, Idaho
Roscoe Randell	University of Illinois Champaign, Illinois

David L. Matthew	Purdue University West Lafayette, Indiana
Harold J. Stockdale	Iowa State University Ames, Iowa
Donald C. Cress	Kansas State University Manhattan, Kansas
Lorraine D. Rodriguez	University of Kentucky Lexington, Kentucky
D. K. Pollet	Louisiana State University Baton Rouge, Louisiana
Glen Koehler	University of Maine Orono, Maine
James J. Linduska	University of Maryland Salisbury, Maryland
Donald C. Weber	University of Massachusetts Amherst, Massachusetts
Larry G. Olsen	Michigan State University East Lansing, Michigan
Phillip K. Harein	University of Minnesota St. Paul, Minnesota
James H. Jarratt	Mississippi State University Mississippi State, Mississippi
Henry F. DiCarlo Ray Nabor	University of Missouri Portageville, Missouri
Flerney G. Jones James W. Johnson	University of Missouri Columbia, Missouri
Gregory D. Johnson	Montana State University Bozeman, Montana
Shripat T. Kamble	University of Nebraska Lincoln, Nebraska
Harry G. Smith	University of Nevada Reno, Nevada
James S. Bowman	University of New Hampshire Durham, New Hampshire

George C. Hamilton	Rutgers University New Brunswick, New Jersey
Donald A. Rutz	Cornell University Ithaca, New York
Kenneth A. Sorensen	North Carolina State University Raleigh, North Carolina
Dean K. McBride	North Dakota State University Fargo, North Dakota
Acie C. Waldron	Ohio State University Columbus, Ohio
Bob Cartwright	Oklahoma State University Lane, Oklahoma
John Rinehold	Oregon State University Corvallis, Oregon
Manual Silveira	NORPAC Foods Salem, Oregon
Ben Simko	County Agent Ontario, Oregon
Dennis Calvin	The Pennsylvania State University University Park, Pennsylvania
Hipolito O'Farrill-Nieves	University of Puerto Rico Mayaguez, Puerto Rico
Steven R. Alm	University of Rhode Island Kingston, Rhode Island
Randall P. Griffin Robert G. Bellinger Clyde S. Gorsuch	Clemson University Clemson, South Carolina
B. H. Kantack	South Dakota State University Brookings, South Dakota
Carroll J. Southards Jaime Yanes, Jr. Edward E. Burgess	University of Tennessee Knoxville, Tennessee
Rodney L. Holloway	Texas A&M University College Station, Texas
Howard Deer	Utah State University Logan, Utah

George B. MacCollom	University of Vermont Burlington, Vermont
Geoff Zehuder	Virginia Polytechnic Inst. & State University Painter, Virginia
Walter I. Knausenberger	University of The Virgin Islands Kingshill, St. Croix
Dan Mayer	Washington State University Prosser, Washington
John F. Baniecki	West Virginia University Morgantown, West Virginia
Katherine P. Schmidt Phil Pellitteri	University of Wisconsin Madison, Wisconsin
Mark A. Ferrell	University of Wyoming Laramie, Wyoming
Craig Osteen	ERS-USDA Washington, D.C.

TABLE OF CONTENTS

Title Page	i
Preface	ii
Acknowledgements	iv
Contents	viii
Executive Summary	1
Introduction	5
Synopses of the Crops, Pests and Use Parameters	10
Pesticide Characterization	22
Mode of Action	22
Physical and Chemical Properties	22
Formulations Marketed	22
Physiological Activity in Soil and Plant and Animal Tissue	22
Methodology of the Benefits Assessment Study	24
Extent of Use	28
Specific Use Analysis	29
Fruit and Nut Crops	29
Apples/Peaches/Pears/Plums.....	31
Cherries	32
Citrus: Oranges/Lemons/Grapefruit	32
Grapes	32
Raspberries	36
Strawberries	36
Walnuts	36
Field Crops	38
Alfalfa for Hay	38

Alfalfa Seed-Production	40
Field Corn	44
Sorghum	50
Vegetable Crops	53
Artichokes	54
Beans (Snap, Lima)	54
Beets	56
Broccoli	56
Cabbage	56
Cauliflower	62
Carrots	62
Celery	66
Corn, Sweet	66
Cucumbers	70
Eggplant	73
Peppers	73
Lettuce	73
Melons	73
Okra	78
Onions	83
Parsley	83
Peas	86
Potatoes	86
Spinach	86
Squash	92

Tomatoes	95
Leafy Greens - Turnip/Mustard/Kale	95
Leafy Greens - Collards	99
Greenhouse	102
Aerial Application	103
Sewage Plants	104
Economic and Social Impacts	105
Methodology and Data	105
Quantitative Use of Mevinphos	106
Yield, Price and Crop Value Benefits	111
Consumer Benefits	115
Macroeconomic Effects	116
Social/Community Impacts	116
Exposure Considerations and their Relationship to Risk Analysis	118
Recommendations	120
References	122
Appendix I, Survey Questionnaire	170
Appendix II, Survey Responses by Commodity, U.S. Production	175
Appendix III, Documentation and Assumptions for Survey Adjustment	198
Appendix IV, Price-quantity Relationships	207

EXECUTIVE SUMMARY

Mevinphos is a highly toxic organophosphorus insecticide that is registered for use on 109 different sites against a wide variety of pests. It may enter the body by ingestion or by dermal contact; it is absorbed by plant roots, translocated readily to leaves and growing shoots, and degraded rapidly. As a systemic insecticide, it is translocated unchanged. Metabolism does not increase its toxicity. A moderate teratogenic effect was produced when mevinphos was injected into hen eggs before or during the early period of incubation. However, the acute toxicity (oral LD₅₀ for rats is 3.7 to 12 mg/kg; acute dermal LD₅₀ for rabbits is 5.3 to 33.8 mg/kg) is of major concern and the trigger for this review.

The principal risk currently known to be associated with mevinphos is that of acute poisoning due to extremely high oral and dermal mammalian toxicity. Poisoning reports from California indicate that mevinphos was among the top five pesticides causing occupational poisoning in the state during 1981-1985. The main avenue of worker applicator exposure was through dermal contact, with more limited inhalation exposure. The California data show 165 mevinphos related poisonings during the period 1980-1986 with an average of 82 poisonings per million pounds of mevinphos used. Because of its high acute toxicity to avian, mammalian and aquatic species, mevinphos is classified as a Category I pesticide.

A questionnaire developed to assess the benefits of mevinphos was designed to elicit key information regarding the present use of mevinphos. Each state was requested to complete a questionnaire for each crop/site for which the pesticide was used in significant quantities. Data were collected relating to the averages of production, costs and use of mevinphos for the crop years 1982-1986. Questionnaires were sent to each of the 50 states plus territories of the United States as well as Washington, DC. Responses were received from all 50 states as well as the Virgin Islands, Guam, Washington, D.C.; and Puerto Rico, and the results serve as the basis of this assessment.

Mevinphos had important commercial use on 33 crops in 25 states. During 1982-1986, an annual average of 666,723 acres were treated with 756,815 pounds of active ingredient of mevinphos. The various use sites include small fruits, citrus, tree fruits and nuts, various vine crops, vegetables, field crops, ornamental plants and sewage systems. The greatest use was on lettuce, followed by uses on strawberry, broccoli, greens (collards, kale, turnip and mustard), cabbage, alfalfa seed, and cauliflower. The amount of mevinphos used on these 7 commodities amounted to 632,751 pounds annually of active ingredient, or 83.6% of the total combined use of mevinphos in the United States during the period covered in the study.

A significant percentage of the total acreage of the following crops grown in the United States was treated with mevinphos: alfalfa seed (69%), broccoli (55%), Brussels sprouts (97%), cabbage (56%), southern pea (83%), cauliflower (67%), celery (51%), greens (75%), cucumber (30%), muskmelons (70%), lettuce (90%), onions (55%), parsley (44%), pepper (47%), strawberry (43%), artichoke (96%), okra (70%) and spinach (62%). Of the 756,815 pounds active ingredient of mevinphos used in the United States, the greatest amount

was used on vegetables (594,777 pounds a.i.), followed by fruits (103,977 pounds a.i.) and field crops (58,061 pounds a.i.). The geographic regions where mevinphos was used in the greatest quantities were the West (445,400 pounds a.i.) and the South (107,035 pounds a.i.), which probably represent treatments on alfalfa for seed production, strawberry, and vegetable crops.

Comparative static equilibrium analysis was used to demonstrate the benefits of mevinphos. The estimated impact of mevinphos use was determined by comparing yields, prices, and production with the expected use of mevinphos's alternatives.

The economic benefits of mevinphos are higher yields and lower costs relative to many other insecticides available for controlling pests on major vegetable and fruit crops. In addition, mevinphos is important in a few major field crops.

The benefits of mevinphos were calculated as the sum of reduced revenues plus added costs that would result from the withdrawal of mevinphos. Presumably, growers are now using mevinphos because it is more profitable than alternative pesticides or control methods.

The yield benefit of mevinphos with respect to price is assumed to be a result of higher supplies and lower prices to consumers. That is, considering total production, the higher yield causes prices to be lower, holding other factors constant. For producers of a commodity exhibiting relatively stable prices, higher production means higher total crop value. At the farm level, opposite changes in production and price are offsetting in celery, lettuce, muskmelons, strawberry, and spinach. Due to the estimated inflexibility of prices for broccoli and especially cauliflower, decreases in U. S. production are not fully offset by increases in price. Therefore, the yield benefit of mevinphos is greater on these two crops compared to the others. The benefit of mevinphos use, based on changes in crop value, totals \$59 million. The largest benefit comes from use on cauliflower, followed by broccoli, celery, lettuce, and strawberry.

In addition to the benefit of mevinphos on crop value, the net change in production costs for mevinphos's alternatives amounts to \$1.9 million. The major cost increases of substituting chemical and other means to control pests previously controlled by mevinphos would occur on cabbage (+\$1.9 million), greens (+\$1.6 million) and strawberry (+\$1.3 million). Offsetting reductions in costs for substituting mevinphos's alternatives could occur in lettuce (-\$1.7 million), alfalfa seed production (-\$0.8 million), and cauliflower (-\$0.5 million).

The average annual net benefit of mevinphos in U. S. crop production, adding the change in total crop value to the net change in control costs, is \$61 million, or about \$91 per acre treated. This estimate of total benefit is sensitive to different estimates of yield change and price flexibility. For example, a lower absolute value for price flexibility produces a higher value of any yield benefit from mevinphos use. Also, the net difference in mevinphos's alternative control costs are assumed to be borne entirely by the producer and not passed on to the consumer. The availability of mevinphos is

likely to be a risk reducing (and thus cost reducing) opportunity for producers, suggesting an unmeasured benefit of mevinphos use.

The benefits to consumers of a product treated with mevinphos are assumed to be measured by the difference in retail price that would be paid for the same product treated with mevinphos's alternatives. Based on the farm-level yield and price effects of mevinphos, assuming mevinphos is unavailable and assuming the price change is passed on to the retail level of distribution, consumers benefit from mevinphos use on broccoli, cabbage, cauliflower, celery, greens, cantaloupe and honeydew (other melons), lettuce, strawberry and spinach.

The consumers' total benefit from mevinphos use results from the lower prices paid for total production caused by the lower marginal value of the increased yield. The product of the change in price times the total production of crops whose yield is increased by using mevinphos equals \$94 million, comprising broccoli (\$4 million), cabbage (\$7 million), cauliflower (\$3.6 million), celery (\$3.7 million), greens (\$2.6 million), muskmelons (\$1.6 million), lettuce (\$65 million), strawberry (\$5.6 million), and spinach (\$0.9 million).

The total benefit of mevinphos use is the sum of the increased production costs expected with mevinphos's alternatives (\$1.9 million), the higher farm value of production (\$59 million), and the higher consumer value of production (\$94 million). The total benefit of mevinphos use is \$155 million.

Of course, in the event that mevinphos is unavailable, higher prices will put pressure on the market to adjust by increasing supply, possibly through increasing imports or through changes in production practices. The analysis presented herein does not attempt to estimate market adjustments; it is intended to indicate where the pressure for adjustment is more likely to occur.

Employment effect might occur in local areas where particular crops are important might occur. These could result from one or both of the following: (1) a crop may no longer be grown in specific local areas if mevinphos is unavailable; and (2) reduction in volume or shutdown in packing houses and with shippers where the rejection rate for produce due to insect damage becomes high or due to the loss of a crop. These are very important events to the affected individuals and communities, but are fairly minor for individual states or nationally. For the most part, crops on which mevinphos is used (i.e. fresh produce) are produced in areas where soil and climate are favorable for a wide range of crops. Therefore, there are usually alternative uses for land and labor.

Mevinphos is the chemical of choice for many vegetable crops because of its short residual life and quick degradation characteristics which make possible the achievement of legal tolerances within a short pre-harvest interval. This feature is important in all vegetable crops; but is especially critical in artichokes, where the only registered alternative for plume moth control has a 30 day pre-harvest interval; in broccoli during head

formation; and in other vegetables where harvest is continuous over an interval of several days; in celery, where there is no registered alternative for a broad spectrum of pests at harvest; in okra and in leafy greens, where there is no single alternative superior to mevinphos for a broad spectrum of pests. This is especially true for collards and for spinach, where mevinphos is the only "cleanup" treatment close to harvest and in lettuce, where all the reporting states considered mevinphos the standard for control of aphids and the lepidopterous pest complex.

Mevinphos is either preferred or considered to have an important role in production of green and snap beans, cabbage, carrots, cauliflower, melons, peas, and potatoes. It is not considered essential for production of beets, eggplant, peppers, onions, cucumbers, sweet corn, squash or tomatoes, where alternatives are available. However, for several of these crops, mevinphos is the least expensive and/or the most effective of the registered chemicals. Mevinphos confers sizable yield benefits in production of many vegetables and is considered a valuable tool in resistance management because of its quick action and short residual.

In fruit crops, mevinphos is considered necessary for strawberry production, less so in production of apples and grapes.

KEYWORDS: Mevinphos, Phosdrin, insecticides, pesticide registration, aphids, mites, grasshoppers, Lygus bugs, red-banded leafroller, fruit tree leafroller, orange tortrix, omnivorous leafroller, leaf folder, leafhoppers, strawberry leafroller, salt-marsh caterpillar, Western tussock moth, alfalfa caterpillar, climbing cutworms, alfalfa weevil, cabbage looper, imported cabbage worm, false chinch bug, dipterous leafminer, plume moth, Mexican bean beetle, thrips, tobacco budworm, green stink bug, velvet bean caterpillar, corn earworm, apples, peaches, pears, plums, cherries (sour), oranges, lemons, grapefruit, grapes, raspberries, strawberries, walnuts, alfalfa, clover, corn, sorghum, mustard greens, turnip tops, pea vines, artichoke, beans, beets, broccoli, cabbage, Brussels sprouts, cauliflower, collards, kale, carrots, celery, cucumbers, egg plant, peppers, lettuce, melons, okra, onions, parsley, peas, potatoes, spinach, summer squash, tomatoes, turnips.

INTRODUCTION

The Mevinphos Benefits Assessment Team was organized by the USDA, NAPIAP Program in June 1988 for the purpose of developing an accurate, objective summary of the uses and benefits of mevinphos to agricultural productivity in the United States and to examine the economic and social impacts of present and continued use of mevinphos. This benefits assessment was initiated in anticipation of a Special Review of Mevinphos by the United States Environmental Protection Agency. As of July 1989, a Special Review has not been initiated.

Mevinphos is the common name of the insecticidally active chemical 3-[(dimethoxyphosphinyl) oxy] - butenoic acid methyl ester. Mevinphos is a broad spectrum, fast acting, short residual organophosphate insecticide originally developed by Shell Chemical Company in 1953 (Thomson 1977). Its pesticide characteristics are summarized in the report that follows. Other names for mevinphos include Apavinphos, Duraphos, Gestid, Menite, Mevinox, OS-2046, Phosdrin and Phosfene (Thomson 1977, EPA Pesticide Fact Sheet No. 156 {1988}). In the United States, mevinphos is currently produced by AMVAC Chemical Corporation and was marketed by Shell Chemical Company until recently when E. E. DuPont de Nemours and Company acquired license to the Phosdrin 4EC and Phosdrin 1PA4 formulations of mevinphos from Shell. More recently, DuPont has announced that it will not pursue reregistration of mevinphos. AMVAC Chemical Corporation, however, has indicated plans to pursue reregistration of mevinphos for selected agricultural uses (source: Jack Prieur, AMVAC Chemical Corporation, 4100 East Washington Blvd., Los Angeles, CA 90023).

Because mevinphos was developed and initially registered for use as an insecticide more than 30 years ago, and because several companies have been involved, the Assessment Team has been able to develop neither a detailed chronology of registrations, nor an accurate historical picture of mevinphos use. However, through the use of questionnaires sent to all State Pesticide Impact Assessment Liaison Representatives in the United States, we have developed what we believe to be an accurate assessment of the current usage patterns of mevinphos in the United States.

Over the years, mevinphos usage in the United States has averaged between 500,000 and 750,000 pounds active ingredient per year, with the fluctuations being related to variation in acreages of various major use crops, such as lettuce, and in insect pest populations (source: Jack Prieur, AMVAC Chemical Corporation). As of May 1988 there were 134 products containing mevinphos registered in the United States for use on a total of 109 sites (Table 1). Most (98) of these sites were represented by agricultural commodities; but there were an additional 7 sites represented by ornamental plant groups and 4 sites associated with sewage systems. Of the agricultural sites, there were 14 in the fruit and nut grouping, 29 in the forage crop grouping and 55 in the vegetable crop grouping. Responses to our survey questionnaires indicated that mevinphos is currently used on only a small proportion of registered sites. Those uses and the benefits associated with them are discussed in detail in the sections that follow.

For insect control on agricultural and ornamental plants, mevinphos is applied as a foliar spray using ground or aerial equipment. Recommended application rates typically range from 0.125 to 1.0 pound a.i./acre but reach 2.0 pounds a.i./acre for control of Western tussock moth larvae, citrus cutworm, variegated cutworm, and pink scavenger caterpillar on citrus (Phosdrin 4EC and Phosdrin IPA4 labels).

The principal risk currently known to be associated with mevinphos is that of acute poisoning due to its extremely high oral and dermal mammalian toxicity (acute oral LD₅₀ for rats 3.7 to 12 mg/kg; acute dermal LD₅₀ for rabbits 5.3 to 33.8 mg/kg {Wiswesser 1976}). Poisoning reports from California indicate that mevinphos was among the top five pesticides causing occupational poisoning in that state during 1981-1985. The main avenue of worker (applicator) exposure is through dermal contact, with more limited inhalation exposure (EPA Pesticide Fact Sheet No. 156, 1988). The California data show 165 mevinphos related poisonings during 1980-1986, with an average of 82 poisonings per million pounds of mevinphos used. On average, an individual suffering mevinphos poisoning spent 53 days in the hospital (Jay S. Ellenberger, USEPA - presentation at June 28, 1988 Assessment Team Meeting in San Antonio). Because of its high acute toxicity to avian, mammalian and aquatic species, mevinphos is classified as a Category I pesticide.

In the issuance of a Registration Standard, the Environmental Protection Agency (EPA) cited major data gaps in the areas of toxicology, environmental fate, and residue chemistry. Because of extensive data gaps, the EPA has indicated that no significant new uses will be granted until it has received data sufficient to evaluate dietary exposure to mevinphos (EPA Pesticide Fact Sheet No. 156, 1988). In addition the EPA enacted more restrictive re-entry restrictions changing them from "Do not enter treated areas without protective clothing until sprays have dried" (Phosdrin 4EC and Phosdrin IPA4 labels 1988) to: "four days for citrus groves, grape vineyards and nectarine and peach orchards and 2 days for all other treated areas unless protective clothing specified on the label is worn" (EPA Pesticide Fact Sheet No. 156, 1988).

In the document that follows, we present a description of the methods used by the Assessment Team in acquiring and analyzing data on mevinphos use and benefits followed by a thorough discussion of uses deemed significant based on the amount of crops/acreage treated per year. For each use, we identify the specific target insects and identify potential alternatives to mevinphos use. Finally, we present an analysis of the economic and social impacts of mevinphos use and a discussion of the consequences of mevinphos not being available.

Table 1. Registered Sites for Mevinphos Use.

Fruit and Nut Crops

Apples (Foliar Treatment)
Cherries (Foliar Treatment)
Cherries (Sour) (Foliar Treatment)
Citrus (Foliar Treatment)
Grapefruit (Foliar Treatment)
Grapes (Foliar Treatment)
Lemons (Foliar Treatment)
Oranges (Foliar Treatment)
Peaches (Foliar Treatment)
Pears (Foliar Treatment)
Plums (Foliar Treatment)
Raspberries (Foliar Treatment)
Strawberries (Foliar Treatment)
Walnuts (Foliar Treatment)

Forage Crops

Alfalfa (Foliar Treatment)
Barley (Foliar Treatment)
Beans (Forage - Fodder) (Foliar Treatment)
Birdsfoot Trefoil (Foliar Treatment)
Celery (Foliar Treatment)
Clover (Foliar Treatment)
Corn (Field) (Forage) (Foliar Treatment)
Corn (Field) (Grain) (Foliar Treatment)
Corn (Forage) (Foliar Treatment)
Corn (Pop) (Foliar Treatment)
Corn (Pop) (Forage) (Foliar Treatment)
Corn (Pop) (Grain) (Foliar Treatment)
Corn (Sweet) (Forage) (Foliar Treatment)
Corn (Foliar Treatment)
Corn (Field) (Foliar Treatment)
Fennel (Foliar Treatment)
Flax (Foliar Treatment)
Oats (Foliar Treatment)
Peas (Foliar Treatment)
Pea Vines (Forage) (Foliar Treatment)
Sesame (Seed Crop Foliar Treatment)
Small Grains (Foliar Treatment)
Sorghum (Foliar Treatment)
Sorghum (Forage) (Foliar Treatment)
Sorghum (Grain)
Sorghum (Grain) (Foliar Treatment)
Wheat (Foliar Treatment)

Table 1 Continued

Vegetable Crops

Artichokes (Foliar Treatment)
 Asparagus (Foliar Treatment)
 Beans (Foliar Treatment)
 Beans (Lima) (Foliar Treatment)
 Beans (Green) (Foliar Treatment)
 Beans (Snap) (Foliar Treatment)
 Beets (Foliar Treatment)
 Beets (Greens) (Foliar Treatment)
 Broccoli (Foliar Treatment)
 Brussels Sprouts (Foliar Treatment)
 Brussels Sprouts (Soil Treatment)
 Cabbage (Foliar Treatment)
 Cantaloupes
 Cantaloupes (Foliar Treatment)
 Carrots (Foliar Treatment)
 Cauliflower (Foliar Treatment)
 Chinese Cabbage (Foliar Treatment)
 Collards (Foliar Treatment)
 Corn (Sweet) (Foliar Treatment)
 Corn (Sweet) (Grain) (Foliar Treatment)
 Cucumbers (Foliar Treatment)
 Eggplant (Foliar Treatment)
 Endive (Escarole) (Foliar Treatment)
 Honeydew Melons (Foliar Treatment)
 Kale (Foliar Treatment)
 Lettuce (Foliar Treatment)
 Lettuce (Bibb) (Foliar Treatment)
 Lettuce (Bibb) (Greenhouse)
 Lettuce (Head) (Foliar Treatment)
 Lettuce (Leaf) (Foliar Treatment)
 Lettuce (Leaf) (Greenhouse)
 Lettuce (Romaine) (Foliar Treatment)
 Lettuce (Romaine) (Greenhouse)
 Melons (Foliar Treatment)
 Melons (Soil Treatment)
 Muskmelon (Foliar Treatment)
 Mustard (Greens) (Foliar Treatment)
 Okra (Foliar Treatment)
 Onions (Foliar Treatment)
 Onions (Dry) (Foliar Treatment)
 Onions (Green) (Foliar Treatment)
 Parsley (Foliar Treatment)
 Peppers (Foliar Treatment)
 Potatoes (Foliar Treatment)
 Pumpkin (Foliar Treatment)

Table 1 Continued.

Vegetable Crops

Spinach (Foliar Treatment)
Squash (Foliar Treatment)
Squash (Summer) (Foliar Treatment)
Squash (Winter) (Foliar Treatment)
Tomatoes (Foliar Treatment)
Turnips (Foliar Treatment)
Turnips (Greens) (Foliar Treatment)
Turnips (Roots) (Foliar Treatment)
Watercress (Foliar Treatment)
Watermelon (Foliar Treatment)

Ornamental Uses

Bermudagrass (Seed Crop Foliar Treatment)
Ornamental Herbaceous Plants (Foliar Treatment)
Ornamental Plants (Foliar Treatment)
Ornamental Plants (Field) (Foliar Treatment)
Ornamental Woody Shrubs (Foliar Treatment)
Ornamental Woody Vines (Foliar Treatment)
Waterlilies (Foliar Treatment)

Sewage Uses

Sewage Biofilters
Sewage Plants
Sewage Systems
Sewage Systems (Water Treatment)

SYNOPSIS OF THE CROPS, PESTS AND USE PARAMETERS FOR MEVINPHOS

FRUIT AND NUT CROPS

Use recommended dosages in sufficient water to obtain thorough spray coverage, but do not exceed indicated maximum gallonage for that crop. If concentrate ground application is employed, do not apply less than 10 gallons of spray per acre, and adjust dosage to obtain same dosage per acre as recommended for full coverage application. Do not apply with hand equipment.

Formulations: 4EC, IPA4, raspberries, 4EC only

Apples/peaches/pears/plums

Aphids, mites, grasshoppers,
Lygus bugs, red-banded
leafroller

Federal label directions (ground equipment)

Apples/peaches/pears/plums

Aphids and mites - Apply 1/4 to 1/2 pt. (0.125 lb. to 0.25 lb. a.i.) per 100 gal. water, 40 to 800 gal./acre recommended. Do not exceed 1200 gal. of finished spray/acre on apples and plums when 1/2 pt. is used.

Grasshoppers, Lygus bugs and red-banded leafroller - Apply 1/2 to 2/3 pt. (0.25 to 0.33 lb. a.i.) per 100 gal. water, 40 to 800 gal./acre recommended. Do not exceed 900 gal. of finished spray/acre on apples and pears when 2/3 pt. is used. Do not exceed 750 gal. of finished spray/acre on peaches or plums.

Do not treat apples, pears, or plums within 2 days of harvest or peaches within 4 days of harvest or apply more than 5 pts./acre on peaches and plums, 6 1/2 pts/acre on apples and pears.

Peaches only. Workers should not be permitted to enter treated areas to engage in any activity requiring substantial contact with treated foliage for 4 days following treatment. When a mixture of two or more organophosphate pesticides is applied, the re-entry interval shall be extended by adding to the longest applicable interval an additional 50% of that interval.

Cherries (sour only)

Aphids and mites - Apply 1/2 to 2/3 pt. (0.25 to 0.33 lb. a.i.)/100 gal. water.

Do not exceed 750 gal finished spray/acre.

Do not treat within 2 days of harvest or apply more than 5 pt./acre.

Citrus: oranges/lemons/grapefruit

Aphids, fruit tree leafroller,
orange tortrix and omnivorous
leafroller, Western tussock
moth larvae, citrus cutworm,

variegated cutworm, pink
scavenger caterpillar

Citrus: oranges/lemons/grapefruit

Aphids - Apply 1 to 2 pt. (0.5 to 1.0 lb. a.i.)/acre in 200 or more gal. water.

Fruit tree leafroller, orange tortrix and omnivorous leafroller -
Apply 1 qt. (1.0 lb. a.i.)/acre in 500 gal. water.

Western tussock moth larvae, citrus cutworm, variegated cutworm and pink scavenger caterpillar - Apply 2 qt. (2.0 lb. a.i.)/acre in 1200 gal. water.

Allow at least 7 days between applications. Do not treat within 4 days of harvest. Workers should not be permitted to enter treated areas to engage in any activity requiring substantial contact with treated foliage for 4 days following treatment. When a mixture of two or more organophosphate pesticides is applied, the re-entry interval shall be extended by adding to the longest applicable interval an additional 50% of that interval.

Grapes	Aphids, leaf folder, leafhoppers, mites, red-banded leafroller and Lyqus bugs
--------	---

Grapes

Aphids - Apply 1/2 to 1 pt. (0.25 to 0.5 lb. a.i.)/acre. Do not treat within 2 days of harvest.

Leaf folder, leafhoppers, mites, red-banded leafroller and Lygus bugs -
Apply 1 to 2 pts. (0.5 - 1.0 a.i.)/acre. At the 2 pt. dosage do not apply
within 5 days of harvest.

Workers should not be permitted to enter treated areas to engage in any activity requiring substantial contact with treated foliage for 4 days following treatment. When a mixture of two or more organophosphate pesticides is applied, the re-entry interval shall be extended by adding to the longest applicable interval an additional 50% of that interval.

Raspberries	Aphids, mites, leafhoppers, fruit tree leafroller and orange tortrix
-------------	--

Raspberries

Aphids - Apply 1/4 to 1/2 pt. (0.125 to 0.25 lb. a.i.)/100 gal. water, up to 200 gal. water/acre.

Mites, leafhoppers, fruit tree leafroller and orange tortrix - Apply 1/2 to 2/3 pt. (0.25 to 0.33 lb. a.i.) per 100 gal. water, up to 200 gal. water/acre.

Do not treat within 3 days of harvest.

Strawberries

Aphids, mites, grasshoppers, strawberry leafrollers, salt-marsh caterpillar and Lygus bugs

Strawberries

Aphids and mites - Apply 1/4 to 1/2 pt. (0.125 to 0.5 lb. a.i.) per 100 gal. water. Do not treat within 2 days of harvest.

Grasshoppers, Strawberry leafroller, salt-marsh caterpillar and Lygus bugs - Apply 1/2 to 1 pt. (0.25 to 0.5 lb. a.i.) per 100 gal. water.

Do not treat within 2 days of harvest.

For hard-to-kill insects including aphids and mites, use 1 qt./100 gal. water but do not treat within 2 days of harvest or apply more than 1 qt./acre. Do not exceed 100 gal. finished spray/acre.

Walnuts

Aphids, mites, omnivorous looper, orange tortrix, fruit tree leafroller and Western tussock moth

Walnuts

Aphids - Apply 1/4 to 1/2 pt. (0.125 to 0.25 lb. a.i.)/100 gal. water.

Mites, omnivorous looper, orange tortrix, fruit tree leafroller and Western tussock moth - Apply 1/2 pt. (0.25 lb. a.i.)/100 gal. water.

Do not treat within 2 days of harvest. Do not exceed 1,000 gal. finished spray, or 5 pts./acre.

FIELD CROPS

Formulations: 4EC, IPA4; pea vines, 4EC only

Alfalfa/clover

Aphids, alfalfa caterpillar, grasshoppers, leafhoppers, cutworms (climbing), mites, alfalfa weevil larvae, and Lygus bugs

Federal label directions (ground equipment)

Alfalfa/clover

Aphids and alfalfa caterpillar - Apply 1/4 to 1/2 pt. (0.125 to 0.25 lb. a.i.) in 10-125 gal. water/acre

Grasshoppers, leafhoppers, cutworms (climbing), mites, alfalfa weevil larvae, and lygus bugs - Apply 1/2 to 1 pt. (0.25 to 0.5 lb. a.i.) in 10-

125 gal. water/acre.

Do not treat within 2 days of harvest.

Corn (field, sweet and popcorn)
for forage only.

Aphids

Corn (field, sweet and popcorn) for forage only

Aphids - Apply 1/2 to 1 pt. (0.25 to 0.5 lb. a.i.) in 10-125 gal. water/acre.

Sorghum (for forage and grain)

Aphids, corn earworm, webworm
and fall armyworm

Sorghum (for forage and grain)

Aphids - Apply 1/4 to 1/2 pt. (0.125 to 0.25 lb. a.i.) in 10-125 gal. water/acre.

Corn earworm and webworm - Apply 1/2 to 1 pt. (0.25 to 0.5 lb. a.i.) in 10-125 gal. water/acre.

Fall armyworm - Apply 1 pt. (0.5 lb. a.i.) in 10-125 gal. water/acre.
Do not treat within 3 days of harvest.

Pea vines (for forage only)

Aphids, grasshoppers,
leafhoppers, mites and cutworms
(climbing)

Pea vines (for forage only)

Aphids - Apply 1/4 to 1/2 pt. (0.125 to 0.25 lb. a.i.) in 10-125 gal. water/acre.

Grasshoppers, leafhoppers, mites and cutworms (climbing) - Apply at 1/2 to 1 pt. (0.25 to 0.5 lb. a. i.) in 10-125 gal. water/acre.
Do not treat within 2 days of harvest.

VEGETABLE CROPS

Formulations: 4EC, IPA4; okra and parsley, 4EC only

Federal label directions (ground equipment)

Artichokes

Aphids, plume moth

Artichokes

Aphids - Apply at 1/4 to 1/2 pt. (0.125 to 0.25 lb. a. i.) in 10-125 gal. water/acre.

Plume moth - Apply at 1 to 2 pt. (0.5 - 1.0 lb. a.i.) in 10-125 gal. water/acre.

Do not treat within 2 days of harvest.

Beans

Aphids, grasshoppers,
leafhoppers, mites and Mexican
bean beetle

Beans

Aphids - Apply at 1/4 to 1/2 pt. (0.125 to 0.25 lb. a.i.) in 10-125 gal. water/acre.

Grasshoppers, leafhoppers, mites and Mexican bean beetle - Apply at 1/2 to 1 pt. (0.25 to 0.5 lb. a.i.) in 10-125 gal. water/acre.
Do not treat within 2 days of harvest.

Beets (including tops)

Aphids, cabbage looper, cutworm
(climbing), dipterous leafminer
(adult), false chinch bug,
grasshoppers, imported cabbage
worm, leafhopper, mites and
salt-marsh caterpillar

Beets (including tops)

Aphids - Apply at 1/4 to 1/2 pt. (0.125 to 0.25 lb. a.i.) in 10-125 gal. water/acre.

Cabbage looper, cutworm (climbing), dipterous leaf miner (adult), false chinch bug, grasshoppers, imported cabbage worm, leafhoppers, mites and salt-marsh caterpillar - Apply at 1/2 to 1 pt. (0.25 to 0.5 lb. a.i.) in 10-125 gal. water/acre.
Do not treat within 3 days of harvest.

Broccoli, cabbage

Aphids, cabbage looper, imported
cabbage worm, grasshoppers,
leafhoppers, salt-marsh cater-
pillar, mites, cutworms
(climbing), dipterous leafminer
(adult) and Lygus bugs

Broccoli, cabbage

Aphids - Apply at 1/4 to 1/2 pt. (0.125 to 0.25 lb. a.i.) in 10-125 gal. water/acre.

Cabbage looper, imported cabbage worm, grasshoppers, leafhoppers, salt-marsh caterpillar, mites, cutworms (climbing), dipterous leafminer (adult) and Lygus bugs - Apply at 1/2 to 1 pt. (0.25 to 0.5 lb. a.i.) in 10-125 gal. water/acre.
Do not treat within 2 days of harvest.

For hard-to-kill insects, including aphids, apply at 1 qt. (1.0 lb. a.i.) in 10-125 gal. water/acre, but do not treat within 3 days of harvest.

Brussels sprouts, cauliflower,
collards, kale

Aphids, cabbage looper,
imported cabbage worm,
grasshoppers, leafhoppers,
salt-marsh caterpillar mites,
cutworms (climbing), dipterous
leafminer (adult) and Lygus
bugs

Brussels sprouts, cauliflower, collards, kale

Aphids - Apply at 1/4 to 1/2 pt. (0.125 to 0.25 lb. a.i.) in 10-125 gal. water/acre.

Cabbage looper, imported cabbage worm, grasshoppers, leafhoppers, salt-marsh caterpillar, mites, cutworms (climbing), dipterous leafminer (adult) and Lygus bugs - Apply at 1/2 to 1 qt. (0.5 to 1.0 lb. a.i.) in 10-125 gal. water/acre.

Do not treat within 3 days of harvest.

For hard-to-kill insects, including aphids, apply at 1 qt. (1.0 lb. a.i.) in 10-125 gal. water/acre, but do not treat collards and kale within 7 days of harvest; 3 days on Brussels sprouts and cauliflower.

Carrots

Aphids, leafhoppers, Lygus
bugs, mites, cabbage looper,
dipterous leafminer (adult),
cutworm (climbing), and salt-
marsh caterpillar

Carrots

Aphids - Apply at 1/4 to 1/2 pt. (0.125 to 0.25 lb. a.i.) in 10-125 gal. water/acre.

Leafhoppers, Lygus bugs, mites, cabbage looper, dipterous leafminer (adult), cutworm (climbing), and salt-marsh caterpillar - Apply at 1/2 to 1 qt. (0.25 to 0.5 lb. a.i.) in 10-125 gal. water/acre.
Do not treat within 2 days of harvest.

Celery

Aphids, dipterous leafminer
(adult), Lygus bugs, salt-marsh
caterpillar, leafhopper,
cabbage looper and mites

Celery

Aphids - Apply at 1/4 to 1/2 pt. (0.125 to 0.25 lb. a.i.) in 10-125 gal. water/acre.

Dipterous leafminer (adult), Lygus bugs, salt-marsh caterpillar,

leafhopper, cabbage looper and mites - Apply at 1/2 to 1 qt. (0.25 to 0.5 lb. a.i.) in 10-125 gal. water/acre.

Do not treat within 3 days of harvest.

For hard-to-kill insects, including aphids, use 1 qt. (1.0 lb. a.i.) in 10-125 gal/acre, but do not treat within 5 days of harvest.

Corn (field, sweet and popcorn)
for grain only

Aphids

Corn (field, sweet and popcorn) for grain only

Aphids - Apply at 1/4 to 1 pt. (0.125 to 0.5 lb. a.i.) in 10-125 gal. water/acre.

Do not treat within 2 days of harvest.

Cucumbers (outdoor)

Aphids, grasshoppers,
leafhoppers and mites

Cucumbers (outdoor)

Aphids - Apply at 1/4 to 1/2 pt. (0.125 - 0.25 lb. a.i.) in 10-125 gal. water/acre.

Grasshoppers, leafhoppers and mites - Apply at 1/2 to 1 pt. (0.25 - 0.5 lb. a.i.) in 10-125 gal. water/acre.

Do not treat within 2 days of harvest.

Eggplant, peppers

Aphids, grasshoppers,
leafhoppers and mites

Eggplant, peppers

Aphids - Apply at 1/4 to 1/2 pt. (0.125 - 0.25 lb. a.i.) in 10-125 gal. water/acre.

Grasshoppers, leafhoppers and mites - Apply at 1/2 to 1 pt. (0.25 - 0.5 lb. a.i.) in 10-125 gal. water/acre.

Do not treat within 2 days of harvest.

Lettuce

Aphids, corn earworm, cutworms (climbing), dipterous leafminer (adult), cabbage looper, imported cottage worm, grasshoppers, mites, Lygus bugs, salt-marsh caterpillar, false chinch bug and thrips

Lettuce

Aphids - Apply at 1/4 to 1/2 pt. (0.125 - 0.25 lb. a.i.) in 10-125 gal. water/acre.

Corn earworm, cutworms (climbing), dipterous leafminer (adult), cabbage looper, imported cabbage worm, grasshoppers, mites, Lygus bugs, salt-marsh caterpillar, false chinch bug and thrips - Apply at 1/2 to 1 pt. (0.25 - 0.5 lb. a.i.) in 10-125 gal. water/acre.

Do not treat within 2 days of harvest.

For hard-to-kill insects, including aphids, apply at 1 qt. (1 lb. a.i.) in 10-125 gal. water/acre, but do not treat within 4 days of harvest.

Melons (including cantaloupes, honey-dew melons, muskmelons, watermelons)

Aphids, cabbage looper, imported cabbage worm, dipterous leafminer (adult), leafhoppers, Lygus bugs, mites, false chinch bugs, salt-marsh caterpillars, cutworms (climbing), grasshoppers; tobacco budworms on watermelon

Melons (including cantaloupes, honeydew melons, muskmelons, watermelons)

Aphids - Apply at 1/4 to 1/2 pt. (0.125 - 0.25 lb. a.i.) in 10-125 gal. water/acre.

Dipterous leafminer (adult), leafhoppers, Lygus bugs, mites, false chinch bugs, and grasshoppers - Apply at 1/2 to 1 pt. (0.25 - 0.5 lb. a.i.) in 10-125 gal. water/acre.

Cabbage looper, salt-marsh caterpillar, cutworms, (climbing) - Apply at 1/2 to 1 pt. (0.25 - 0.5 lb. a.i.) in 10-125 gal. water/acre (except watermelons).

Rindworms (cabbage looper, cutworms, salt-marsh caterpillars, tobacco budworms) on watermelons - Apply at 1/2 pt. in 10-125 gal. water/acre. Do not treat cantaloupes, honey-dew melons, muskmelons or watermelons within 2 days of harvest.

Okra

Aphids, cutworms (climbing), corn earworms, green stink bug, mites and velvet bean caterpillar

Okra

Aphids - Apply at 1/4 to 1/2 pt. (0.125 - 0.25 lb. a.i.) in 10-125 gal. water/acre.

Cutworms (climbing), corn earworms, green stink bug, mites and velvet bean caterpillar - Apply at 1/2 to 1 pt. (0.25-0.5 lb. a.i.) in 10-125 gal. water/acre.

Do not treat within 2 days of harvest.

Onions (including green onions)

Thrips and cutworms (climbing)

Onions (including green onions)

Thrips and cutworms (climbing) - Apply at 1/2 to 1 pt. (0.25-0.5 lb. a.i.) in 10-125 gal. water/acre.
Do not treat within 2 days of harvest.

Mustard greens, turnip tops

Aphids, cabbage looper,
imported cabbage worm, false
chinch bug, dipterous leafminer
adult, grasshoppers,
leafhoppers and mites

Mustard greens, turnip tops

Aphids - Apply 1/4 to 1/2 pt. (0.125 to 0.25 lb. a.i.) in 10-125 gal. water/acre.

Cabbage looper, imported cabbage worm, false chinch bug, dipterous leafminer adult, grasshoppers, leafhoppers and mites - Apply 1/2 to 1 pt. (0.25 to 0.5 lb. a.i.) in 10-125 gal. water/acre.
Do not treat within 3 days of harvest.

Parsley

Aphids

Parsley

Aphids - Apply 1 to 2 pt. (0.5 - 1.0 lb. a.i.) in a minimum of 40 gal. water/acre. For light infestations use lower dosage of 1 pt. (0.5 lb. a.i.). For hard-to-kill insects and increased populations use 2 pt. (1.0 lb.).

Begin application when insects first appear and repeat as often as necessary to maintain control.

Do not treat within 5 days of harvest for the low dosage or within 8 days of harvest for the high dosage. Do not treat more than 3 times between harvests.

Peas (including vines)

Aphids, grasshoppers,
leafhoppers, mites and cutworms
(climbing)

Peas (including vines)

Aphids - Apply 1/4 to 1/2 pt. (0.125 to 0.25 lb. a.i.) in 10-125 gal. water/acre.

Grasshoppers, leafhoppers, mites and cutworms (climbing) - Apply at 1/2 to 1 pt. (0.25 - 0.5 lb. a.i.) in 10-125 gal. water/acre.
Do not treat within 2 days of harvest.

Potatoes

Aphids, grasshoppers,
leafhoppers, and mites

Potatoes

Aphids - Apply 1/4 to 1/2 pt. (0.125 - 0.25 lb. a.i.) in 10-125 gal. water/acre.

Grasshoppers, leafhoppers and mites - Apply at 1/2 pt. (0.25 lb. a.i.) in 10-125 gal. water/acre.

Do not treat within 2 days of harvest.

Spinach

Aphids, cabbage looper, imported cabbage worm, grasshoppers, leafhoppers, mites, dipterous leafminer (adult), cutworms (climbing), salt-marsh caterpillar and false chinch bug

Spinach

Aphids - Apply at 1/4 to 1/2 pt. (0.125 - 0.25 lb. a.i.) in 10-125 gal. water/acre.

Cabbage looper, imported cabbage worm, grasshoppers, leafhoppers, mites, dipterous leafminer (adult), cutworms (climbing), salt-marsh caterpillar and false chinch bug - Apply at 1/2 to 1 pt. (0.25 - 0.5 lb. a.i.) in 10-125 gal. water/acre.

Do not treat within 4 days of harvest.

For hard-to-kill insects, including aphids, apply at 1 qt. (1 lb. a.i.) in 10-125 gal. water/acre, but do not treat within 7 days of harvest.

Summer squash

Aphids, cabbage looper, dipterous leafminer (adult), leafhoppers, Lygus bugs, mites, false chinch bugs, salt-marsh caterpillar, cutworms (climbing), and grasshoppers

Summer squash

Aphids - Apply at 1/4 to 1/2 pt. (0.125 - 0.25 lb. a.i.) in 10-125 gal. water/acre.

Cabbage looper, dipterous leafminer (adult), leafhoppers, Lygus bugs, mites, false chinch bugs, salt-marsh caterpillar, cutworms (climbing), and grasshoppers - Apply at 1/2 to 1 pt. (0.25 - 0.5 lb. a.i.) in 10-125 gal. water/acre.

Do not treat within 2 days of harvest.

Tomatoes (outdoor)

Aphids, grasshoppers, leafhoppers and mites

Tomatoes (outdoor)

Aphids - Apply at 1/4 to 1/2 pt. (0.125 - 0.25 lb. a.i.) in 10-125 gal. water/acre.

Grasshoppers, leafhoppers and mites - Apply at 1/2 to 1 pt. (0.25 - 0.5 lb. a.i.) in 10-125 gal. water/acre.

Do not treat within 2 days of harvest.

Turnips

Aphids, cabbage looper, imported cabbage worm, grasshoppers, leafhoppers, mites, false chinch bug and dipterous leafminer (adult)

Turnips

Aphids - Apply at 1/4 to 1/2 pt. (0.125 - 0.25 lb. a.i.) in 10-125 gal. water/acre.

Cabbage looper, imported cabbage worm, grasshoppers, leafhoppers, mites, false chinch bug and dipterous leafminer (adult) - Apply at 1/2 to 1 pt. (0.25 - 0.5 lb. a.i.) in 10-125 gal. water/acre.

Do not treat within 3 days of harvest.

GREENHOUSE

Diluted PHOSDRIN 4 EC should be applied in a closed greenhouse (close all doors, windows and ventilators). Lock or barricade all entrances, post warning signs, and take whatever precautions are necessary to prevent unprotected humans and domestic animals from entering the treated area. The operator must wear a full-face mask of a type found adequate for mevinphos (PHOSDRIN Insecticide) protection.

Apply any time the greenhouse ventilators can remain closed for two hours without endangering crops from high temperatures, such as early morning, cloudy days or late afternoons. If application is made during the day, keep the greenhouse closed tightly for at least two hours, then ventilate the enclosure thoroughly for an hour before resuming regular work. If application is made in the late afternoon, the house can be kept closed all night. On the following morning, ventilate for one hour to render the air safe for regular work.

Lettuce (including leaf, bibb and romaine)

Aphids, corn earworms, cutworms (climbing), dipterous leafminer (adult), cabbage looper, imported cabbage worm, grasshoppers, mites, Lygus bugs, salt-marsh caterpillars, false chinch bugs and thrips - Apply 1 1/4 to 2 1/3 pt. (0.625 - 1.155 lb. a.i.) diluted, per 50,000 sq. ft.

Do not treat within 10 days of harvest.

AERIAL APPLICATION

Vegetable and Field Crops

General - Apply at rates indicated, to vegetables and crops as listed above. Use 5-20 gallons water/acre.

Parsley

Aphids

Parsley

Aphids - Apply Phosdrin 4EC, 1 to 2 pt., (0.5 - 1.0 lb. a.i.) in 5-20 gallons water/acre. For light infestations use the lower dosage.

Do not treat within 5 days of harvest.

For hard-to-kill insects and increased populations use the higher dosage.

Do not treat within 8 days of harvest.

Begin application when insects first appear and repeat as often as necessary to maintain control.

Do not treat more than 3 times between harvests.

Fruit and Nut Crops

Grapes

Aphids, leaf folder,
leafhoppers, mites, red-banded
leafroller and Lygus bugs

Grapes

Aphids - Apply at 1/2 to 1 pt. (0.25 - 0.5 lb. a.i.) in 7-25 gal. water/acre.

Do not treat within 4 days of harvest.

Leaf folders, leafhoppers, mites, red-banded leafroller and Lygus bugs-

Apply at 1 to 2 pts. (0.5 - 1.0 lb. a.i.) in 7-25 gal. water/acre.

At the 2 pt. dosage do not treat within 5 days of harvest.

Orchard Crops

General - Apply at rates indicated, to fruit and nut crops listed above.
Use 10-20 gallons water/acre.

PESTICIDE CHARACTERISTICS

Mode of Action

Mevinphos is highly toxic, and may enter the body by ingestion or by dermal contact. It is a cholinesterase inhibitor (EPA 1988, Hayes 1982) which acts directly, as differentiated from others which are not active until they have been altered by chemical or enzymatic action (Doull 1976).

Physical and Chemical Properties

Mevinphos exists in two isomeric forms, the E or cis-crotonate isomer sometimes designated alpha, and the Z or trans-crotonate, designated as the beta isomer. The two differ in NMR spectra, melting point, stability, and toxicity to insects and mammals (Fukuto et al 1961, Hayes 1982). The E-isomer is about 100 times more toxic to both insects and mammals than the Z-isomer (Cremlyn 1978). The empirical formula is $C_7H_{13}O_6P$, with a molecular weight of 224.16. Preparation is described by A. R. Stiles (U. S. pat. 2,685,552 to Shell 1954). Technical mevinphos is a pale yellow to orange liquid with a mild odor or without odor. The boiling point ranges from 99 - 103°C at 0.03 mm Hg. The E-isomer has a melting point of 21°C, the Z- of 6.9°C. The density is 1.24 at 16°C, with densities of the two isomers 1.2345 and 1.245, respectively. The vapor pressure is 0.003 mm Hg at 21°C. Most commercial products contain about 60% E- and 30% Z-isomer. The technical material is miscible with water, alcohols, ketones, chlorinated hydrocarbons, and aromatic hydrocarbons, but is only slightly soluble in aliphatic hydrocarbons. It is stable at ordinary temperatures. Mevinphos is hydrolyzed in water with a half-life of 1.4 hours at pH 11, 3 days at pH 9, 35 days at pH 7, and 120 days at pH 6. Lime sulphur results in rapid decomposition (EPA 1988, Hayes 1982). The cis-isomer is hydrolyzed faster than the trans, at pH 11 and 28°C the half-life is 1.8 and 3.0 hr respectively (Gatterdam et al 1959). Exposure to UV will increase the proportion of the trans-isomer (Casida 1955).

Formulations Marketed

Mevinphos has been marketed in the past as a dust, emulsifiable concentrate, soluble concentrate/liquid, and as a ready to use formulation. It is now marketed as an emulsifiable concentrate at 4 lb. per gallon, or as a solution in isopropyl alcohol at 4 lb. per gallon (EPA 1988).

Physiological Activity in Soil or Plant and Animal Tissue

Mevinphos is absorbed by plant roots, translocated readily to leaves and growing shoots, and degraded rapidly (EPA 1988). Mevinphos as a systemic insecticide is translocated unchanged. Metabolism does not increase its toxicity (Eto 1974). The major residues found in plants are the E- and Z-isomers of mevinphos and dimethyl phosphate. Mevinphos acid is a minor metabolite in plants, which is converted to desmethyl mevinphos acid (EPA 1988). In cows dimethyl phosphate is the primary metabolite in the urine, other hydrolysis products are found in the feces (Matsumura 1985). The E-

isomer is readily degraded by the mouse liver glutathione S-alkyltransferase (Morello et al 1968). The Z-isomer is not demethylated, but is degraded by esterases splitting the vinyl ester linkage (Spencer 1972).

A moderate teratogenic effect was produced when mevinphos was injected into hen eggs before or during the early period of incubation (Eto 1974).

The acute oral LD₅₀ for rats is 4 to 7 mg/kg (Eto 1974). In a study in which rats were fed 110 ppm mevinphos, all died within 3 weeks. At 50 ppm for 60 days, rats showed reduced growth, slight tremor, and brain cholinesterase 20% of normal. At 6.3 ppm, rats showed slight tremor and brain cholinesterase 90% of normal (Kodama et al 1954). Dogs fed at a dietary level of 5 ppm (ca. 0.1 mg/kg/day) for 14 weeks showed erythrocyte and plasma cholinesterase levels reduced to 70 and 90% of controls, respectively (Cleveland & Treon 1961). Animal metabolism data indicate that twelve hours after dosing 57 to 65% of the ³²P-residue was excreted, 45-50% in the urine and 12-15% in the feces, and only mevinphos hydrolysis products were present (EPA 1988).

In an experiment using human volunteers, a dose of 2.5 mg/man/day (ca. 0.036 mg/kg/day) produced a maximal decrease of 25% in red blood cell (RBC) cholinesterase by the 27th day, but did not affect plasma cholinesterase activity or result in any other symptoms. Dosages of 1.5 and 2.0 also reduced RBC cholinesterase, whereas a dose of 1.0 for 30 days produced no detectable effect. After dosing was stopped, return of RBC cholinesterase to normal was determined entirely by the rate of replacement of red cells (Rider et al 1972, 1975). Accidental poisoning may result in the rapid appearance of symptoms. In more than one instance this has involved the impairment of judgement (Hayes 1982), which is not explained by studies in animals that indicate that penetration of mevinphos into the brain is slow and limited, and that brain cholinesterase is less susceptible than the enzymes derived from plasma and red cells (Sharma et al 1973). Other accident reports have shown that symptoms may be delayed until 48 hours after last exposure (Roche et al 1961).

METHODOLOGY OF THE BENEFITS ASSESSMENT STUDY

A questionnaire was developed to assess the benefits of mevinphos (see APPENDIX I). This questionnaire was designed to elicit key information regarding the present use of mevinphos. Each state was requested to complete a questionnaire for each crop/site for which the pesticide was used in significant quantities. Data were collected relating to the averages of production, costs, and use of mevinphos for the crop years 1982-1986. Questionnaires were mailed to each State Pesticide Liaison Representative on July 19, 1988. Questionnaires were sent to each of the 50 states plus territories of the U.S. (Virgin Islands, Guam, Puerto Rico) as well as Washington, D.C.

Responses were received from all 50 states, as well as the Virgin Islands; Guam; Washington, DC; and Puerto Rico. Mevinphos had important commercial use on 33 crops in 25 states. Survey responses by commodity and U. S. production by commodity are given in APPENDIX II.

Reporting states used a variety of methods to complete the questionnaires. In some states (notably Florida), all responses were formulated by a single individual working under the supervision of the State Pesticide Liaison Representative. At the other end of the spectrum, some State Liaison Representatives sent the questionnaire to extension agents and/or industry field men in key production areas. In these cases, a single individual may have responded for one or several sites. Other states fell in between these two extremes.

Rules were established to deal with missing data. In general, production data (yields, acres, prices) were used for the affected crops as reported by respondents. Major adjustments to account for missing or inaccurate data were as follows:

- a) All production data as given by the respondents were used when it appeared to be reasonable. Internal checks were run by converting all production data to hundredweight and computing price per hundredweight and revenue per acre. In this manner all "unreasonable" data were checked against other data sources such as the National Agricultural Statistics Service (NASS) and the Economic Research Service.
- b) When missing data were encountered, we used NASS data where available for crop production, acreage, prices, etc. Missing data on yield changes and alternative control costs were taken from similar sites in the same region.
- c) Some sites were eliminated where the crop or mevinphos use was very minor.
- d) When respondents reported ranges, we took the mid-point of the range.

All survey adjustments are reported in Appendix III.

Survey responses are summarized in Tables 2 and 3. The 33 crops on which mevinphos is used in significant quantities are indicated in Table 2. A total of 130 questionnaires were used in the analysis. These crops account for 2.9 million acres in production. Mevinphos was used on about 667,000 acres, or 23 percent of the total acreage of these crops. The largest use in terms of acreage was (1) lettuce (209,500 acres or 90 percent of lettuce acreage); (2) alfalfa seed (92,800 acres or 69 percent of alfalfa seed acreage); and (3) broccoli (61,892 acres or 55 percent of broccoli acreage).

The crops on which mevinphos is used accounted for an aggregate value of production of \$4.8 billion (Table 3).

Table 2. Survey responses: States reporting, acreage, and acreage treated with mevinphos (average annual), 1982-86.

Crop	States reporting	Acreage reported		Acres treated (percent)
		in production	treated	
Alfalfa seed	CA,OR,WA	133,862	92,800	69
Apple	IL,NJ,NY	83,680	4,740	6
Broccoli	AL,CA,FL,NJ OR,TX	112,622	61,892	55
Brussels sprouts	CA	3,094	3,000	97
Cabbage	AL,CA,DE,FL, GA,IL,NC,NJ,NY SC,TX,UT,WI	83,158	46,310	56
Pea, southern	TN	300	250	83
Carrot	FL,OR,WI	15,060	3,535	23
Cauliflower	CA,FL,NJ,OR,TX	51,337	34,400	67
Celery	CA,FL,MI	30,973	15,743	51
Greens	AL,AZ,FL,GA,IL, LA,MD,NJ,NY,NC, LA,OK,TN,TX,VA	33,142	24,985	75
Corn, field	CO,FL	706,200	820	--
Corn, sweet	FL	67,020	12,712	19
Cucumber	FL,GA,NJ,TX,OR	33,280	10,126	30
Eggplant	FL,NJ	3,396	401	12
Grape	CA	676,700	11,419	2
Muskmelons	CA,FL,GA,NC, NJ,TX	55,230	38,463	70
Lettuce	AZ,CA,CO,FL,MI, NJ,NY,TX,WI	233,496	209,500	90
Onion (all)	CA,FL,OR,WI	10,110	5,550	55
Parsley	FL	910	400	44
Pea, english	WI	88,140	500	1
Pepper	FL,NJ,TN,TX	38,100	17,769	47
Potato	FL,ID	32,900	658	2
Sorghum	FL	45,045	1,700	4
Squash	FL,GA	25,360	4,650	18
Strawberry	FL,NJ,CA	19,102	8,177	43
Tomato	AR,FL,NJ,TX	58,370	6,922	12
Turnip	FL	1,199	200	17
Artichoke	CA	10,417	10,000	96
Bean, lima	FL	242	23	10
Bean, snap	AR,FL,NC,TN,WI	142,497	10,169	7
Okra	FL	2,440	1,855	76
Spinach	MD,NJ,NY,FL,VA, OK,TX,CA	22,299	13,910	62
Watermelon	NJ,FL,GA,NC	109,610	13,145	12
Total		2,929,291	666,723	23

Table 3. Survey responses: Acreage, production, value (average annual), 1982-1986.

Crop	Acreage	Production (1,000 cwt.)	Value of production		
			Total (1,000 dollars)	Average dollars (per cwt.)	Average dollars (per acre)
Alfalfa seed	133,862	1,085	105,390	97.13	787
Apple	83,680	12,585	135,530	10.77	1,620
Broccoli	112,622	10,570	260,034	24.60	2,309
Brussels sprouts	3,094	294	9,226	31.41	2,982
Cabbage	83,518	24,601	184,248	7.49	2,216
Pea, southern	300	10	185	18.97	617
Carrot	15,060	2,699	20,865	7.73	1,385
Cauliflower	51,337	5,380	134,364	24.97	2,617
Celery	30,973	10,923	138,292	12.66	4,465
Greens	33,142	4,874	110,569	22.68	3,336
Corn, field	706,200	66,735	296,925	4.45	420
Corn, sweet	67,020	6,997	83,894	11.99	1,252
Cucumber	33,280	4,717	64,163	13.60	1,928
Eggplant	3,396	706	11,289	15.98	3,324
Grape	676,700	49,315	1,043,366	21.16	1,542
Muskmelons	55,230	7,967	105,199	13.20	1,905
Lettuce	233,496	67,218	783,094	11.65	3,354
Onion	10,110	5,144	44,414	8.63	4,393
Parsley	910	142	4,162	29.31	4,574
Pea, english	88,140	2,643	25,737	9.74	292
Pepper	38,100	4,201	109,272	26.01	2,868
Potato	32,900	7,486	66,227	8.85	2,013
Sorghum	45,045	8,394	10,969	1.31	244
Squash	25,360	2,946	44,098	14.97	1,739
Strawberry	19,102	7,555	332,503	44.01	17,407
Tomato	58,370	35,021	396,046	11.31	6,785
Turnip	1,199	---	---	---	---
Artichoke	10,417	795	30,838	38.77	2,960
Bean, lima	242	8	513	64.13	2,120
Bean, snap	142,497	6,656	79,869	12.00	560
Okra	2,440	83	2,829	34.08	1,159
Spinach	22,299	2,857	54,761	19.17	2,456
Watermelon	109,610	12,604	78,516	6.23	716
Total	2,929,291		4,767,386		1,626

EXTENT OF USE

Mevinphos was registered for use on 109 different sites as of May 9, 1988 (Table 1). The various use sites include small fruits, citrus, tree fruits and nuts, various vine crops, vegetables, field crops, ornamental and sewage systems. During the 1982-86 use period, 666,723 acres (Table 2) of 33 various agricultural commodities were treated with a total of 756,815 pounds of mevinphos active ingredient (Table 34).

The greatest use was on lettuce (241,748 lbs a.i.) followed by uses on strawberry, broccoli, greens, cabbage, alfalfa seed, and cauliflower. The amount of mevinphos used on these 7 commodities was 632,751 lbs a.i. or 83.6 % of the total combined use on all other crops that reported use in the United States. A significant percentage of the total United States acreage of the following crops was treated with mevinphos: alfalfa seed (69%), broccoli (55%), Brussels sprouts (97%), cabbage (56%), southern pea (83%), cauliflower (67%), celery (51%), greens (75%), cucumber (30%), muskmelons (70%), lettuce (90%), onions (55%), parsley (44%), pepper (47%), strawberry (43%), artichoke (96%), okra (70%), spinach (62%), and watermelon (23%) (Table 2).

Of the total 756,815 lbs a.i. of mevinphos used in the U.S., the greatest amount was used on vegetables (594,777 lbs), followed by fruits (103,977 lbs), and field crops (58,061 lbs), (Table 34).

Geographic regions in the U.S. that reported the greatest mevinphos usage were the West (445,400 lbs a.i.) and the South (107,035 lbs a.i.). This probably represents treatments on strawberry and vegetable crops.

Until alternative materials and methods are developed that are equal to or better than mevinphos and have a short pre-harvest interval, the foreseeable use patterns will probably stay about the same. That is, a significant percent of the strawberry and certain vegetable acreage will continue to use mevinphos, whereas most of the field and fruit crops are currently using safer and more cost-effective pesticides and will probably continue to do so.

The average annual use of mevinphos, 756,815 pounds active ingredient, which was estimated from the surveys is above the indicated usage in the United States which has averaged 500,000 to 750,000 pounds active ingredient per year. This could result from several factors. Entomologists, extension specialists, and industry personnel who responded to the survey may have based their estimates of use on events or years which were extremes rather than averages for the period 1982-1986. Furthermore, their contacts may tend to be based on more progressive grower or growers who are more aware of particular insect problems. Apparently the estimate is about 21 per cent above actual use. The committee believes that our estimate is defensible given the inevitable data limitations, time and funding available for this study.

SPECIFIC USE ANALYSIS

FRUIT AND NUT CROPS

The results of our survey questionnaire suggest that mevinphos is currently being used only on apples, grapes, and strawberries in the United States, although it is registered for use on nine other fruit/small fruit/nut crops. It was reported as being used against leafrollers (redbanded, obliquebanded and variegated), mites and fruitworms (green and sparganothis), aphids, Lygus bugs and codling moth on apples; grape leafhopper, mites and grape leafroller on grape and aphids, mites and strawberry leafroller on strawberry.

Leafrollers. Damage to apple is caused by the larval stage. Larvae are indiscriminate feeders with most serious injury occurring around petal fall and shortly after when developing fruit is damaged. Fruit damaged at this stage usually drops prematurely, but some remains on the tree and develops deep corky scars and indentations by harvest. Although the larvae are primarily foliage feeders, they often feed on the surface of the fruit during the season (Reissig 1980, Metcalf et al. 1962). If not controlled, 80-90% of the crop may be damaged in years of abundance (Metcalf et al. 1962).

Mites. Two species, the European red mite and the twospotted spider mite, are pests of apples. All motile stages of both species feed on individual leaf cells removing the contents, including the chlorophyll. The injury results in off-color foliage. The leaf efficiency and productivity is reduced significantly. Heavy feeding in June and July reduces tree growth and yield and can drastically affect fruit bud formation reducing yield the following year (Metcalf et al. 1962, Lienk 1980).

Fruitworms. This complex is an early season pest of apples. Most damage is observed within 3 - 4 weeks after petal fall. The larval stage feed on the developing fruit. Often, the whole side of an apple is consumed. A single larva will severely damage several fruit on a branch. Often the fruit will remain on the tree. At harvest it is worthless (Metcalf et al. 1962).

Aphids. The major aphid pest of apple is the rosy apple aphid. It is an early season pest found on the trees from bud-break until late spring. Feeding on the foliage causes severe distortion of the leaves. Feeding on the leaves of fruit clusters often results in bunching, stunting, and malformation of the fruit. This becomes noticeable as the fruit develops and renders it unmarketable. Honeydew produced by the aphids provides a media for sooty mold fungus growth. This can affect fruit finish. Populations levels fluctuate and can cause significant fruit loss if not controlled (Wieres and Leeper 1980). The apple aphid and the spirea aphid also are pests. Both of these species remain on the tree throughout the season. Their feeding does not cause leaf curling or fruit distortion. However, the copious amounts of honeydew produced by these aphids provides a media for the growth of sooty mold with the same results as described for the rosy apple aphid (Metcalf et al. 1962).

Lygus bugs. These insects all have piercing-sucking type mouthparts which are inserted into the plant material. A toxic saliva is injected during the feeding process which kills the plant cells in the immediate area. On apple, this may cause abortion of the developing fruit bud or, if shortly after bloom, it causes a deeply depressed area in the fruit. Numbers can vary considerably from year-to-year, but considerable damage occurs if no controls are applied at pink and petal fall (Metcalf et al. 1962).

Codling moth. The larval stage enters the side or blossom end of the apple fruit and tunnels in to the core. The seeds and core are eaten by the larvae rendering the fruit unsuitable for human consumption. Small larvae may partially enter the fruit after pesticide treatment and produce "stings" which reduce the fruit grade. If left uncontrolled, 20 - 90% of the fruit may be infested by this pest (Metcalf et al. 1962).

Grape leafhopper. Both adults and nymphs feed on grape foliage and puncture individual cells and remove the contents. Feeding punctures leave a white spot. This appears on the foliage and fruit. Heavy feeding on the leaves reduces photosynthetic activity. Vines can withstand 20% defoliation a month after fruit set without adverse yield affect. Excessive fruit spotting on table grapes can lead to economic loss before any effect on soluble solids or yield is noted. Adults and nymphs are annoying to pickers when populations are high (Jensen and Flaherty 1981a). In some instances, extremely high populations may reduce the quantity of fruit harvested by as much as 30% (Metcalf et al. 1962).

Spider mites. The Pacific spider mite is the primary pest species while the Willamette spider mite occasionally causes problems and the twospotted spider mite is only occasionally found on grapes. Both the Pacific and Willamette mites cause chlorosis of the leaves while high densities of the Pacific mite can cause shoot tip distortion.

Pacific mites can affect fruit quality or maturity due to the loss of productive leaf surface. The following year vine growth and crop may also be affected. Occasionally, heavy populations of Pacific mites have killed grape vines.

Willamette mites seldom produce economic damage to grapevines. Spotty damage may occur in the spring when high numbers overwinter (Flaherty et al. 1981).

Grape leafroller. Injury to grapes is caused by the larval stage. The primary damage is by rolling and feeding on the leaves which reduces the photosynthetic function. Extreme population densities may feed on fruit, but economic damage usually occurs only with massive, late season infestations. In table grapes, defoliation of the top of the vine can result in sunburned, discolored fruit (Jensen and Flaherty 1981b).

Aphids. Damage to grapes occurred in the 1950s on an occasional basis. The species was not determined. At this time no economic damage is attributed to aphids (Jensen and Kido 1981).

Aphids. Several species of aphids are potential problems on strawberries. Direct damage is caused by the removal of plant juices during feeding. When present in large numbers they weaken the plants. The large amounts of honeydew excreted during feeding supports the growth of sooty mold. When this occurs on the fruit, quality is lost. The major problem with aphids on strawberries is their role in transmitting virus diseases to the plants. When aphids are not controlled and virus sources are nearby, nearly 100% infection can occur in as little as two weeks but generally occurs within two or three years (Mass 1984).

Spider mites. Mites remove the contents of strawberry leaf cells during feeding. This results in dry areas in the leaves and when populations are very high may result in stunting and die back of the plants. This can affect the current year's crop and significantly reduce the following year's yield. Plant death may occur when damage is combined with drought stress (Goulart et al. 1988, Mass 1984).

Strawberry leafrollers. Damage to strawberry is caused by larval feeding on the foliage. Continuous feeding causes leaves to turn brown and die. One species, the omnivorous leaf tier, is a pest in California. A high proportion of the larvae feed on the berries and may remain inside. Even low populations cause serious damage by contaminating the processed fruit (Mass 1984).

Lygus bugs. Adults suck sap from strawberry buds, blossoms, newly formed green fruits, and tender growing leaf tips. While feeding a toxic saliva is injected into the plant tissue. When young fruit are attacked the toxic saliva kills some of the tissue causing the berry to develop unevenly. This produces a fruit with a knobby button-like appearance (Goulart et al. 1988, Mass 1984).

Apples/Peaches/Pears/Plums

Controls are recommended primarily during the early part of the season for leafrollers, fruitworms, aphids, Lygus bugs, and codling moth. Exceptions are for mites and obliquebanded leafrollers which often require treatment during the middle of the season. Numerous sampling schedules and treatment thresholds are suggested. The treatment thresholds vary, depending on the pest and the time of year. Rosy apple aphids are scouted for at pink-bud. New York recommends treatment if 5% of the leaves in fruit clusters have rosy apple aphids (Agnello et al. 1988) while North Carolina suggests a threshold of 10% of the terminals infested after Second Cover Spray (Walgenbach 1988). The threshold for apple aphid is 50% of the terminals infested in North Carolina (Walgenbach 1988).

All states surveyed recommend a pesticide application at petal fall to reduce insect pest populations. Thresholds for leafrollers are 3% of the leaf clusters infested during bloom in New York (Agnello et al. 1988) and five larvae or fresh feeding scars observed on 300 fruit in every 10 acres (Walgenbach 1988). In New York, if the threshold during bloom is reached, the petal fall insecticide must be selected to provide control of leafrollers

as well as the other targeted pests. In early July, the New York threshold is 12% of the leaf and fruit clusters infested (Agnello et al. 1988).

New York uses a variable threshold for mites depending on the time of year. Through June 15 the threshold is 2.5 mites/leaf; June 16-30, 5 mites/leaf; July 1-15, 7.5 mites/leaf; July 16 - August 1, 10 mites/leaf (Agnello et al. 1988). North Carolina used a threshold of 85% of the leaves infested with one or more mites (Walgenbach 1988).

Other pests such as Lygus bugs and codling moth are generally controlled with the petal fall and first cover sprays recommended by nearly all apple producing states. Codling moth can be sampled with pheromone traps. North Carolina uses a threshold of four moths caught each week for two consecutive weeks if one trap/3 acres is used or ten moths caught each week for two consecutive weeks if there is one trap/10 acres (Walgenbach 1988).

In general, mevinphos was used on only 5.7% of the acreage in the three states reporting its use on apples. The alternatives to mevinphos, rates, and number of acres to be treated are given in Table 4. In all three states, single applications would be used for all alternatives except chlorpyrifos which would require 1.5 applications on the average.

Both Illinois and New Jersey did not feel that there was a need for mevinphos or that there was any yield or quality reduction due to the use of the alternatives. The only advantage of mevinphos was that it could be used close to harvest. The alternatives were all safer to use.

New York felt that the alternatives were less hazardous, however all had longer preharvest intervals, were more expensive and with the possible exception of methomyl were less effective against the leafroller complex. Also, leafrollers are becoming more tolerant to the alternatives. Loss of mevinphos would increase losses due to leafrollers by 1-2% in years of heavy leafroller populations. For the three states using mevinphos on apples there is no economic gain from its use.

There were no reported uses of mevinphos on peaches, pears or plums.

Cherries

There were no reported uses of mevinphos on cherries.

Citrus: Oranges/lemons/grapefruit

There were no reported uses of mevinphos on citrus.

Grapes

California, the only state reporting use of mevinphos on grapes, treats 1.7 percent of its grape acreage with mevinphos annually. This indicates

that the alternatives to mevinphos, listed in Table 5, are the primary pesticides used for arthropod management.

Treatment thresholds for Grape Leafhopper vary with the grape variety, its end use, and the time of year. The following is an example using Thompson Seedless (Jensen and Flaherty 1981a):

<u>Use</u>	<u>1st Brood</u>	<u>2nd Brood</u>
raisins wine	20 nymphs per leaf	10 - 15 nymphs/leaf (Do not permit >20% leafloss)
		or
		600 nymphal days from beginning of 2nd brood until harvest
Table grapes	10 - 15 nymphs/leaf	Early varieties: 10 nymphs/leaf
		Midseason varieties: 5 - 10 nymphs/leaf
		Late varieties: 5 - 8 nymphs/leaf
		or
		300 nymphal days from beginning of the second brood until harvest

The need for treating mites depends on the vigor of the vineyard, the soil type and the species of mite present. Average to strong vineyards can tolerate fairly heavy populations. Poor soils result in weaker vines. Dusty conditions promote outbreaks of Pacific spider mites. Spot infestations require treatment only when beneficials are absent. General infestations with moderate or heavy Pacific mite populations and very few predators require immediate treatment. Low mite populations with few predators indicate delayed treatment. Moderate to heavy mite populations with predaceous mites frequent to numerous should be monitored (Flaherty et al. 1981).

Grape leaffolder management is not well defined. Vineyards should be monitored for young larvae and for leaf rolling. Wine and raisin grapes can withstand more defoliation than table grapes. The first brood rarely requires treatment. Second and third broods should be treated as the bulk of the brood begins to make new leaf rolls (Jensen and Flaherty 1981b).

Aphids are not a problem at this time in California vineyards. Controls

Table 4. Use of Mevinphos on Apples (average annual), 1982-1986.

State	Acres Treated	Pests	Rate (lbs. a.i./A)	No. Appl.	Alternatives	Rate lbs. a.i./A (No. A treated)	Limitations
Illinois	1491	leafrollers mites codling moth	0.2	1.5	azinphosmethyl phosmet	0.625 (795) 0.625 (693)	longer PHI
New Jersey	37	leafrollers mites aphids Lygus bugs	0.75	1-2	dimethoate methomyl endosulfan	1.5 0.34 1.5	
New York	3212	leafrollers fruitworms	1.0	1	chlorpyrifos endosulfan fenvalerate methomyl methyl parathion (Pennacap-M) permethrin	1.5 (642) 2 (160) 0.2 (482) 0.9 (1124) 1 (642) 0.4 (160)	longer PHI (all alternatives) more expensive less effective against leafrollers

Table 5. Use of Mevinphos on Grapes (average annual), 1982-1986.

State	Acres Treated	Pests	Rate (lbs. a.i./A)	No. Appl.	Alternatives	Limitations
California	11,504	grape leafhopper	0.25-0.5	1-2	carbaryl parathion endosulfan methomyl dimethoate	disruptive to mite predators (except parathion and endosulfan)
		mites			dicofol propargite	
		grape leaf folder			carbaryl Bt parathion	disruptive to mite predators

are not warranted or recommended.

Mevinphos is very disruptive to the beneficial predators that aids in control of mites, and leafhoppers. Several of the alternatives used to control leafhoppers and leafhoppers are also disruptive to the mite predators. Bacillus thuringiensis, parathion, endosulfan, dicofol and propargite are considered relatively easy on the mite predators.

The primary advantage of mevinphos is that it can be used close to harvest for late season pest outbreaks. A 6% yield benefit is realized when mevinphos is used. However, the costs of the alternatives are \$8.38/acre less. In the aggregate, however, there is negative economic benefit to the use of mevinphos.

Raspberries

There were no reported uses of mevinphos on raspberries.

Strawberries

Use of mevinphos on strawberries was heavier than use on apples and grapes. Florida apparently relies completely on mevinphos for arthropod management. California treats 22.5% of its crop with mevinphos while New Jersey treats only 2.8% of its crop with mevinphos.

Alternatives listed by New Jersey include azinphosmethyl, parathion, carbaryl and endosulfan for aphids; dicofol and oxydemeton-methyl for mites; and diazinon, azinphosmethyl and carbaryl for leafrollers. Each would be used once a year. Rates and acreages treated with alternatives are given in Table 6.

All of the alternatives have longer preharvest intervals but are safer to use. New Jersey did not have any yield benefit from the use of mevinphos while California experienced a 60% yield benefit and Florida a 15% yield benefit. Control costs were \$152.99/acre more for the alternatives in California, \$67.74/acre less for the alternatives in Florida, and the same in New Jersey.

In summary, mevinphos appears to be needed for strawberry production.

Walnuts

There were no reported uses of mevinphos on walnuts.

Table 6. Use of Mevinphos on Strawberries (average annual), 1982-1986.

State	Acres Treated	Pests	Rate (lbs. a.i./A)	No. Appl.	Alternatives	Rate lbs. a.i./A (No. A treated)	Limitations
California	3002		0.25-0.5	3			
Florida	5160		1.0	14-21			
New Jersey	17	aphids	0.5	1	azinphosmethyl ethyl parathion carbaryl endosulfan	0.5 (1.7) 0.5 (0.9) 1.0 (13) 0.99 (1.7)	longer PHI (all alternatives)
		mites			dicofol oxydemeton-methyl	0.7 0.75	
		leafrollers			diazinon azinphosmethyl carbaryl	0.5 (7) 0.5 (7) 1.0 (8.5)	

FIELD CROPS

Alfalfa for Hay

Mevinphos (Phosdrin 4EC or Phosdrin IPA4) may be applied to alfalfa for control of aphids and alfalfa caterpillar at a rate of 1/4 to 1/2 pint per acre; for control of grasshoppers, leafhoppers, cutworms (climbing), mites, alfalfa weevil larvae, and Lygus bugs use 1/2 to 1 pint per acre. Do not treat within two days of harvest. It can be used in conventional hydraulic sprayers, low-volume ground applicators, or airplane sprayers.

Alfalfa grown for hay is regularly subject to economic injury caused by several of the insects for which mevinphos is registered, such as alfalfa weevil larvae, leafhoppers, and aphids. Others, such as the alfalfa caterpillar, grasshoppers, and cutworms occasionally cause damage. Losses ranging up to 100% for a given cutting have been observed, although losses are usually much less. Losses from the alfalfa weevil and from leafhoppers are generally restricted to only one or two harvests per year. In some areas hymenopterous parasites and a fungal disease have reduced the alfalfa weevil to sporadic pest status.

Most recommendations combine thresholds for either chemical application or early harvest, with resistant cultivars for spotted alfalfa aphid or winter harvest and parasite conservation for the alfalfa weevil. There are a number of insecticides available.

Only four states reported using mevinphos on alfalfa for hay (Table 7). Based upon their reports, ca. 54,200 acres of alfalfa are treated, this constitutes ca 0.2% of the average 26,078,000 acres of alfalfa harvested for hay 1986-88 (Va. Agr. Stat. Serv. 1988). In the two states which reported the acreage of alfalfa grown in their states, the per cent of the crop treated with mevinphos was <1% in Pennsylvania, and <0.1% in Nevada. There is obvious variation in use from year to year depending on the pest densities. There is usually only one treatment per year except in Nevada where there are usually two. The rate of application of mevinphos varies from 0.125 to 0.5 pounds actual per acre.

Mevinphos is most likely to be applied to alfalfa for hay either just prior to harvest to avoid late pest injury, or when there are flowering weeds in the field which are being visited by bees when controls must be applied to control a serious pest. A late evening treatment with a short residual insecticide such as mevinphos could reduce pest injury without killing many bees (similar to the situation on alfalfa grown for seed). However, on alfalfa grown for hay, neither situation is necessary. Advancing the harvest date is almost always preferable to application of a spray just before harvest. Regarding the second situation, weeds in alfalfa should be controlled as in any other crop. If weeds are present, and a spray is necessary, the beekeeper should be notified to move or protect his bees. Bees may also be attracted to the honeydew produced by the blue alfalfa aphid or the spotted alfalfa aphid.

Table 7. Use of Mevinphos on Alfalfa for Hay (average annual), 1982-1986.

State	Acres Treated	Pests	Alternatives	Rate (lbs a.i/A)	Applic. Method	Limitations ²
Calif.	45267 est.	Aphids Loopers Armyworm Cutworm G-hoppers Mites	MSR ¹ trichlorfon	.25-.5	Gr & Air	MSR- 2/day PHI Trichlorfon- acceptable substitute
Nevada	225	Aphids <u>Lygus</u> Alf.Weev. G-hoppers		.125-.5	Gr & Air	MSR- 2/day PHI Trichlorfon- acceptable substitute
Oregon	725	Aphids		.25-.5	Gr & Air	MSR- 2/day PHI Trichlorfon- acceptable substitute
Penn	8000 est.	Alf.Weev.	methomyl carbofuran chlorpyrifos dimethoate malathion azinphosmethyl methoxychlor	.25-.5	Gr & Air	Malathion- less effective others >2 days PHI

¹MSR = Meta-Systox R, oxydemetonmethyl

²PHI = Preharvest interval

The reports by Stoltz and Baird (1986) and Peairs (1986) show that we now have a number of insecticides which are not only more effective than mevinphos, but have a lower mammalian toxicity. Their data also cast doubt on the idea that mevinphos residues disappear very quickly in the field, the control shown by their 3 day counts for aphids and 7 day counts for webworms in the mevinphos (Phosdrin) treatments were not markedly different from earlier counts.

Mevinphos is applied both by ground and airplane applicators. Thus, there may be spray drift onto nearby crops or noncrop areas. This represents a real danger to man, domestic animals, and wildlife which are present during or shortly after an application. The short residual effect of mevinphos reduces the hazard to individuals or animals which enter an area a day or two after it has been treated, as well as reducing concerns over toxic residues on nontarget areas.

The insecticides which have been suggested as alternatives by our survey form respondents (Table 8), are all less toxic and have a greater residual longevity than mevinphos. The alternatives are quite variable in their toxicity. We do not have information regarding their effectiveness under various field conditions and for individual pest species.

Alfalfa-seed production

Mevinphos is used in alfalfa seed production to control aphids, alfalfa caterpillar, grasshoppers, leafhoppers, cutworms (climbing), mites, alfalfa weevil larvae, and Lygus bugs. Rates for the first two cited pests vary from 0.125-0.25 lb a.i. in 10-125 gal. water/acre. For other cited pests, registered rates range from 0.25-0.50 lb a.i. in 10-125 gal. water/acre. See Table 9.

A major benefit of mevinphos stems from its short pre-harvest interval (PHI) of 2 days. Major hazards include toxicity to non-target organisms, pollinators, beneficial insects, and high user (man) toxicity. (See below).

Production of alfalfa seed is a highly specialized and management intensive process. Management components include three primary facets: 1) control of detrimental insects; 2) the supply of effective pollinators (various species of bees); and 3) irrigation water at time of bloom.

The short residual organophosphates, oxydemeton-methyl, naled, mevinphos and trichlorfon are presently the cornerstones of the insecticide component of pest control - especially in the Pacific Northwest production areas. These chemicals (and mevinphos in particular) provide short residual control of primary and certain secondary pests during the flowering period when pollinators are active. University and industry personnel who deal with this important commodity feel that the loss of these short-residual chemicals would devastate alfalfa pest management systems and would result in seed fields receiving 8-10 applications of other insecticides per year.

Table 8. A Comparison of Toxicities Between Mevinphos and Other Insecticides Recommended as Alternatives on Alfalfa for Hay.

Insecticide	Rat acute oral LD ₅₀	Rat acute dermal LD ₅₀
Mevinphos	3-7	3-90
Azinphosmethyl	11-16	220
Carbofuran	8-14	>10200 (Rb) ¹
Chlorpyrifos	82-245	202
Dimethoate	28-500	>150-<1150
Malathion	885-2800	4000->4444
Methomyl	17-24	>5000 (Rb) ¹
Methoxychlor	5000-7000	>2820->6000
Naled	250-430	800
Oxydemetonmethyl	47-75	100-250
Trichlorfon	450-630	>2000

¹Acute dermal toxicity for rabbits.

Data from: Larson, L. L., E. E. Kenaga and R. W. Morgan. 1985. Commercial and experimental organic insecticides. Entomol. Soc. Amer. 105 p.

Table 9. Use of Mevinphos on Alfalfa for Seed (average annual), 1982-1986.

State	Acres Treated	Pests	Alternatives	Rate (lbs. a.i./A)	Applic. method	Limitations ^{3/}
Oregon plus Idaho	2,500	aphids ^{1/} <u>Lygus</u> ^{2/}	dimethoate oxydemeton-methyl trichlorfon parathion malathion diazinon carbofuran	.125-.50	ground or air	Trichlorfon- acceptable substitute malathion- less effective others >2 days PHI
Nevada	1,037	aphids <u>Lygus</u>	same as for Oregon	.25-.50	ground	Trichlorfon- acceptable substitute malathion- less effective others >2 days PHI
Washington	6,000	aphids <u>Lygus</u>	same as for Oregon	.25-.50	ground or air	Trichlorfon- acceptable substitute malathion- less effective others >2 days PHI

^{1/} 3 aphid spp. - pea aphid, alfalfa aphid and blue aphid
^{2/} Lygus spp. L. elisus and L. hesperus
^{3/} PHI = Preharvest interval

The primary pest for which mevinphos is applied is late season Lygus bug control - a primary seed pest. Users report good control of grasshoppers, control of alfalfa weevil, and some use to suppress seed chalcid. Mevinphos has very severe impact on some beneficials and has a high applicator hazard, but is relatively safe on pollinating bees with only a 5-hr or less residual activity. With timing of mevinphos applications after sunset, the hazard to bees is minimized.

A synopsis of on-farm usage of mevinphos is supplied in Table 9 for states reporting. A noticeable gap is the lack of specific use reports from California.

Mevinphos usage ranges from 0%-70% of commodity area treated, with Nevada reporting the highest percent usage. Few data are available concerning seasonal and regional variability in the use of mevinphos. In general, mevinphos was not recommended for major uses on a very large percentage of commodity areas. Mevinphos has been used in 1988 in Nevada and may be recommended in isolated cases in 1989. The frequency of application for mevinphos ranged from 1.0-4.0 within a single season, with the lower figure predominant. Only Oregon reported more than 2.5 applications in a single season. Where more than one application of mevinphos is applied, the interval is 2 or more weeks. The median for active ingredient applied per acre was 0.375 lb. In 1988, all applications were at this rate, while in earlier years the range was 0.25-1.0 lb/acre.

The usage of mevinphos for alfalfa seed production was concentrated on late season Lygus bug control. Some usage during flowering stages is common because of the short residual and minimum impact on pollinating bees.

In most northwest production areas, alfalfa seed producers apply insecticides with field sprayers pulled by closed-cab tractors. In the Columbia Basin of Washington, the insecticide is applied by aircraft. The preference for ground application stems from the better performance (% kill) when sprayed 2-3 feet over the canopy with high gallons per acre.

In agroecosystems in California, mevinphos is used only occasionally as a pre-harvest treatment after honey-bee hives are removed from fields. There the crop rotation and pest complex differ because of the proximity of alfalfa seed cropping with cotton and safflower. There honeybees are relied upon to pollinate seed fields, whereas in the major seed producing areas in the Northwest, the wild bees (alkali bee and leafcutting bee) are depended upon almost exclusively for pollination. Honeybees differ from the wild bees with respect to insecticide tolerance and, where appropriate, mevinphos is used in Fresno and Kin Counties because of its efficacy and low hazard to bees. Overall, the extremely short residual of mevinphos does not affect cropping rotation or re-use of land.

The production of alfalfa seed is closely tied to integrated pest management systems developed in Oregon, Idaho, Nevada and the San Joaquin Valley area of California. The classic components of alfalfa seed IPM are: 1) weekly sampling to monitor pest and beneficial insect populations, 2) use of research-derived economic thresholds, 3) conservation of beneficials and

4) selective use of pesticides. Mevinphos plays an important role in 4 (above) because of its safety to pollinating bees when applied after sunset, and its short residual, and hence, short pre-harvest interval of one day. Additionally, mevinphos is highly effective for Lygus bug control.

All pesticides which are highly efficacious and destroy high percentages of the target population exert high selective pressures and consequent opportunity for build-up of pest resistance. Mevinphos is used only sparingly in most areas, and with the exception of Nevada, is restricted to a small percentage of commodity land area. The short residual on plant surfaces is a benefit in that immigrating insects are not affected, allowing in-flow of non-treated gene material. Overall, in alfalfa seed production, the potential for pest resistance to mevinphos is low. However, some Lygus resistance has been observed in California where insecticide treatments to the larger agroecosystems encourages area-wide resistance and cross-resistance especially to the older organophosphates that have been used extensively in California for many years.

Despite a highly evolved IPM for alfalfa seed production, pesticides occupy a key place in control of primary and secondary pests. The alternatives to mevinphos, presently, are four other pesticides under special review: carbofuran, oxydemeton-methyl (ODM), naled, and trichlorfon. Each has unique properties which impact the selectivity needed to control specific pests at specific times in the growth and harvest of alfalfa seed. The comparative efficacy or economic value of these insecticides cannot be evaluated in a vacuum, one against another, for a single pest species. Rather, their value must be judged in the context of a complex insect pest-pollinator management system with its subtle overlay of environmental factors. Table 10 modified from the Pacific Northwest producer's group report "Biological and economic assessment of ODM, trichlorfon, naled, and mevinphos in alfalfa seed production" compares alternative insecticides for use in the flowering-pollinating period critical to production of seed.

Field Corn

Constraints to corn production in the United States include a multitude of insect pests which may affect corn from planting to maturity. Management of the more important insect pests requires planned cultural and insecticide programs that reduce real or potential damage throughout the production season. The extent of insect management efforts in corn should be, and is, largely to prevent yield reduction. In most areas of the U.S. where insecticides are used in management programs, some degree of insecticide persistence is required. This is particularly true in the major corn-growing region of the Midwest. In fringe areas or areas where corn may be grown for special purposes, unique insect problems often occur. In these areas, insecticide requirements may also be unique.

Only two states, Colorado and Florida, responded to field corn production questions on a survey to determine mevinphos use and need in the U.S. Within these two states, an average of 820 acres was treated annually with mevinphos from 1982 through 1986 (Table 11). In Colorado, mevinphos was

Table 10. Effectiveness of Insecticide Use on Alfalfa in the Northwest During Flowering-Pollination Period.

Chemical	Rates (lbs. a.i./A)	Effect on 1° Pest <u>Lygus</u> bugs	Effect on 2° Pests	Impact on Beneficials	Other Advantages or Disadvantages
Oxydemeton methyl (ODM)	0.375-0.5	Excellent on early season Lygus, good to fair on late season Lygus.	Good on pea aphid Some mite suppression	Moderate	Systemic action, Relatively bee safe.
Naled	0.5-1.0	Very good on late season Lygus	Good on grasshopper, Fair on Lepidoptera	Severe	High bee risk, weather conditions need to be carefully monitored when applying.
Mevinphos	0.25-0.5	Very good on late season Lygus	Good on grasshopper Fair on alfalfa weevil. Some use to suppress Chalcid.	Very Severe	Relatively safe on bees. High application hazard.
Trichlorfon	0.5-1.0	Poor on Lygus except in Montana and other select areas.	Very good on small Lepidoptera	Moderate	Relatively bee safe. Resistance problem with Lygus.
Permethrin	0.1-0.2	Inconsistent on Lygus	Inconsistent on aphids	Very Severe	Mite outbreak following treatment. Seed specialists recommend against use because of high risk of bee kills.

Table 11. Use of Mevinphos on field corn for grain and forage (average annual), 1982-1986.

State	Acres Treated	Pests	Alternatives	Rate (lbs. a.i./A)	Applic Method	Limitations ¹
Corn - Grain						
Colorado	320	Fall Armyworm Earworm	permethrin	0.10	Air	1 day
			chlorpyrifos	0.50	Air	30 days
			fenvalerate	0.10	Air	21 days
			trichlorfon	0.50	Air	1 day
Corn - Forage						
Florida	500	Aphids	Parathion (ethyl)	.20	Ground	1 day
			Parathion (methyl)	1.00	Ground	1 day

¹Pre Harvest Interval

applied to control earworm [Heliothis zea (Boddie)] and fall armyworm [Spodoptera frugiperda (Smith)]. In Florida, mevinphos was used to obtain control of corn leaf aphid [Rhopalosiphum maidis (Fitch)] in the same application with another insecticide or fungicide.

Insect biology is an important factor when considering pest management. In the mevinphos use survey, corn earworm and fall armyworm were the most important field corn pests reported from Colorado, and corn leaf aphid was the most important pest reported from Florida.

The corn earworm, Heliothis zea (Boddie), and fall armyworm, Spodoptera frugiperda (Smith), overwinter as pupae (in southern latitudes of the U.S. and in South and Central America, fall armyworm may overwinter in all stages). Adults emerging from overwintering stages lay eggs on wild and cultivated hosts during the early season. Following egg, larval and pupal development, moths from this first and subsequent generations migrate to more northern regions. In these northern regions, corn is the preferred host.

Damage by corn earworm results from larval feeding in the whorls of immature plants and on silks and kernels of developing ears during and following pollination. Injury appears in the form of ragged leaves in the whorl and overall reduced plant vigor. This damage may result in delayed silking, poor pollination and yield loss for plants severely affected. Direct losses of 2 to 4% per developing ear may result from larval feeding in the silk channel and on developing kernels (Blanchard and Douglas, 1953; USDA Handbook, 1965).

Fall armyworm damage to corn usually occurs in Colorado during the same time period or slightly later than for corn earworm damage. Fall armyworm damage usually results from whorl and ear feeding by developing larvae.

Fall armyworm is found throughout the year in South America, Central America, and the southeastern U.S. But each year it migrates northward, arriving in the central U.S. and Colorado in late June to late July. After arrival, fall armyworm usually undergoes two generations. The first generation adults oviposit on leaves, and emerging larvae feed on leaves in the unfolding whorl and on developing ears. The fall armyworm life cycle is most frequently synchronized with the appearance of the silks during the pollination period. Developing larvae then feed primarily on the ear. Damage to the ear is often similar to that of earworm. However, unlike earworm, this species feeds on and into the basal portion of the ear. This type of damage is usually the most destructive, and under heavy infestation levels can result in severe yield losses. Fortunately, damage by this insect is less frequent than for corn earworm.

Colorado growers elected to treat approximately less than 1% annually of planted corn acres (660,000) with mevinphos during 1982-86 for either (or both) corn earworm and fall armyworm to avoid perceived and real potential yield reduction. Mevinphos was reported as having no distinct advantages for corn earworm and fall armyworm control over the four alternative insecticides listed in Table 11.

The corn leaf aphid, Rhopalosiphum maidis (Fitch), was identified by Florida entomologists as the most important aphid pest of corn used for forage. In the warmer climates of the U.S. (including California, Florida and Texas), aphids can reproduce throughout the year and multiply rapidly under favorable conditions. Corn leaf aphids feed on tassels and silks and coat them with honeydew, which results in reduced pollination. Severely infested tassels may become depleted of moisture and drop from the plant. Thus, overall vigor of the plant may be affected. Corn leaf aphid is also noted for transmission of pathogenic viruses (including maize dwarf mosaic) which adversely affect plant growth, yield and quality of grain and forage.

Mevinphos applied for control of aphids (all species) was approximately 1% or 500 of 46,200 acres per year during 1982-86. This insecticide apparently offers no distinct advantage over alternatives ethyl and methyl parathion (Table 11). Mevinphos is slightly more effective for aphid control, and only one application is needed to obtain control. This application is often made in conjunction with other insecticides or fungicides.

In Colorado, corn is grown in sandy soil regions where irrigation is available and provides an economical return on investment. Corn is followed by corn in production fields and is grown exclusively for grain. In these sandy soils, herbicides are not always effective for grass control and because of intensive irrigation and rapid corn growth, only one cultivation for weed control is possible. In some areas, cultivation may not be possible because of increased erosion potential.

Late planting, coupled with grassy weeds, attracts corn earworm and fall armyworm adults and provides an ideal mating and oviposition habitat. To prevent yield losses during years of heavy infestation by one or both pests, one application of an effective insecticide, mevinphos or alternatives (Table 11), is made during the late-whorl to full pollination stage.

Field corn produced in Florida is used primarily as forage for beef cattle production. Ear development is important for maximum yields and food value. Total yield is dependent on adequate rainfall, as well as proper fertility and agronomic practices. In Florida, the corn leaf aphid and other aphids are present every year and become important to corn production when populations peak during tassel emergence and silking. Pollination may be reduced or prevented.

Efforts to control aphids by one application of mevinphos or other short residual insecticide are usually made during years of adequate rainfall. During dry or unproductive seasons, profitable returns on insecticide investment for aphid control are usually not anticipated. Aphids are not controlled during such seasons and are made part of the forage if they are still present at time of ensilage chopping.

Extension entomologists in both Florida and Colorado reported no hazard to other crops, livestock or the surrounding environment.

In Florida, corn is grown in sparsely populated regions. Mevinphos or alternative parathion (Table 11) is applied only by experienced applicators.

Corn production fields are usually rotated to one or more crops including soybeans, peanuts, tobacco and infrequently, vegetables. Corn is an essential crop to reduce nematode problems in these rotation schemes.

In Colorado, corn is produced in isolated sites away from populated areas. Aerial application is the choice method because of extremely low hazard potential.

Insecticide resistance in highly mobile species usually occurs only when insecticides with long residual activity are applied frequently at sub-lethal dosages over extensive regions. Corn earworm, fall armyworm and aphids are mobile, but mevinphos does not fit into the mode of resistance development. Because mevinphos is used only once per annum and has no residual activity, potential for development of resistance is negligible.

Insecticide controls remain an important method of managing insects. Where used prudently and judiciously, losses and environmental disruption are minimized. This is particularly true where insecticide use is necessary only once per season and residues are non-existent. Mevinphos and certain other organophosphate-type insecticides fit this description. The insects described previously may be sporadic in building to economic levels requiring an insecticide application. According to returned questionnaires, only one application of mevinphos or alternative compounds is required during the season. Alternative controls for corn earworm and fall armyworm in Colorado and aphids in Florida are limited (Table 11). Cultural control options are few and limited primarily to planting date. Early planting is weather dependent but is practiced when possible because the later the planting, the greater the infestation.

The insecticides listed as alternatives by Colorado for corn earworm and armyworm control include two organophosphates, trichlorfon and chlorpyrifos; and two synthetic pyrethroids, fenvalerate and permethrin. These alternative insecticides provide comparable control, are slightly less hazardous to the applicator and provide residual activity.

Sorghum for Grain and Forage

Insects pose a sorghum production constraint wherever it is grown in the United States. From planting through grain formation insects can influence yield. As a result management of the more important insects pests requires planned cultural and insecticide programs that reduce economic yield loss throughout the production season. In different geographic locations insecticide usage may be unique.

Florida and Colorado were the only two states that responded to a questionnaire to determine if mevinphos is indeed needed for sorghum production. In Colorado, between 1982 and 1986, mevinphos was applied annually to control greenbugs on an average of 640 acres: 320 acres for grain and 320 acres for forage. Similary in Florida, between 1982 and 1986, mevinphos was applied annually to an average of 450 acres of grain sorghum and 1250 acres of forage sorghum. Florida respondents did not state specifically what pest was treated.

In Colorado, aphids, corn earworms, fall armyworms, climbing cutworms and armyworms were listed as sorghum insect pests. In Florida, aphids, corn earworms, fall armyworms, and sorghum webworms were listed as sorghum insect pests. Also, both states recommend the use of mevinphos for controlling these insect pests.

Insect biology is an important factor when considering pest management. In the mevinphos survey the greenbug aphid and fall armyworm are the most important in Colorado. In Florida, aphids, corn earworms, fall armyworms and sorghum webworms have equal importance.

The greenbug, Schizaphis graminum (Rondani), is prolific and of general distribution and has great prolificacy. In some years, it has reduced sorghum production 25%. While feeding the aphid injects a toxic saliva which causes discoloration and tissue destruction. In the southern states this insect passes the winter in the active nymphal and adult stages and this is generally true in both Florida and Colorado. Economically, the decision to treat is based on the amount of damage to the plant at different stages of growth. Example: Emergence to six-inch, treat when plants begin to yellow and visible colonies are present on the leaf; Six-inch to pre-boot, treat when colonies cause red spotting or yellowing on leaves but prior to the death of any one leaf; Pre-boot to heading, treat before the death of one functional leaf; Heading to hard dough, treat when numbers are sufficient to kill two normal sized leaves.

The female aphid gives birth to living young which may become either winged or wingless. Each female begins reproduction between 7 and 18 days of age and continues to reproduce for 20 to 30 days giving birth to 50 to 60 offspring. The winged form can migrate to nearby plants, or for several hundred miles.

The corn leaf aphid, Rhopalosiphum maidis, was also identified by Florida entomologists as an important sorghum aphid. Again, this aphid can

reproduce throughout the year and multiply rapidly under favorable conditions. Corn leaf aphids feed on the underside of the leaf and in the whorl area as the sorghum head emerges. The feeding of this aphid can reduce plant vigor and/or reduce grain fill by feeding on the newly emerging head.

The corn earworm, Heliothis zea (Boddie), and fall armyworm, Spodoptera frugiperda (Smith), are both important pests of sorghum. The corn earworm over winters as a pupa in the southern latitudes of the United States. The fall armyworm may overwinter in all stages. Adults emerging from over wintering stages lay eggs on wild and cultivated hosts during the early season. Following egg, larval and pupal development, moths from first and subsequent generations migrate to more northern regions.

In Colorado both pests feed on foliage and heads. Generally they affect yield when they feed in the whorl and damage the growth bud, or when they feed on the head. In Colorado, they are considered economical if there are two or more larvae per head.

The sorghum webworm, Celama sorghiella (Riley), is also considered a major pest of sorghum in Florida. It is particularly damaging to the ripening kernels of grain in the sorghum heads.

In Colorado the insect pest problems occur within a time frame that will allow control with the alternative insecticides listed in Table 12, and still allow for the P.H.I. of the alternatives. However, according to survey results Florida entomologists, none of the alternatives listed in Table 12 will allow sorghum to be made into silage. This probably accounts for the 1,250 acres of mevinphos used on forage sorghum (1982-1986). Survey results from Florida respondents state that a 5-10% yield reduction would possibly occur without the use of mevinphos for forage sorghum.

Also, in Florida sorghum seed heads are vulnerable to the sorghum webworm and fall armyworm late in the season. As a result, a short P.H.I. is necessary some years. Carbaryl has a zero day P.H.I. and mevinphos has a three day P.H.I. for grain sorghum. Carbaryl does not provide as effective control for these pests, therefore the Florida entomologists feel that a 5% yield loss could result if Mevinphos were not available for use near harvest.

Florida entomologists reported no hazard to other crops, livestock or the surrounding environment. In Florida, sorghum is grown in sparsely populated regions. Mevinphos or most of the alternatives on Table 12 are applied by commercial applicators, so mevinphos doesn't present an unusual risk problem.

Currently in Florida, mevinphos is used on 5% of the forage sorghum acreage and 2.2% of the grain sorghum acreage annually. This accounts for a total of 1,700 acres. The remainder of the acreage is treated with the alternatives listed in Table 12.

The three day P.H.I. is probably the most important advantage of mevinphos regarding sorghum production plus the fact that it is the only insecticide registered for use on sorghum intended for silage.

Table 12. Use of Mevinphos on Sorghum (average annual), 1982-1986.

<u>State</u>	<u>Acres Treated</u>	<u>Pests</u>	<u>Registered Insecticides</u>	<u>Rates (lbs. a.i./A)</u>	<u>Limitations P.H.I.</u>
SORGHUM FOR FORAGE					
Florida	1,250	Aphids	mevinphos dimethoate parathion	0.125 to 0.50 0.4 0.2	3 day 28 day 15 day
		Corn Earworm> Fall Armyworm> Sorghum Webworm>	methomyl parathion carbaryl	0.4 0.2 0.2	14 day 15 day 0 day
Colorado	320	Aphids	mevinphos chlorpyrifos dimethoate	0.125 to 0.50 0.25 0.25 to 0.50	3 day 30 day 28 day
		Armyworm> Climbing Cutworm> Corn Earworm> Fall Armyworm>	methomyl chlorpyrifos mevinphos carbaryl	0.25 to 0.50 0.50 to 1.00 0.25 to 0.50 1.00 to 2.00	14 day 30 day 3 day 0 day
SORGHUM FOR GRAIN					
Florida	450	Aphids	mevinphos dimethoate parathion	0.125 to 0.50 0.4 0.2	3 day 28 day 15 day
		Corn Earworm> Fall Armyworm> Sorghum Webworm> Climbing Cutworm>	mevinphos parathion carbaryl methomyl	0.125 to 0.50 0.2 0.2 0.4	3 day 15 day 21 day 14 day
Colorado	320	Aphids	dimethoate chlorpyrifos mevinphos	0.25 to 0.50 0.25 0.125 to 0.50	28 day 30 day 3 day
		Armyworm> Climbing Cutworm> Corn Earworm> Fall Armyworm>	methomyl chlorpyrifos mevinphos carbaryl	0.25 to 0.50 0.50 to 1.00 0.25 to 0.50 1.00 to 2.00	14 day 30 day 3 day 21 day

VEGETABLE CROPS

Mevinphos is a vital pesticide to agricultural crops because of its wide spectrum-control of insects. It is especially important for vegetable crops because the short residual allows growers to control aphids and other insects close to harvest and allows them to market commodities without a loss.

Aphids are one of the most important pests controlled by mevinphos; they are economically important on most vegetable crops in all the states reported. Aphids are small, variously colored depending on species, and occur in clusters. At times, they are so dense that the entire plants may be covered if they are not controlled. When aphid populations reach high levels, they secrete honeydew, transmit diseases, cause young plants to lose vigor, decrease the maturation rate, and may kill the plant. If not controlled, heavy yield losses occur from contamination of the edible portion from honeydew, or as a result of diseases dissemination by aphids. In warm climates, aphids may remain active throughout the year, and some species give rise to living young all year. Most aphids are wingless when plants are young, but when plants harden-off or when populations expand and crowding occurs, winged forms are produced for migration.

Cabbage looper, a migratory lepidopteran, is a general foliage feeder on vegetable crops and is usually found in all regions where vegetables are produced. The destructive larva is light green with white or pale yellow stripes, has three pairs of prolegs, and moves with a characteristic looping motion. The adult white moth emerges in the spring and deposits dome-shaped, pale green eggs on host plants, mostly at night. After hatching, the larva reaches full development in 2-4 weeks. After 8-10 days in the pupal stage, new adults emerge. Three or more generations are produced each season.

Cabbage looper can be a severe pest to lettuce and cole crops. Loopers damage plants by eating ragged holes in leaves, boring into heads, and contaminating head and leaves with their bodies and frass. High populations can chew seedlings severely enough to kill or slow down the maturation of the crop. Heads contaminated with, or tunneled into, by loopers are not marketable.

Diamondback moth is a primary microlepidopterian pest of cole crops. In California and throughout the south, it is a year-round pest. The small yellowish-white eggs are laid singly, or less commonly in groups of two or three, on the undersides of leaves. The eggs are scalelike and very difficult to detect. In a few days the very small greenish larvae hatch and feed mostly on the underside of outer or older leaves of older plants, chewing small holes, or at the growing points of younger plants. Growing point injury causes the most serious losses. Damage to crowns of young plants can severely stunt growth. The larvae may also chew the growing points, or bore into developing heads of broccoli, cauliflower, and cabbage, causing serious damage or contamination (U.C., Pest Management Manual on Cole Crops, 1985). They become full grown in 10-14 days, depending on temperature, and then spin loose white cocoons, which they attach to leaves or stems.

Adult moths are small and slender, grayish or brownish. When the moth is in its normal resting position, the wings of male moths display three diamond-shaped markings on their back that inspire the name "diamondback moth". The folded wings flare-up and out at the tip.

Imported cabbageworm is a widely distributed primary lepidopteran pest of cole crops. Adults are white to yellowish white butterflies with one to four black spots on their wings. They lay tiny white to yellow eggs, usually on undersides of the leaves. These hatch in seven or more days into velvety green, pubescent caterpillars, each having a narrow orange stripe dorsally and paler broken stripes laterally. Caterpillars are sluggish, but they eat voraciously. Young caterpillars chew holes in both the outer and inner leaves; older caterpillars move toward the center of the plant, often feeding at the base of the wrapper leaves or boring into the heads of cabbage, broccoli or cauliflower.

Imported cabbageworm damage is similar to that caused by cabbage looper. The caterpillars chew large, irregular holes in the leaves, bore into heads, and drop greenish brown fecal pellets which contaminate the marketable leaves and heads. Seedlings may be destroyed or crop maturity may be delayed with high populations. The most serious losses occur when the imported cabbageworm bores into the heads or feeds on marketable leaves.

Artichokes

Mevinphos is registered for use on artichoke for control of aphids at 0.125 lb ai/acre and artichoke plume moth at 0.25 lb ai/acre. Both uses have a two day preharvest interval. Artichoke production in the U.S. is predominantly in California where the sole use of mevinphos on artichoke is for plume moth control. Mevinphos has typically been applied once or twice per season to an average of 10,000 acres of artichokes.

The only alternative is azinphosmethyl, which is an unsatisfactory replacement for mevinphos because of its long (30-day) preharvest interval. Non-chemical controls do not give reliable insect control on artichokes. Mevinphos is needed for this crop.

Beans (Snap, Lima)

Mevinphos is used on snap bean acreage in some states for insect control close to harvest (Table 13). Arthropod pests controlled include aphids, leafhoppers, thrips, and mites. Five states reported usage of mevinphos on snap beans with the major advantage over alternative insecticides being the short pre-harvest interval. Other alternatives include dimethoate and acephate. The new synthetic pyrethroids labeled in recent years will control leafhoppers but not aphids, mites, or thrips. They have a longer post harvest interval (fenvalerate, 21 days), as do acephate and dimethoate (14 days).

Thrips control is an important use in North Carolina for mevinphos. It

Table 13. Use of Mevinphos on Beans (average annual), 1982-1986.

State	Acres treated	Pests	No. Appl.	Rate (lbs. a.i./A)	Alternatives	Limitations
<u>Snap beans</u>						
Tennessee	3500	aphids	2	1/4	acephate, dimethoate	longer PHI
Wisconsin	1000	potato leafhopper	1	1/4-1/2	acephate dimethoate fenvalerate	longer PHI
Arkansas	1500	mites	2	1/4-1/2	dimethoate	
Florida	2269	aphids mites leafhoppers	1	1/4-1/2	acephate methomyl dimethoate dicofol	longer PHI
North Carolina	2000	thrips	1	1/4	dimethoate methomyl carbaryl	less control increases mites, kills bees
<u>Lima beans</u>						
Florida	23	aphids grasshoppers mites	2	1/4-1/2	methomyl acephate dimethoate	longer PHI

was preferred over dimethoate for that use and also preferred over carbaryl for caterpillar control (Table 13).

Beets

Mevinphos is registered for use on table beets for control of aphids, cabbage looper, climbing cutworm, dipterous leafminer adults, false chinch bug, grasshoppers, imported cabbage worm, leafhopper, mites and salt marsh caterpillar; it has a 3 day preharvest use restriction. Apparently the only significant use of mevinphos on table beets is for control of aphids and leafminers in Wisconsin. The principal advantage of mevinphos is its extremely short residual activity; however on table beets this is not of major importance since insect damage occurs on a portion of the crop which is not harvested. Diazinon, trithion and parathion are all viable alternatives to mevinphos for aphid and leafminer control on table beets.

Broccoli

Most of the commercial broccoli is grown in nine states with a total of 108,236 acres. Mevinphos is recommended in all nine states primarily to control aphids, cabbage looper, diamondback moth, and imported cabbageworm. Mevinphos was applied to broccoli at least one time in all nine states; in some states e.g., Alabama, it was applied 5 times. In California, there are several effective alternatives for control at early crop stages but growers must turn to mevinphos close to harvest because of its short preharvest interval. Other states (e.g. Alabama, Florida, Illinois, Louisiana, New Jersey, and Texas) have several effective alternatives registered to control the lepidopterous pest complex. However in Alabama, the diamondback moth develops resistance to the pyrethroids rapidly. Mevinphos in combination with Dipel has proven to give the most effective control. Controlling insects on broccoli is most critical during head formation, therefore a quick knock down short residual material such as mevinphos is necessary at maturity because it can be applied until 2 days before harvest (Table 14).

Cabbage

Fourteen states responded to the survey, representing a total of approximately 67,818 acres. Mevinphos is recommended in all the states with each reporting at least one application; the greatest number of applications, 3, was reported from Florida. The major target pests were aphids and diamondback moth.

Mevinphos plays an important role in 8 of the 14 states in controlling aphids and diamondback moth. Without mevinphos, marketable head loss varies from 3-5% in Wisconsin to as high as 80% in Florida due to contamination or feeding damage to the head. Pyrethroids are effective alternatives used in North Carolina and Wisconsin but are more expensive, and there exists potential resistance. In Alabama and Louisiana, mevinphos used alone against diamondback moth was not effective, however, used in combination with B.

Table 14. Use of Mevinphos on Broccoli (average annual), 1982-1986.

State	Alternatives					Limitations
	Acres Treated	Rate (lbs. a.i./A)	Chemicals	Rates (lbs. a.i./A)	Pests ¹	
Alabama ²	300	0.5	<u>B. thuringiensis</u> dipel/Javelin esfenvalerate fenvalerate permethrin	Label 0.05 0.10 0.10	DBM DBM DBM DBM	longer PHI longer PHI longer PHI
California ³	40,372	0.25-0.5	permethrin diazinon dimethoate azinphosmethyl methamidophos oxydemeton-methyl methomyl carbaryl	0.05-0.1 0.25-0.5 0.25-0.5 0.5-0.75 0.5-1.0 0.25-0.5 0.9	C, ICW, DBM A, CL, ICW, DBM A DBM, CL, ICW A, DBM, CK, ICW A A, DBM, CL, ICW CL	longer PHI longer PHI longer PHI longer PHI longer PHI
Florida ⁴	71	0.25-0.5	methamidophos ethyl parathion endosulfan permethrin fenvalerate flucythrinate <u>B. thuringiensis</u> carbaryl ethyl parathion methyl parathion methomyl	1.0 0.2 1.0 0.2 0.1-0.2 0.1 Label 0.2-0.4 0.2 1.0 0.5	A, CL, ICW A, CW, LB, M A CL, LM, ICW CL, CW, GH, ICW CL, SM CL, SM GH, SM, CW LM LH ICW, SM	Toxic warm-blooded animals & beneficial resistance suspected, longer PHI longer PHI resistance suspected resistance suspected
Oregon	50% of acreage	0.25-0.5	oxydemeton-methyl permethrin methamidophos methomyl	 	A, CL, ICW A, CL, ICW A, CL, ICW	longer PHI, poor control longer PHI, poor control longer PHI

Table 14 Continued

New Jersey	diazinon	0.25	65	0.5	A	longer PHI
	dimethoate			0.5	A	longer PHI
	disulfoton			1.0	A	longer PHI
	oxydemeton-methyl			0.5	A	longer PHI
	methamidophos			1.0	A,CL,ICW	longer PHI
	esfenvalerate			0.03	CL,ICW	longer PHI
	<u>B. thuringiensis</u>			Label	CL,ICW	longer PHI
	azinphosmethyl			0.75	CL,ICW	longer PHI
	methomyl			0.9	CL,ICW	longer PHI
	flucythrinate			0.04	CL,ICW	longer PHI
	permethrin			0.2	CL,ICW	longer PHI
	fenvalerate			0.2	CL,ICW	longer PHI
Texas	acephate	0.25-0.5	50	0.5-1.0	A,LCX	
	azinphosmethyl			0.5-0.75	A,LCX	
	<u>B. thuringiensis</u>			Label	LCX	
	Diazinon			0.25-4.0	A,CW,LCX	
	endosulfan			0.75-1.0	A,LCX,CW	
	esfenvalerate			0.025-0.5	LCX,CW	
	fenvalerate			0.1-0.2	LCX,CW	
	malathion			0.625-1.25	A,LCX	
	methamidophos			0.5-1.0	A,LCX	
	methyl parathion			0.25-1.25	A,LCX,M	
	ethyl parathion			0.5	A,LCX	
	naled			2.0	LCX	
	permethrin			0.05-0.1	A,LCX	
	phosphamidon			0.5-1.0	A,LCX	
	carbaryl			2.0	CW	
	chlorpyrifos			0.5-1.0	CW	
	dimethoate			1.0	A	
	disulfoton			0.25-0.5	A,M	
	oxydemeton-methyl			0.375-0.75	A,M	

¹A=aphids; CL=cabbage looper; CW=cutworm; DBM=diamondback moth; GH=grasshopper; ICW=Imported cabbageworm; LB=Lygus bug; LXC=caterpillar complex; LM=leafminers; M=mites; SM=saltmarsh caterpillar.

225% loss

3 15-20% loss

4 15% loss

Table 15. Use of Mevinphos on Cabbage (average annual), 1982-1986.

State	Acres Treated	Rate (lbs. a.i./A)	Alternatives		
			Chemicals	Rates (lbs. a.i./A)	Pests ¹
Alabama ²	100	0.5	esfenvalerate fenvalerate methomyl permethrin <u>B. thuringiensis</u>	0.05 0.1 0.45 0.1 Label	DBM DBM DBM DBM DBM
California ³	6,178	0.25-0.5	permethrin diazinon dimethoate azinphosmethyl methamidophos oxydemeton-methyl endosulfan methomyl trichlorfon	0.05-0.1 0.25-0.5 0.25-0.5 0.25-0.5 0.5-1.0 0.25-0.75 0.25-0.75 0.45-0.9 0.25-0.5	DBM, CL, ICW DBM, CL, ICW, A A A, DBM, CL, ICW A, DBM, CL, ICW A, DBM, CL, ICW A, DBM, CL, ICW A, DBM, CL, ICW A, DBM, CL, ICW A, DBM, CL, ICW
Florida ⁴	10,461	0.25-0.5	methamidophos ethyl parathion endosulfan permethrin fenvalerate flucythrinate <u>B. thuringiensis</u> carbaryl methomyl methyl parathion	1.0 0.2-0.4 1.0 0.2 0.2 0.1 32 billion (Btu) 0.2-0.4 0.5 1.5	A, CL, LM A, CW, LM, M, BL A CL, ICW, LM CL, CW, ICW, GH CL, SM CL, SM CW, GH, SM ICW LH
Delaware ⁵	500	0.25	fenvalerate flucythrinate methamidophos Bt(k) new strain methomyl permethrin	1.7 oz 2 oz 2 pts 2 pts 2-4 pts 8 or 12.8 oz	A, DBM A, DBM A, DBM A, DBM A, DBM A, DBM
					longer PHI, resistant longer PHI, resistant
					longer PHI
					longer PHI
					resistant

Table 15 Continued

Georgia ⁶	5,000	0.75	<u>B. thuringiensis</u> fenvalerate esfenvalerate methomyl methamidophos	8-10 BTU 0.1-0.2 0.03-0.05 0.45-0.9 0.5-1.0	A, DBM A, DBM A, DBM A, DBM A, DBM	longer PHI
Illinois	800		permethrin fenvalerate esfenvalerate dimethoate	0.2 0.2 0.07 0.5	A, DBM, ICW, CL A, DBM, ICW, CL A, DBM, ICW, CL A, DBM, ICW, CL	
Missouri			No Alternative		LCX	
New Jersey	484	0.25	diazinon dimethoate oxydemeton-methyl methamidophos disulfoton esfenvalerate <u>B. thuringiensis</u> azinphosmethyl methomyl methamidophos flucythrinate methyl parathion permethrin	0.5 0.5 0.5 1.0 1.0 0.03 Label 0.75 0.9 1.0 0.04 1.15 0.2	A A A A A CL, ICW CL, ICW CL, ICW CL, ICW CL, ICW CL, ICW CL, ICW CL, ICW CL, ICW	longer PHI
North Carolina ⁷	500	0.25	Bt Dipel permethrin fenvalerate Bt Javelin Pyrethroids + PBO	8 oz 0.1 0.2	DBM DBM DBM DBM DBM	longer PHI
New York ⁸	4,000	0.5	permethrin Bt Dipel Bt Javelin esfenvalerate endosulfan	0.1 1.0 1.0 0.1 0.5	CL, DBM, TH CL, DBM CL, DBM CL, DBM CL, DBM, TH	
South Carolina ⁹	450	0.5	endosulfan <u>B. thuringiensis</u> methomyl methamidophos	1.0 + 0.5 0.45 1.0	CL, DBM, ICW CL, DBM, ICW CL, DBM, ICW CL, DBM, ICW	longer PHI expensive

Table 15 Continued

Utah ¹⁰	200	methamidophos	0.5-1.0	A, CL	longer PHI
		methomyl	0.45-0.9	A, CL	
		<u>B. thuringiensis</u>	1-2 qts	A, CL	
		naled	2.0	A, CL	
		endosulfan	1.0	A, CL	
		permethrin	0.05-0.1	A, CL	
		parathion	0.5	A, CL	
		Malathion	0.75-2.5	A, CL	
		azinphosmethyl	0.5-0.75	A, CL	
		fenvalerate	0.1-0.2.	A, CL	
Wisconsin ¹¹	1, 200	azinphosmethyl	0.5	A	longer PHI
		demeton	0.5	A	longer PHI
		diazinon	0.5	A	longer PHI
		dimethoate	0.25-0.5	A	longer PHI
		disulfoton	1.0	A	longer PHI
		endosulfan	0.5	A, CL, DBM, ICW	longer PHI
		malathion	0.5	A	
		methamidophos	0.5	A, CL, DBM, ICW	longer PHI
		naled	1.0-2.0	A, DBM, CL, ICW	longer PHI
		oxydemeton-methyl	0.5	A	
		ethyl parathion	0.25	A, CL, DBM, ICW	
		<u>B. thuringiensis</u>	0.5	CL, ICW, DMB	
		fenvalerate	0.05-0.2	DBM, CL, ICW	longer PHI, resistant
		methomyl	0.25-0.9	DBM, CL, ICW	
		permethrin	0.05-0.2	DBM, CL, ICW	

¹A=aphids; CL=cabbage looper; CW=cutworm; DBM=diamondback moth; GH=grasshopper; ICW=Imported cabbageworm; LB=Lygus bug; LCX=caterpillar complex; LH=leafhoppers; LM=leafminers; M=mites; SM=saltmarsh caterpillar; TH=thrips.

thuringiensis satisfactory control was achieved (Table 15).

Cauliflower

Cauliflower is grown in six of the reporting states with a total of approximately 47,665 acres. California is the leading state in total acreages with 45,566 followed by Florida, Tennessee, Louisiana, New Jersey, and Oregon. Mevinphos is registered and used in all six states for control of aphids and the lepidopterous larvae. One to two applications are made each year.

Several effective alternative chemicals are used. Some of the alternatives used for aphid control are methamidophos in California, Tennessee, New Jersey, and Oregon; oxydemeton methyl, diazinon, dimethoate, and endosulfan in California. Permethrin, esfenvalerate and methomyl are other alternatives which are used effectively against the lepidopterous complex in California, Louisiana, New Jersey, and Oregon. Florida and Louisiana had a problem controlling the diamondback moth with methomyl or mevinphos because of resistance. However, in Louisiana, mevinphos used in combination with Bacillus thuringiensis gave excellent control (Table 16).

Carrots

Three states (Florida, Texas and Wisconsin) reported the use of mevinphos (Phosdrin) on carrots. Mevinphos is registered on carrots for the following pests: aphids, leafhoppers, Lygus bugs, mites, cabbage looper, dipterous leafminer (adult), cutworm (climbing) and saltmarsh caterpillar (Table 17).

The rates of mevinphos range from 1/4 to 1/2 pint per acre for aphids and from 1/2 pint to 1 pint per acre for the other pests. The PHI is 2 days. The total acreage treated with mevinphos is 5,035 acres with an average of 1 application on carrots.

Survey results indicate that if mevinphos were not available and aphids were a problem within 1 week of harvest, a 10% loss in yield could result. Aphids could cause some damage to carrots, both early and late in the growing season. The control costs would rise some 25% when mevinphos is compared to other known chemical or nonchemical control alternatives. As few materials are registered on carrots, mevinphos is important to insect management on carrots. Mevinphos' high performance activity coupled with a short preharvest interval (2 days) make it superior to all alternatives.

Registered alternatives for aphids are malathion, diazinon, ethyl parathion, parathion (methyl) and endosulfan. The alternatives have longer PHI's: malathion (7 days), diazinon (10 days), ethyl parathion (15 days), methyl parathion (15 days), and endosulfan (7 days). These materials cannot be used on carrots should aphid populations increase rapidly within 1 week of harvest. Hence, it is important to keep mevinphos registered on carrots because of its high performance activity and its short preharvest interval.

Table 16. Use of Mevinphos on Cauliflower (average annual), 1982-1986.

State	Acres Treated	Rate (lbs. a.i./A)	Alternatives			Limitations
			Chemicals	Rates (lbs. a.i./A)	Pests ¹	
California ²	31,231.94	0.25-0.75	acephate	0.5-1.0	A, CL, ICW, DBM	longer PHI
			permethrin	0.1-0.2	CL, ICW, DBM	longer PHI
			diazinon	0.25-0.5	A, CL	longer PHI
			dimethoate	0.25-0.5	A	longer PHI
			azinphosmethyl	0.5-0.75	A, CL, ICW, DBM	longer PHI
			methamidophos	0.5-1.0	A, CL, ICW, DBM	longer PHI
			oxydemeton-methyl	0.25-0.5	A	longer PHI
			endosulfan	0.25-0.75	A, CL	longer PHI
			trichlorfon	0.25-0.5	CL, ICW, DBM	longer PHI
			carbaryl	0.5-1.0	CL	longer PHI
			methomyl	0.45-0.9	A, CL, ICW, DBM	longer PHI
			methamidophos	1.0	A, ICW, LM	longer PHI, highly toxic
			acephate	0.2	A, LH	
			endosulfan	1.0	A	
Florida	285	0.5-1.0	permethrin	0.2	CL, ICW, LM	longer PHI, expensive
			fenvalerate	0.1-0.2	CL, ICW, CW	longer PHI, expensive
			flucythrinate	0.1	CL, SM	longer PHI, expensive
			<u>B. thuringiensis</u>	Label	CL, SM	
			carbaryl	0.2-0.4	GH, CW, SM	longer PHI, expensive
			ethyl parathion	0.4	CW, LM, LB, M	longer PHI, expensive
			methyl parathion	1.0	LH	longer PHI, expensive
			methomyl	0.5	ICW, SM	longer PHI, resistant
			oxydemeton-methyl	A		
			permethrin	CL, ICW		longer PHI
			methamidophos	A, CL, ICW		
			methomyl	A, CL, ICW		
			permethrin	0.1-0.2	CL, ICW	longer PHI
			fenvalerate	0.1-0.21	CL, ICW	longer PHI
Louisiana	550	0.25	esfenvalerate	0.025-0.05	CL, ICW	longer PHI
			methomyl	0.25-0.5	CL, ICW	longer PHI
			<u>B. thuringiensis</u>	Label	CL, ICW	
Oregon		0.25-0.5	oxydemeton-methyl	A		
			permethrin	CL, ICW		longer PHI
			methamidophos	A, CL, ICW		
			methomyl	A, CL, ICW		

Table 16 Continued

New Jersey	57	0.25				longer PHI
			diazinon	0.5	A	
			dimethoate	0.5	A	
			disulfoton	1.0	A	
			oxydemeton-methyl	0.5	A	
			methamidophos	1.0	A, CL, ICW	
			acephate	0.997	A, CL, ICW	
			esfenvalerate	0.03	CL, ICW	
			<u>B. thuringiensis</u>	Label	CL, ICW	
			azinphosmethyl	0.75	CL, ICW	
			methomyl	0.9	CL, ICW	
			flucythrinate	0.04	CL, ICW	
			permethrin	0.2	CL, ICW	
Tennessee	1,000	0.125-0.5	methamidophos	0.5-1.0	A, DBM	
			ethyl parathion	0.5	A, DBM	
			disulfoton	1.0	A	

1 A=aphids; CL=cabbage looper; CW=cutworm; DBM=diamondback moth; ICW=imported cabbageworm;
LB=Lygus bug; LH=leafhoppers; LM=leafminers; M=mites; SM=saltmarsh caterpillar.

2 40-50% loss

Table 17. Use of Mevinphos on Carrots (average annual), 1982-1986.

State	Acres Treated	No. Appl.	Rate (lbs. a.i./A)	Pests ¹	Alternatives	Limitations
Florida ²	3,285	1	0.25 (aphids) 0.50 (others)	aphids CL cutworms LH, LM Lygus mites SM	ethyl parathion endosulfan methomyl, BT ethyl parathion carbaryl ethyl parathion ethyl parathion methomyl carbaryl trichlorfon	very toxic/beneficials moderate control/fish kills suspected resistance to methomyl moderate control expensive
Wisconsin ³	250	1	0.13-0.15 (aphids) 0.5 (aster leafhopper)	aphids	endosulfan diazinon malathion ethyl parathion	all except endosulfan have a long PHI hazardous/4 day PHI
Texas	1,500	1	0.125-0.25	aphids	carbaryl fenvalerate methomyl ethyl parathion diazinon endosulfan malathion methyl parathion ethyl parathion carbaryl fenvalerate malathion methoxychlor methyl parathion methyl parathion ethyl parathion	fewer pests controlled

¹CL=cabbage looper; LH=leafhoppers; LM=leafminers; SM=saltmarsh caterpillar.²Need is minor unless aphids cause a problem within one week of harvest, in which case 10% loss in yield could result.³Need is minimal.

Celery

Four states (Michigan, Florida, Wisconsin and California) reported on the use of mevinphos on celery. Mevinphos is registered on celery for the following pests: aphids, dipterous leafminer (adult), Lygus bug, saltmarsh caterpillar, leafhopper, cabbage looper and mites.

The rates of mevinphos range from 1/4 to 1/2 pt/acre for aphids, from 1/2 to 1 pt. for other pests and 1 qt/acre for "hard-to-kill" insects. The time limitations between last spray and harvest are from 3 to 5 days depending on the rate used. The total acreage treated with mevinphos is 19,792 with an average of 1.7 applications on celery (Table 18).

Based on survey results yield decrease ranges from 2 to 45%. Aphid, lepidoptereous larval, and to a lesser extent mite, outbreaks within 1 week of harvest may result in some yield loss as well as in loss of marketability due to contamination. There is little change in control costs when mevinphos is compared to other chemical or nonchemical control alternatives. Mevinphos is a useful broad spectrum insecticide with cleanup capability that can be used close to harvest.

Registered alternatives for aphids are endosulfan, naled, parathion (ethyl) and acephate. These alternatives have longer PHI's: endosulfan (4 days), naled (7 days), parathion (ethyl) (30 days) and acephate (21 days). Alternatives do not provide adequate mite and aphid control near harvest.

Registered alternatives for cabbage looper are permethrin, acephate, B.t. and endosulfan. Other than B.t. these have a longer PHI, are extremely toxic to fish; B.t. is slow acting. Alternatives for leafhoppers are carbaryl and azinphosmethyl. Carbaryl may induce mite and aphid outbreaks. For leafminers, cyromazine may be used but it is expensive. For Lygus parathion or carbaryl may be used but they give moderate control only. And for mites, both parathion and demeton suppress mites, but are highly toxic to animals.

California considers mevinphos use necessary on its 11,562 acres of celery. There are no effective alternatives to the use of mevinphos on celery at harvest time for a broad spectrum of insect pests. Hence mevinphos continues as a viable option to late season pest control and cleanup.

Sweet Corn

Three states (Florida, Texas and California) reported the use of mevinphos on sweet corn. Mevinphos is registered on corn for control of aphids. However, mevinphos also provides some suppression of lepidopterous larvae especially as a quick knockdown product close to harvest.

The rates of mevinphos are from 1/4 to 1/2 pt. per acre for aphids. The time limitation between last spray and harvest is 2 days for mevinphos. The total acreage treated with mevinphos is 13,712 acres with an average of 1 application on sweet corn (Table 19).

Table 18. Use of Mevinphos on Celery (average annual), 1982-1986.

State	Acres Treated	No. Appl.	Rate (lbs. a.i./A)	Pests ¹	Alternatives	Limitations
Michigan	3,100	1 8 others	0.25-0.5	aphids	naled	
					endosulfan	less effective longer PHI
Florida	1,080	1	0.25 0.50	aphids others	acephate endosulfan	moderate control and toxic to fish
					naled ethyl parathion	not effective control toxic/bad on beneficials
					permethrin acephate	toxic to fish
					<u>B. thuringiensis</u> endosulfan	
				CL	carbaryl azinphosmethyl	lost effectiveness
					cyromazine	expensive
					ethyl parathion methyl parathion carbaryl	moderate control
					ethyl parathion methyl parathion demeton	highly toxic to animals
					naled	
Wisconsin	50	1	0.25-0.5	CL	B.T. endosulfan methomyl	fewer pests controlled longer PHI
					azinphosmethyl methomyl parathion	longer PHI
				Lygus		

Table 18 Continued

California	11,562	4	0.25-0.5	aphids armyworm CL	acephate diazinon azinphosmethyl methamidophos oxydemeton-methyl methomyl carbaryl	Alternative have longer PHI
------------	--------	---	----------	--------------------------	--	--------------------------------

1CL=cabbage looper; LH=leafhoppers; LM=leafminers; SM=saltmarsh caterpillar.

Table 19. Use of Mevinphos on Sweet Corn (average annual), 1982-1986.

State	Acres Treated	No. Appl.	Rate (lbs. a.i./A)	Pests ¹	Alternatives	Limitations
Florida	5,100	1	0.25	aphids	ethyl parathion methomyl endosulfan	toxic/beneficials toxic/beneficials resistance high fish kill
Texas	1,000	1	0.25	aphids	ethyl parathion oxydemeton-methyl fenvalerate endosulfan chlorpyrifos	toxic toxic phytotoxic high fish kill state registration
California	7,613	1	0.25-0.5	aphids/ lepidopterious larvae	diazinon dimethoate methomyl trichlorfon carbaryl	

There is no change in yield with or without mevinphos. There is a savings of 50% in control costs when mevinphos is compared to other chemical or nonchemical control alternatives. However, mevinphos does provide growers with another alternative in aphid control.

Registered alternatives for aphids are parathion, methomyl, endosulfan, fenvalerate and chlorpyrifos. The PHI's for these compounds are as long or longer; some are toxic to fish and beneficials, ineffective, phytotoxic or only registered in two states. The PHI's for alternatives are: parathion (12 days), methomyl (0 days), endosulfan (0 days), fenvalerate (1 day) and chlorpyrifos (21 days).

As aphids are only a problem on corn early in the season, the use of mevinphos and alternatives is not restricted or limiting. However, the use of mevinphos does provide growers with another alternative for aphid control and its use in IPM is important.

Cucumbers

Mevinphos is registered for use on cucumbers at 0.125 to 0.25 lb ai/acre for control of aphids and 0.25 to 0.50 lb ai/acre for control of grasshoppers, leafhoppers and mites. In Georgia, mevinphos is also applied to cucumber at 0.5 lb ai/acre for control of cucumber beetle and pickleworm. Use of mevinphos on cucumber occurs primarily in Florida, Georgia, New Jersey and Texas.

With the exception of pickleworm which feeds directly on the fruit, the target pests feed on the foliage and do not directly damage the harvested product. The foliage feeding pests can be effectively controlled on an "as-needed" basis when damaging infestations develop. Typically there is only limited need to control foliage feeding species once harvest begins, although aphid infestations requiring control can develop during the several week harvest period. It is in combatting these aphid infestations that mevinphos is particularly useful.

Mevinphos is used for pickleworm control only in Georgia. Pickleworm must be controlled from the onset of fruiting through harvest. Because cucumbers are harvested every 3 to 4 days, any insecticides used during the harvest period must have a very short preharvest interval; it is this feature of mevinphos (2 days PHI) that makes it well suited for this use. There are, however, several viable alternatives to mevinphos for aphid and pickleworm control during the fruiting period.

Because cucumber is a short term (65 to 75 day) crop and the insect pest problems are highly variable in intensity and frequency, naturally occurring biological control cannot be depended upon to prevent significant crop damage. Insecticidal control is important to successful cucumber production during certain periods in most, if not all, production areas. There are a number of currently registered insecticides which are viable but more expensive alternatives to mevinphos (Table 20). At present mevinphos is not essential to profitable cucumber production in any area of the US.

Table 20. Use of Mevinphos on Cucumbers (average annual), 1982-1986.

State	Acres Treated	No. Appl.	Rate (lbs. a.i./A)	Pests	Alternatives	Limitations
Florida ¹	784	1 to 2	0.25	aphids	methamidophos endosulfan emthomy ethyl parathion	cost moderate efficacy resistance suspected ---
					carbaryl fenvalerate	bee kills ---
					carbaryl fenvalerate ethyl parathion	bee kills ---
					ethion dicofol ethyl parathion	limited efficacy --- ---
Georgia ²	1,000	1 to 2	0.5	aphids	endosulfan dimethoate	none poor control of cucumber beetles
					carbaryl	poor control of aphids; bee kill
					endosulfan esfenvalerate	none poor control of aphids
New Jersey	242	2	0.5	pickleworm aphids	endosulfan esfenvalerate	none poor control of aphids
					methamidophos diazinon oxydemeton-methyl methomy endosulfan	long PHI long reentry

Table 20 Continued

Texas	700	1	0.125-0.25	aphids	diazinon dimethoate endosulfan malathion methamidophos methyl parathion naled oxydemeton-methyl parathion phosphamidon
			0.25-0.50	leafhoppers	azinphosmethyl carbaryl diazinon dimethoate esfenvalerate ethion fenvalerate malathion methamidophos naled ethyl parathion phosphamidon
			0.25-0.50	mites	diazinon dicofol ethion malathion methamidophos naled oxydemeton-methyl ethyl parathion phosphamidon
			0.25-0.50	mites	

¹Limitations are mild and mevinphos is not essential to cucumber production in Florida except in those instances when there is a severe aphid problem very close to harvest.

²Limitation are not severe. Mevinophos would be needed only if aphids develop resistance to other compounds.

Eggplant

Mevinphos was applied to eggplant acreage in the two reporting states (Florida and New Jersey) for effective and economical late-season, close-to-harvest control of leafhoppers, aphids, and mites. The alternative insecticides labeled for eggplant include endosulfan, oxymyl, carbaryl, methamidophos, and fenbutatin-oxide (Table 21).

Peppers

Mevinphos was used on 9500 acres of green bell pepper in four reporting states (New Jersey, Florida, Texas and Tennessee), probably for late-season outbreaks of mites or aphids. It was applied for the control of these two pests as well as for other insects in Florida. One to two applications were applied during the crop season. Registered alternative insecticides such as acephate and dimethoate have a longer pre-harvest interval but provide good green peach aphid control on peppers (Table 22).

Lettuce

Lettuce is grown primarily in nine states, comprising approximately 241,950 acres. California and Arizona comprise seventy to eighty percent of those acreages in the United States. All nine states recommended mevinphos for control of aphids and lepidopterous larval. In California and Arizona, mevinphos is a second choice to newer aphicides and larvicides that are recommended. These alternatives are not generally available to other states. Mevinphos is applied once or twice to the nation's lettuce crop each year.

Several effective alternative chemicals are used. However, most of them have longer pre-harvest intervals than does mevinphos. Colorado and New York reported no satisfactory alternative controls for lettuce pests (Table 23).

Melon

Mevinphos is registered for use on melons, including muskmelons and watermelons, at 0.125 to 0.25 lb ai/acre for control of aphids and at 0.25 to 0.5 lb ai/acre for control of cabbage looper, dipterous leafminer adults, leafhoppers, Lygus bugs, mites, false chinch bugs, salt-marsh caterpillars, climbing cutworms and grasshoppers. In addition it is registered at 0.25 lb ai/acre for control of the rindworm complex on watermelons. This complex includes cabbage loopers, cutworms, salt-marsh caterpillars and tobacco budworms which feed superficially on the developing fruit. There is a 2 days preharvest use restriction on all melons.

Mevinphos is used most extensively on melons in Florida and Georgia where it is applied primarily to watermelon with lesser quantities applied to cantaloupe. Mevinphos is also used in Texas on cantaloupe and honeydew melons and in California, New Jersey and North Carolina on cantaloupe and watermelons.

Table 21. Use of Mevinphos on Eggplant (average annual), 1982-1986.

State	Acres treated	Pests	No. Appl.	Rate (lbs. a.i./A)	Alternatives	Limitations
Florida	240	grasshoppers leafhoppers	1	1/2	carbaryl methamidophos fenbutatin-oxide	poor control expensive expensive
New Jersey	160	aphids	2	1/4	endosulfan oxamyl oxydemeton-methyl	longer PHI

Table 22. Use of Mevinphos on Peppers (average annual), 1982-1986.

State	Acres treated	Pests	No. Appl.	Rate (lbs. a.i./A)	Alternatives	Limitations
New Jersey	1129	aphids mites	2	1/4	dimethoate acephate methomyl endosulfan	longer PHI
Florida	6980	aphids leafhoppers grasshoppers mites	1	1/4-1/2	acephate methomyl carbaryl dicofol	longer PHI highly toxic to fish
Texas	1000	aphids mites	1	1/4-1/2	dicofol acephate dimethoate	-
Tennessee	500	aphids	1	1/8-1/4	endosulfan acephate	longer PHI

Table 23. Use of Mevinphos on Lettuce (average annual), 1982-1986.

State	Acres Treated	Rate (lbs. a.i./A)	Chemicals	Alternatives		
				Rates (lbs. a.i./A)	Pests ¹	Limitations
Arizona		1.0	methamidophos		A, TH, AW A, TH, AW	longer PHI
California ²	137,898	0.25-0.5	acephate permethrin diazinon dimethoate methamidophos oxydemeton-methyl endosulfan methomyl <u>B. thuringiensis</u>	0.5-1.0 0.1-0.2 0.25-0.5 0.25 0.5-1.0 0.25-0.5 0.25-0.5 0.45-0.9 Label	A, CL, M, TH CL, BAW A, CL, M, BAW A, TH A, CL, BAW A, TH A, TH, BAW, CL A, CL, BAW, TH CL	longer PHI
Colorado	200	0.25	No Alternatives			
Florida ³	1,300	0.25-0.5	acephate methamidophos methomyl endosulfan ethyl parathion methyl parathion permethrin flucythrinate <u>B. thuringiensis</u> carbaryl cyromazine dimethoate	1.0 0.5 0.9 0.7 0.2 1.0 0.2 0.1 Label 0.1-0.15 0.2 1.0	A, CL, TH A, CL, TH A, CEW, TH A, ICW A, CB, ICW, BL, M M CL, CEW, LM CL, SM CL CB, LB, ICW LM M	longer PHI
New Jersey	484	0.25	dimethoate malathion oxydemeton-methyl methomyl acephate carbaryl	0.25 1.25 0.5 0.45 0.998 2.0	A A A A A TPB	longer PHI

Table 23 Continued

				No Alternatives			longer PHI, expensive
New York	925	0.25		dimethoate	0.125	A	
				ethyl parathion	0.5	A	
Michigan ⁴	250	0.25-0.5		methomyl	0.9	A	
Texas	300	0.25-1.0		acephate	0.5-1.0	A, CL	
				diazinon	0.25-0.5	A	
				dimethoate	0.25	A	
				endosulfan	0.75-1.0	A, CL, ICW	
				methoxychlor	1.0-2.25	ICW	
				malathion	1.25-1.875	A, CL	
				methyl parathion	0.5-1.0	A, CL, ICW	
				oxydemeton-methyl	0.375-0.5	A	
				ethyl parathion	0.5	A, CL	
				<u>B. thuringiensis</u>	Label	CL, CEW, ICW	
				permethrin	0.1-0.2	CL, CEW, CW,	
						ICW	
				carbaryl	1.0-2.0	ICW, CEW	
				lindane	0.4-0.5	CW	
				trichlorfon	0.5-1.0	CW	
Wisconsin ⁵	600	0.125-0.5		acephate	0.75-1.0	A, CL	longer PHI
				demeton	0.25-0.5	A	
				diazinon	0.25-0.5	A	
				dimethoate	0.25	A	
				disulfoton	1.0-2.0	A	
				malathion	1.25	A, CL	
				oxydemeton-methyl	1.5-2.0	A	
				ethyl parathion	0.5	A, CL	
				endosulfan	0.75	CL	
				methomyl	0.2-0.9	CL	

¹A=aphids; AW=armyworm; BAW=beet armyworm; CEW=corn earworm; CL=cabbage looper; CW=cutworm; ICW=Imported cabbageworm; LB=Lygus bug; LM=leafminers; M=mites; TPB=tarnished plant bug; TH=thrips.

215-20% loss
310-15% loss
45% loss
510% loss

Because of its short residual activity and 2-days preharvest interval, the principal use of mevinphos on melon crops is for insect control during the harvest period. On watermelon the primary targets are aphids and rindworms. Generally, severe damage levels of these pests can be prevented with any of several registered alternatives to mevinphos, even though some have a longer preharvest interval, if the grower monitors the developing populations and anticipates the problems.

On cantaloupe and honeydew, the targets of mevinphos use vary from state to state: in Georgia, aphids and cucumber beetles on foliage and pickleworm on fruit (see cucumber section); in Texas, various foliage feeders including aphids, cabbage looper, cutworms, leafhopper, leafminer and mites. Acceptable control in these pest situations can be achieved with registered alternatives, as is the case with control of aphids on cantaloupe and watermelon in New Jersey. In California, the alternatives diazinon and dimethoate have longer preharvest intervals and are less effective; hence, yield benefits are associated with mevinphos use in California melons (Appendix Table II.18). In North Carolina, thrips are an early to mid-season problem on foliage and fruit damage is not direct; dimethoate and carbaryl are viable mevinphos alternatives for control, although repeated applications of carbaryl can stimulate outbreaks of mites and leafminer and improperly timed applications during bloom can cause extensive bee kills. It is these factors that account for the yield benefits attributed to mevinphos relative to alternatives as shown in Table 24.

Nonchemical controls are not dependable and cannot replace the use of insecticides on melons in the foreseeable future. Naturally-occurring biological controls (parasitoids, predators and pathogens) do not provide the level of control needed. Aphid-resistant cantaloupe varieties adapted to California have been developed which will provide acceptable aphid control. Unfortunately, aphid resistant varieties adapted to areas where mevinphos is used most extensively for aphid control are not available.

Although mevinphos is used on approximately 15,500 acres of melons annually for control of a variety of insect pests, a number of acceptable alternative insecticides are currently available to replace these uses with little decrease in the level of insect control. The continued availability of mevinphos, however, would provide a valuable tool for use in the management of insecticide resistance among the insect pests of melons (Table 24).

Okra

Only Florida reported the use of mevinphos on okra. Mevinphos is registered on okra for aphids, cutworms, corn earworm, green stink bug, mites and velvetbean caterpillar.

The rate of mevinphos use on okra ranges from 1/4 to 1/2 pint per acre for aphids and from 1/2 to 1 pint for the other pests on the label. The PHI is 2 days. And with okra harvested every other day under some conditions, this short preharvest interval is critical. The total acreage treated with

Table 24. Use of Mevinphos on Cantaloupe and Watermelons (average annual), 1982-1986.

State	Acres Treated	No. Appl.	Rate (lbs. a.i./A)	Pests	Alternatives	Limitations
California	10,752	1-4	0.25	aphids	dimethoate methomyl	less efficacious less efficacious
			0.5	mites	dimethoate	less efficacious
Florida	30 (cantaloupe)	1	0.25	aphids	dimethoate endosulfan methomyl	less efficacious less efficacious less efficacious
	6,000 watermelon	1			ethyl parathion	
			0.5	loopers	fenvalerate <u>Bacillus thuringiensis</u> endosulfan methomyl	resistance suspected resistance suspected controls only Lepidoptera low efficacy resistance suspected
			0.5	cutworms	carbaryl	limited efficacy and provides no aphid control
					fenvalerate methomyl	resistance suspected resistance suspected
			0.5	grasshoppers	carbaryl fenvalerate	resistance suspected
			0.5	leafhoppers	dimethoate carbaryl ethyl parathion fenthion	low efficacy
			0.5	leafminers	methamidophos naled	low efficacy
			0.5	<u>Lygus</u>	fenvalerate	

Table 24 Continued

Georgia watermelons	5,000	1 to 2	0.5	mites	dicofo ethyl parathion ethion	low efficacy				
			0.5							
			0.5	saltmarsh caterpillars	None					
				rindworms watermelon only	fenvalerate methamidophos	some resistance high cost				
					endosulfan <u>Bacillus</u> <u>thuringiensis</u>	high cost some resistance				
					methomyl					
				chinch bugs	ethyl parathion					
			0.5	aphids	endosulfan dimethoate	none poor control of cucumber beetle				
cantaloupe					carbaryl	poor control of aphids, kills bees				
					esfenvalerate endosulfan	poor control of aphids none				
500	1	0.5	aphids	endosulfan dimethoate						
				carbaryl	poor control of aphids, kills bees					
			pickleworm	esfenvalerate endosulfan	poor control of aphids none					
				endosulfan esfenvalerate	none					

81

Table 24 Continued

	1	0.25- 0.5	leafminers	azinphosmethyl carbaryl diazinon dimethoate ethion malathion methamidophos naled oxamyl parathion phosphamidon
		0.25- 0.5	mites	diazinon dicofol ethion malathion methamidophos naled oxydemeton-methyl parathion phosphamidon
New Jersey cantaloupe & watermelons	145	0.5	aphids	diazinon dimethoate oxydemeton-methyl methomyl endosulfan
North Carolina ¹ cantaloupe watermelon	2,000	0.25	thrips	carbaryl
				kills bees, stimulates mite and leafminer outbreaks
				Longer PHI
				dimethoate

¹Limitations of alternatives not severe.

mevinphos is 1,855 with an average of 4.5 sprays.

Survey results indicate a reduction in okra yield of 10-40% would result if mevinphos were not available. Okra is susceptible to aphids and stink bugs, especially after fruit has set. As okra is harvested many times after fruiting has begun, mevinphos with its short preharvest interval is far superior to any alternatives. None of the alternatives provide effective control during harvest.

Registered alternatives for aphids are malathion and parathion. Neither is as effective as mevinphos; parathion is very toxic to warm-blooded animals and beneficials and has a longer PHI (21 days). Malathion has a further restriction of not being registered to use after pods form.

Registered alternatives for Lepidoptera and stink bugs are malathion, parathion and carbaryl. Again malathion and parathion have longer PHI's. Carbaryl with a 0 day PHI is good, but corn earworm resistance is thought to exist (Table 25).

With none of the alternatives to mevinphos providing effective control of the broad spectrum of pests on okra, mevinphos has an important role in Florida. Likewise with the very minor nature of okra, from a pest management view it is important to retain the few registered insecticides available. The prospects of new registrations on okra seem very limited.

Onions

Mevinphos is used on both green and dry bulb onions in Florida, but only on the latter in Wisconsin, Texas and California. The primary pests were thrips, with cutworms also controlled in Florida. All states reported that mevinphos is both more effective than the registered alternatives and has a shorter preharvest interval (Table 26). Survey results indicate that if mevinphos were not available for thrips control, average annual yield loss of 10-20 percent in onions would have resulted (1982-1986).

Parsley

Only Florida reported the use of mevinphos on parsley, with 4-6 applications made to 400 acres for aphid control. Aphids are important pests of parsley because their feeding can render parsley leaves unmarketable. Mevinphos is the insecticide of choice for aphid control on parsley, primarily because of its effectiveness and short PHI. Light aphid infestations may be controlled with 0.5 lb ai/acre (PHI= 5 days) whereas high infestations or hard to kill species may require 1.0 lb ai/acre (PHI= 8 days).

The only registered alternative for aphid control on parsley is malathion which is both less effective and has a much longer PHI (21 days). The use of mevinphos provides a 20% savings in control costs as compared to the use of other known chemical or nonchemical alternatives. Further, survey

Table 25. Use of Mevinphos on Okra (average annual), 1982-1986.

State	Acres Treated	No. Appl.	Rate (lbs. a.i./A)	Pests	Alternatives	Limitations
Florida	1,855	4.5	0.05	aphids	malathion parathion	Not as effective Very toxic to warmblooded animals and beneficials, long PHI
				corn earworm	carbaryl	Not effective (suspect resistance)
				cutworms	parathion carbaryl	
				mites	malathion parathion	
				stink bugs velvetbean caterpillar	carbaryl	

Table 26. Use of Mevinphos on Onions (average annual), 1982-1986.

State	Acres treated	Pests	No. Appl.	Rate (lbs. a.i./A)	Alternatives	Limitations
Florida	750	thrips cutworms	6-8	1/2	methomyl malathion parathion	longer PHI
Wisconsin	300	thrips	2	1/2	parathion malathion diazinon	poor control
Texas	2000	thrips	-	1/2	diazinon parathion oxamyl malathion	-
California	4500	thrips	3	1/4	diazinon azinphosmethyl methomyl	longer PHI

results indicate that if mevinphos were no longer available for aphid control, parsley yields would drop an estimated 10 to 20%.

In the absence of suitable alternatives, it is important to maintain the availability of mevinphos for insect control in parsley.

Peas, green and southern

Mevinphos was used on some acreage in Wisconsin for control of pea aphids and cutworms. The alternative insecticides, dimethoate and fenvalerate do not control cutworms (dimethoate) or pea aphids (fenvalerate).

Potatoes

Three states reported the use of one application of mevinphos on potatoes for late season control of aphids, leafhoppers, and mites. Alternative insecticides labeled for control of potato insect pests include methamidophos, methomyl, parathion, carbaryl, dimethoate, and fenvalerate. Carbaryl is less effective against the insect pests, does not control mites and is highly toxic to bees. Methamidophos is more expensive and is toxic to beneficials (Table 28).

Spinach

Mevinphos is registered for use on spinach for control of aphids, cabbage looper, imported cabbageworm, grasshoppers, leafhoppers, mites, adult dipterous leafminers, climbing cutworms, saltmarsh caterpillar and false chinch bug. Registered use rates range from 0.125 to 0.25 lb ai/acre for aphids to 1.0 lb ai/acre for "hard to kill insects including aphids" (Phosdrin 4EC label) with a 4 day preharvest use restriction (Table 29).

Mevinphos is used for insect control on spinach in California, Florida, Maryland, New Jersey, New York, Oklahoma, Texas, and Virginia, with the most extensive use in New Jersey, New York, Oklahoma, Texas and California. Although in Florida and Texas, some use of mevinphos is directed against many or most of the insects listed on the Phosdrin 4EC label, the primary targets of use in all states are aphids. Aphid populations can develop rapidly. Early season infestations, if uncontrolled, can cause severe stunting or even death of the plants. Infestations which develop close to harvest are particularly serious because they constitute a source of insect contamination of the harvested crop. Whereas viable alternatives exist for control of early season aphid infestations (e.g. parathion, dimethoate, diazinon, disulfoton, azinphosmethyl, endosulfan and methyl parathion) none of the alternatives are considered to be effective for controlling aphid infestations close to harvest either because of limited efficacy or preharvest intervals that are too long. Of the remaining insect pests, loopers, cutworms, grasshoppers, imported cabbage worm and salt marsh caterpillars are chewing insects which

Table 27. Use of Mevinphos on Peas (green and southern) (average annual), 1982-1986.

State	Acres treated	Pests	No. Appl.	Rate (lbs. a.i./A)	Alternatives	Limitations
Wisconsin (green)	500	aphids climbing cutworms	1	1/4	fenvalerate dimethoate	poor aphid control
Tennessee (southern peas)	250	aphids	3	1/8-1/4	dimethoate disulfoton	

Table 28. Use of Mevinphos on Potatoes (average annual), 1982-1986.

State	Acres treated	Pests	No. Appl.	Rate (lbs. a.i./A)	Alternatives	Limitations
Idaho	100	green peach aphid	1	1/4	methamidophos endosulfan	none
Florida	658	aphids mites leafhoppers	1	1/4-1/2	methamidophos methomyl carbaryl fenvalerate	toxic to beneficials, expensive, moderate control
Texas	2000	aphids leafhoppers mites	1	1/8-1/4	dimethoate endosulfan	

Table 29. Use of Mevinphos on Spinach (average annual), 1982-1986.

State	Acres treated	No. Appl.	Rate (lb. ai/A)	Pests	Alternatives	Limitations
California	4,000	1-2	0.25	aphids	diazinon	longer PHI
					dimethoate	longer PHI
					azinphosmethyl	longer PHI (14 days)
Florida	125	1	0.5	cabbage- looper beet army- worm	methomyl	longer PHI
					azinphosmethyl	longer PHI
			0.5	aphids	naled	poor control
					ethyl parathion	poor control
					malathion	poor control
			0.5	cabbage- looper	fenvalerate	resistance suspected
					Bacillus	long PHI
					thuringiensis	expensive; controls
					ethyl parathion	only Lepidoptera
			0.5	cutworms	fenvalerate	long PHI
					carbaryl	long PHI
					methomyl	long PHI
			0.5	grass- hoppers	carbaryl	long PHI
					fenvalerate	long PHI
					malathion	poor control
			0.5	imported cabbage worm	fenvalerate	long PHI
					methomyl	long PHI
					Bacillus	
					thuringiensis	reduced control
					endosulfan	

Table 29 Continued

Florida	125	1	0.5	0.5	leaf-hoppers	dimethoate carbaryl parathion	reduced efficacy does not control aphids
			0.5		mites	ethyl parathion methyl parathion	
	125	1	0.5		saltmarsh caterpillar	methomyl parathion	resistance suspected; long PHI
Maryland	100	4-5	0.25		aphids	dimethoate	14 day PHI would lead to increased aphid contamination
New Jersey	597	1	0.25		aphids	diazinon dimethoate disulfoton	longer PHI
New York	600	1-2	0.12-0.25		aphids	none	no other materials with short enough PHI to eliminate insect contamination
Oklahoma	2,000	2	0.25		aphids	dimethoate ethyl parathion	poor control of Lepidoptera 14 day PHI
Texas	1,000	1	0.125-0.25		looper	ethyl parathion methomyl	14 day PHI reduced control
					aphids	azinphosmethyl diazinon dimethoate endosulfan malathion methyl parathion naled	

Table 29 Continued

Texas	1,000	1	0.125-0.5 0.125-0.5	cabbage-looper cabbage-looper	<u>Bacillus thuringiensis</u> methyl parathion naled ethyl parathion permethrin
			0.25-0.5	cutworm	diazinon permethrin
			0.25-0.5	leaf-hopper	carbaryl dimethoate methoxychlor methyl parathion ethyl parathion permethrin
			0.25-0.5	mites	azinphosmethyl methyl parathion ethyl parathion
Virginia	400	2-3	0.25	aphids/ leaf-miner	diazinon dimethoate disulfoton permethrin
					PHI too long can be used at planting only and does not provide full season control (leafminers only) not labeled for aphids

feed on the foliage. Heavy infestations early in the season can slow plant growth and even cause plant death whereas lighter infestations can severely reduce quality through feeding damage on harvested leaves. For this reason damage thresholds for these insects approach zero and infestations are prevented from becoming established. There are a number of equally effective alternatives for control of these pests on spinach, except in the unusual situation where a larval infestation exists close to harvest and must be controlled to prevent insect contamination of the marketable commodity.

Spider mite feeding on spinach results in a yellow stippling of the foliage which reduces the quality of the harvested produce. Mite infestations at the time of harvest thus must be controlled. Methyl and ethyl parathion are alternatives to mevinphos for mite control on spinach.

There are registered alternative insecticides which will control all of the important insect pests attacking commercially grown spinach in the US. The activity spectra and the longer preharvest intervals of the alternatives preclude their use as viable replacements for mevinphos as a "clean up" treatment close to harvest to prevent insect contamination of the marketable crop. Thus, yield benefits are associated with mevinphos use.

There are no effective alternatives to the use of insecticides for spinach. Parasitoids, predators, and pathogens cannot be depended upon to prevent damaging infestations of chewing insects from developing. Although parasitoids and fungal pathogens will often prevent aphid populations from reaching levels which damage the plant through direct feeding injury, aphids killed by these natural enemies become stuck to the spinach foliage and become a major source of insect contamination of the harvested crop. In essence these natural enemies constitute as severe a problem as the aphids and their presence must be prevented through efficient aphid control.

Squash

Mevinphos is registered for use on summer squash at 0.125 to 0.25 lb. ai/acre for control of aphids and at 0.25 to 0.50 lb ai/acre for control of cabbage looper, dipterous leafminer adult, leafhoppers, Lygus bugs, mites, false chinch bugs, salt marsh caterpillar, climbing cutworm and grasshoppers. The short preharvest interval (2 days) makes mevinphos attractive for use on summer squash during the period when the crop must be harvested approximately every three days.

Most mevinphos use on summer squash takes place in Florida and Georgia. It is used in Florida primarily during the harvest period for control of aphids, mites and Lygus. Aphid and mite populations develop on the foliage. Lygus feeding on developing fruits can lead to misshapen and aborted fruits, and, hence, reduced yield and quality. Although, a wide variety of other insect pests can be controlled on summer squash in Florida by mevinphos, other insecticides (Table 30), because of their longer residual activity, are preferred except during the harvest period, at which time the short preharvest interval and broad activity spectrum makes mevinphos the preferred insecticide.

Table 30. Use of Mevinphos on Squash (average annual), 1982-1986.

State	Acres treated	No. Appl.	Rate (lb. ai/A)	Pests	Alternatives	Limitations
Florida						
summer squash	1,500	1	0.25	aphids	ethyl parathion methomyl endosulfan	
			0.5	cabbage- looper	fenvalerate endosulfan <u>Bacillus</u> <u>thuringiensis</u>	
			0.5	cutworms	fenvalerate methomyl ethyl parathion	
			0.5	chinch bugs	ethyl parathion carbaryl	
			0.5	grass- hoppers	fenvalerate carbaryl	
			0.5	leaf- hoppers	naled parathion oxamyl	
			0.5	lygus	ethyl parathion fenvalerate	
			0.5	mites	dicofol ethion	
Florida						
summer squash	1,500	1	0.05	mites	ethyl parathion	
			0.5	saltmarsh caterpillar	methomyl	

Table 30 Continued

Georgia	3,000	2 to 3	0.5	aphids	endosulfan dimethoate	none poor control of cucumber beetle
			0.5	cucumber beetle	carbaryl endosulfan esfenvalerate	poor control on aphids; bee kills none poor control of aphids
			0.5	pickle- worm	endosulfan esfenvalerate	none poor control of aphids

In Georgia, mevinphos use on summer squash is directed against aphids and cucumber beetles, which feed on the foliage, and pickleworm which feeds on the fruit. Aphids and cucumber beetles, if properly controlled during the preharvest period, rarely develop damaging infestations during the harvest period. The pickleworm, because it infests the fruits, must be controlled from first bloom through harvest.

As with most vegetable crops, nonchemical controls and naturally occurring biological control cannot be depended upon to provide a reliable and acceptable level of insect control on summer squash. Insecticides are needed to achieve acceptable levels of control on summer squash. However, viable alternatives to mevinphos (e.g., endosulfan) are registered for use on summer squash. The loss of mevinphos would not impose severe constraints on summer squash production in either Florida or Georgia unless resistance to the alternatives were to develop in one or more of the key pest species. The yield benefits attributed to mevinphos use are associated with its use during the harvest period.

Tomatoes

Four states (Florida, New Jersey, Arkansas and Texas) reported using mevinphos on tomatoes for control of aphids, leafhoppers, leafminers, and mites. Approximately 7,000 acres received 1 to 2 treatments of mevinphos primarily for late season aphid control near harvest. Alternatives labeled for tomatoes are diazinon, dimethoate, methomyl, and dicofol. All have longer preharvest intervals than mevinphos. Based on the survey results the need for mevinphos usage on tomatoes did not appear significant except for late season outbreaks of aphid or mite populations during the harvest period (Table 31).

Leafy Greens - Turnip, Mustard and Kale

Seven states (Illinois, New Jersey, Oklahoma, South Carolina, Tennessee, Georgia and Louisiana) reported the use of mevinphos on leafy greens (turnip, mustard and kale) (Table 32). Mevinphos is registered on leafy greens for the following pests: aphids, cabbage looper (CL), imported cabbage worm, grasshoppers, leafhoppers, salt-marsh caterpillar, mites, climbing cutworms, leafminer (adult), lygus bugs and false chinch bugs. More recent use of mevinphos has been directed toward diamondback moth.

The rates of mevinphos range from 1/4 to 1/2 pint per acre for aphids and from 1/2 to 1 pint for the other pests on the label. The PHI is 3 days. The total acreage treated with mevinphos is 19,031 with an average of 2.5 applications per crop season.

Survey results indicate that yield decrease without mevinphos ranges from 0 to 50%. Some contamination by aphids and caterpillars occurs and is of occasional economic concern in the processing industry. Mevinphos is the best aphicide available for use on greens, the only material registered for mite control, the material of choice for diamondback moth and an excellent

Table 31. Use of Mevinphos on Tomatoes (average annual), 1982-1986.

State	Acres treated	Pests	No. Appl.	Rate (lbs. a.i./A)	Alternatives	Limitations
Florida	4500	aphids leafhoppers grasshoppers mites	1	1/4	dimethoate diazinon methomyl dicofol	longer PHI late season aphid control
New Jersey	1040	aphids mites	2	1/4	dimethoate diazinon methomyl dicofol	longer PHI
Arkansas	1000	mites	2	1/4-1/2	none	-
Texas	400	aphids leafhoppers leafminers	1	1/8-1/4	-	-

Table 32. Use of Mevinphos on Leafy Greens (Turnip, Mustard, Kale) (average annual), 1982-1986.

State	Acres Treated	No. Appl.	Rate (lbs. a.i./A)	Pests	Alternatives	Limitations
Georgia ¹	17,000	3.5	0.75	diamondback moth (DMB) aphids grasshoppers	<u>B. thuringiensis</u> fenvalerate methomyl endosulfan	No cleanup spray for DBM collards only not effective no DBM, long PHI
Louisiana ²	250	3.5	0.25 - 0.5	aphids DBM stinkbugs	malathion azinphosmethyl methyl parathion (Pennacap-M) naled B.T. fenvalerate permethrin	less residual limited Appl. PHI limited crop registrations (not mustard)
Oklahoma ³	800	2	0.25	aphids	naled dimethoate malathion parathion	not for aerial application long PHI not effective for CL long PHI long PHI
South Carolina ⁴	900	4	0.5	DBW imported cabbageworm aphids	endosulfan B.T. methomyl	Phosdrin okay 21 day PHI no control of DBM expensive no control of DBM 10 day PHI
Illinois ⁵	500	2	0.5 - 1.0	green peach aphid cabbage worms	dimethoate B. T.	Longer PHI worm control only
Tennessee ⁶	1,200	6	0.12 - 1.0 0.12 - 1.0	aphids caterpillars	dimethoate parathion	Longer PHI

Table 32 Continued

New Jersey	81	1	0.25	aphids mites	dimethoate carbaryl	mevinphos is only material registered for mite control
				ICW CL	<u>B. thuringiensis</u> methomyl	

- 1 Cleanup spray at harvest; reduction in quality if mevinphos is not used.
- 2 Little change in yield and no change in quality.
Where DBM is a problem mevinphos is an excellent alternative in a "rotating" schedule.
No material alone is effective.
- 3 Crop Rejection est. 5-15% - Loss.
Aphids are the major late season pest and create serious problems in contaminating the crop.
Clean up alternative is needed which cannot be applied by air.
Ground application limited due to irrigation and time constraints.
- 4 Yield loss 50%.
Decreased quality due to insect damage caused by DBM.
Only recommended material for DBM (essential).
- 5 Some contamination by insects.
Need not essential.
- 6 No yield effects.
Aphids present when processing (some concern)
Mevinphos is best aphicide on greens.

material in an insecticide rotation schedule and as a cleanup spray.

Registered alternatives for aphids are dimethoate, malathion, endosulfan, parathion, azinphosmethyl, and naled. They have less residual activity against aphid, have longer preharvest intervals than mevinphos, and none is registered for mites. Some of the alternatives cannot be applied by air. In addition they are expensive and in the case of diamondback moth, ineffective.

With no single, effective alternative superior to mevinphos for a broad spectrum of pests and with their serious limitations, mevinphos remains the product of choice in most situations.

Leafy Greens - Collards

Four states (North Carolina, South Carolina, Alabama and Florida) reported the use of mevinphos on collards (Table 33). Mevinphos is registered on collards for the following pests: aphids, cabbage looper, imported cabbage worm, grasshoppers, leafhoppers, salt-marsh caterpillar, mites, cutworms (climbing), leafminer (adult), Lygus bugs. More recent use of mevinphos has been directed against diamondback moth due to the occurrence of resistance to other registered insecticides. In this use pattern a tank mix of mevinphos and one of the Bacillus thuringiensis formulations has proven most effective in the management of resistant diamondback moth populations.

The rates of mevinphos range from 1/4 to 1/2 pt/acre for aphids, from 1/2 to 1 pt. for other pests and 1 qt/acre for hard to kill insects. The time limitation between last spray and harvest is from 2 to 3 days depending on the rate used. The total acreage treated with mevinphos is 3,781 with an average per state of 5.25 applications on collards.

Survey results indicate that yield decrease without mevinphos ranges from 5 to 50%. Aphids are the major late season pest and create a serious problem in contaminating the crop. The contamination may be direct (insects, cast skins), or indirect (honeydew, sooty mold, necrotic leaves, holes). Processed greens also may be contaminated with grasshoppers, leafminers and diamondback moth (DBM). Mevinphos is the single best material that is effective as a clean-up treatment near harvest. As a result of label limitations and more recently the development of diamondback moth resistance to several pesticides, mevinphos is the only effective material that will control diamondback moth in addition to managing a broad range of pests. Mevinphos has the advantage of having a short preharvest interval (3 days), of being registered for both ground and aerial application, of being relatively inexpensive, and of having a long history in broad spectrum activity as a clean-up treatment. Added value with mevinphos comes from its use in a tank mix with Bacillus thuringiensis (Dipel or Javelin) for the management of resistant diamondback moth populations throughout the southeast and its full season use in a rotating schedule and in an integrated pest management approach. For latest information as to DBM resistance refer to the regionalized cooperative study under the directions of Drs. Anthony M.

Table 33. Use of Mevinphos on Leafy Greens (Collards) (average annual), 1982-1986.

State	Acres Treated	No. Appl.	Rate (lbs. a.i./A)	Pests	Alternatives	Limitations
North Carolina ¹	2,000	8	0.5	diamondback moth (DBM) aphids	methomyl fenvalerate <u>B. thuringiensis</u>	Not effective where resistant populations occur. No aphid control with these materials PHI (methomyl) (10) Slow action
South Carolina ²	1,000	7	0.5	DBM	endosulfan fenvalerate <u>B. thuringiensis</u>	Endosulfan 21 days, PHI, expensive. No DBM control; toxic to fish
Alabama ³	400	5	0.5	DBM	methomyl <u>B. thuringiensis</u> <u>B. thuringiensis</u>	10 day PHI Less control
Florida ⁴	381	1	0.5	imported cabbageworm leafminer lygus mites salt-marsh caterpillar cabbage looper cutworms grasshoppers aphids	fenvalerate methomyl <u>B. thuringiensis</u> endosulfan dimethoate carbaryl parathion naled, diazinon, malathion	DBM resistance, highly toxic to fish DBM resistance expensive provides only moderate control not effective resistance suspected highly toxic to warm-blooded animals and beneficials does not provide effective control

¹Est. 10% loss in yield with alternatives. More where resistant populations exist. Severe cases results in nonmarketable greens.

²Yield 5%. Decreased quality due to DMB damage.

³25% yield loss due to DBM. Quality decreased greatly.

⁴5-10% yield loss
Aphids & LM before harvest results in reduced quality of collards.

Shelton (Cornell University) and Jeffrey Wyman (University of Wisconsin).

Mevinphos as a tank mix with Bacillus thuringiensis (B.T.) has received endorsement of most universities and has been the "stop-gap" measure adopted by growers where DBM resistance exists or is suspected.

Registered alternatives for aphids are naled, parathion, diazinon and malathion and provide inadequate aphid control and all have a longer preharvest interval than mevinphos (3 days at 1/4 to 1/2 pt/acre and 7 days at 1 qt/acre) i.e. naled (4 days), parathion (10 days), diazinon (10 days), and malathion (7 days).

Registered alternatives for control of Lepidoptera (cabbage looper, imported cabbage worm, diamondback moth, salt-marsh caterpillar, climbing cutworms, corn earworm, and velvet bean caterpillar) are fenvalerate, B.T., parathion, carbaryl, methomyl and endosulfan. The limitations of lack of effective and timely control exist for all of the above materials. Slow mode of action (B.T.); ineffectiveness against cabbage looper and diamondback moth by parathion, fenvalerate, carbaryl, methomyl and endosulfan; increased comparative cost of B.T., fenvalerate, carbaryl, methomyl and endosulfan; high toxicity of parathion and methomyl; danger of fish kills with fenvalerate and endosulfan; suspect or documented resistance by diamondback moth to fenvalerate and methomyl and the long preharvest interval between last spray and harvest of fenvalerate (7 days), parathion (10 days), carbaryl (14 days), methomyl (10 days), and endosulfan (21 days and can be used only once per season), individually and collectively make the use of alternatives unacceptable. Likewise, similar limitations exist for the other pests listed earlier. In particular the long preharvest interval for those materials listed earlier and for dimethoate (14 days).

There are no other effective alternatives to the use of insecticides in the management of pests on collards at present. Rotation, natural and biological control, resistant varieties, and new pesticide chemistry offer some hope but their realization and adopted use is neither practical nor likely to be commercially available in the foreseeable future. Hence, mevinphos remains the most viable approach in pest management on collards.

GREENHOUSE

Mevinphos (Phosdrin 4 EC) is registered for use on greenhouse-grown lettuce (including leaf, bibb and romaine). Pests controlled are aphids, corn earworms, climbing cutworms, dipterous leafminer adults, cabbage looper, imported cabbageworm, grasshoppers, mites, Lygus bugs, salt-marsh caterpillars, false chinch bugs and thrips. Rates to be applied are 0.625-0.155 lb. a.i./50,000 sq.ft. Pre-harvest interval is 10 days.

None of the states surveyed reported any greenhouse use of mevinphos.

AERIAL APPLICATION

The aerial application of mevinphos was reported in the Columbia Basin of Washington to alfalfa grown for seed production, in California, Nevada, Oregon and Pennsylvania to alfalfa grown for hay and in Florida to parsley and in Oklahoma to greens. The use of mevinphos on alfalfa grown for seed and for hay is discussed in the Field Crops section of this report and the use of mevinphos on parsley is considered in the Vegetable Crops section. Mevinphos was not applied by air to large acreages and was not considered as essential by any reporting state.

SEWAGE PLANTS, SYSTEMS AND BIOFILTERS

One state, Missouri, reported on our questionnaire that mevinphos is occasionally used to control flies which breed in the filters of sewage disposal plants.

If not controlled, these flies may build up to the point that their larvae and pupae clog filters, and the adults become a nuisance to plant employees and to persons living nearby. It was reported that over the period 1982-86, an average of 1-2 cities per year treated flies at their sewage plants with mevinphos. Because no other state reported the use of mevinphos for this purpose, either the situation is unique to Missouri or is not considered a problem in other areas, or the crop oriented respondents to our questionnaire were unaware of this use in their state.

When used for this purpose, mevinphos is applied 1-2 times per year, at a rate of 0.35 pounds per 100 gal. It is obviously applied only when the flies have become a problem or are expected to do so. The Missouri questionnaire indicated that there is no alternative insecticide. The obvious advantage of using mevinphos is that it has a broad spectrum of toxicity, yet it hydrolyses very rapidly, thus causing little downstream risk. It is not known whether Bacillus thuringiensis var. israelensis has been evaluated for these pests. B. thuringiensis would impact a more limited spectrum of organisms, but would persist further downstream.

ECONOMIC AND SOCIAL IMPACTS

Methodology and Data

Comparative static equilibrium analysis is used to demonstrate the benefits of mevinphos. The estimated impact of mevinphos use is determined by comparing yields, prices, and production with the expected use of mevinphos's alternatives.

The economic benefits of mevinphos are the higher yields attained with its use in addition to its lower costs relative to many insecticides for controlling pests on major vegetable and fruit crops. In addition, it is important in a few major field crops.

The benefits of mevinphos were calculated as the sum of reduced revenues plus added costs that would result from the withdrawal of mevinphos. Presumably, growers are now using mevinphos because it is more profitable than alternative pesticides or control methods. If mevinphos were withdrawn from use, a grower could be expected to experience the following results:

- 1) reduced yield (from decreased quantity or quality),
- 2) increased costs associated with the use of a more expensive alternative treatment, and/or
- 3) increased market prices (from aggregate reductions in crop supplies).

The methodology used is derived from estimating

$$B = \sum_{i=1}^n (R_0 - R_1 + C_1 - C_0)$$

where:

- B = Average annual net benefits of mevinphos,
- R₀ = Average annual revenue in the base period (1982-86) with the use of mevinphos,
- R₁ = Average annual revenue in the future without mevinphos,
- C₁ = Average annual costs of control in the future without mevinphos, and
- C₀ = Average annual costs of control in the base period (1982-86),
- i = a site in a particular state, and
- n = total number of sites.

Benefits were summarized by state, by region, and by site to determine the impact of the withdrawal of mevinphos.

The methodology assumes the following:

- a) A parallel shift in the supply of a commodity due to a change in yield results in a change in the quantity of that commodity demanded, holding other factors constant. The Economic Analysis Simulator (EAS) developed by Walter Ferguson (1989) of USDA Economic Research Service is used to generate the results.
- b) The yield benefit of using mevinphos instead of its alternatives is known for each commodity. Survey responses are used to estimate the yield benefit of mevinphos by asking crop production experts for the yield loss from using mevinphos's alternatives. The percentage shift in aggregate supply is an average, weighted by mevinphos-treated and non-mevinphos-treated production. Non-mevinphos-treated production is the total U.S. production minus the product of mevinphos-treated acreage and yield.
- c) The price response following a yield-induced shift in supply is estimated using a price flexibility of demand. Farm-level flexibilities were obtained from econometric analyses of 1954-1988 data, including own-price, per capita consumption, price of substitutes, and selected other factors (summarized in APPENDIX IV).
- d) The cost change from using mevinphos's alternatives is additional to any yield, price, and value of production changes for a commodity. Survey responses are used to estimate the cost changes associated with mevinphos's alternatives.

Quantitative Use of Mevinphos

The total amount of mevinphos used annually in the United States during 1982-86 averaged 756,815 pounds active ingredient (a.i.) (Table 34). The total acreage treated averaged 666,723 acres. The treated acreage received an average of 1.14 pounds a.i. per acre. The total annual treatment cost of mevinphos use averaged \$13,612,000, about \$20.42 per acre. This total includes application and chemical costs. The total cost of alternatives to mevinphos treatment averaged \$15,491,000 or about \$23.23 per acre.

Vegetables were treated with an average of 594,777 pounds a.i. of mevinphos annually during 1982-86. This amounts to 78.6 percent of the total mevinphos applied (Table 34). Fruit crops (mainly strawberry) received 103,977 pounds a.i. or 13.7 percent of the total. Field crops (mainly alfalfa seed) received 58,061 pounds a.i. or 7.7 percent of the total.

Vegetable acreage treated with mevinphos was 547,067 acres, 82.1 percent of the total acreage treated. Total annual cost of mevinphos treatment on vegetables was \$11,309,000, an 83.1 percent share of the total costs of treatment. The per acre cost of treatment with mevinphos averaged \$20.67 for vegetables, \$18.44 for field crops, and \$22.39 for fruit crops. The per acre

Table 34. Survey responses: Acreage and acreage treated with mevinphos, mevinphos applied, and control costs with and without mevinphos available, by crop type and region (average annual), 1982-86.

Crop Type or Region ¹	Acreage		Mevinphos applied ² (pounds a.i.)	Control Costs	
	reported	treated		Current (—1,000 dollars—)	Alternative ³
Vegetables	1,264,702	547,067	594,777	11,309	12,708
Field crops	885,107	95,320	58,061	1,758	947
Fruits	779,482	24,336	103,977	545	1,836
Northeast	117,780	13,964	7,601	236	221
North Central	204,180	9,900	5,388	166	115
South	592,261	107,035	223,476	900	4,946
South Central	746,380	79,743	75,678	815	2,061
West	1,229,096	445,400	435,104	11,348	8,029
Northwest	39,594	10,682	9,568	147	119
Total	2,929,291	666,723	756,815	13,612	15,491

¹For regional definitions, see Note 1 at end of section.

²Total pounds active ingredient (a.i.) applied equals average number of applications times average application rate times acres treated.

³Acreage treated with mevinphos times average cost of chemical and/or nonchemical control alternatives.

cost of treatment with alternatives to mevinphos averaged \$23.23 for vegetables, \$9.93 for field crops, and \$75.44 for fruit crops.

The West region averaged the largest share of mevinphos use, 57.5 percent. The South region was next with 29.5 percent. The others accounted for 13 percent (Table 34). In terms of acreage treated, the West region had 66.8 percent of the total treated, the South had 16.1 percent, the others had 17.1 percent. In terms of total treatment costs, the West had 83.3 percent of the total, the South had 6.6 percent, and the others had 10.1 percent. Of the total cost of alternatives to mevinphos, the West had 51.8 percent, the South had 31.9 percent, and the others had 16.3 percent.

In terms of the total acreage treated or mevinphos applied, lettuce is the most important crop, followed by alfalfa seed, broccoli, cabbage, greens, and strawberry (Table 35). Other important crops are cauliflower, celery, and muskmelons. For crops such as artichoke, carrot, southern pea, okra, and spinach which do not account for a large proportion of either acreage or mevinphos used, the use of mevinphos is important, as more than half their acreage reported was treated with mevinphos.

The largest shares of the total cost of using mevinphos are in lettuce (41.2 percent), alfalfa seed (12.7 percent), broccoli (11.1 percent), and cauliflower, cabbage, and celery (total 14.7 percent) (Table 35). The largest shares of total cost of alternatives to mevinphos use are in lettuce (25.0 percent), cabbage (16.4 percent), greens (11.5 percent), strawberry (10.2 percent), broccoli (8.8 percent) and alfalfa seed (6.0 percent).

The net cost advantage of using mevinphos over its alternatives on all crops is \$1,879,000, but the difference varies widely over individual crops. Higher total costs of using alternatives to mevinphos occur mainly in cabbage, greens, and strawberry. Lower total costs of using alternatives to mevinphos occur mainly in lettuce, where the yield benefit of 7,501,000 hundredweight (cwt.) offsets the higher cost of using mevinphos. Alternative costs are lower in alfalfa seed production also, but the yield benefit is offsetting.

The estimated yield benefit of mevinphos ranges widely over different crops (Table 36). In descending order, the crops with high relative yield benefits on treated acreage are artichoke (45.0 percent), strawberry (44.3 percent), and cauliflower (40.3 percent). Crops with a yield benefit in the range of 20 percent to 30 percent include celery, onion, and okra. Crops with a yield benefit in the range of 10 to 20 percent include broccoli, Brussels sprouts, cabbage, sweet corn, muskmelons, lettuce, parsley, and spinach.

The last column of Table 36 lists the aggregate yield benefit of mevinphos on the basis of total U.S. acreage (treated and untreated with mevinphos). Nine crops have a significant aggregate yield benefit from mevinphos use: broccoli (6.4 percent), cabbage (8.9 percent), cauliflower (20.4 percent), celery (8.0 percent), greens (3.8 percent), muskmelons (3.1 percent), lettuce (12.1 percent), strawberry (10.2 percent), and spinach (14.2 percent). In the markets for these crops with significant yield

Table 35. Survey responses: Acreage treated with mevinphos, mevinphos applied, mevinphos's yield benefit, and control costs with and without mevinphos available (average annual), 1982-86.

Crop	Acreage treated	Mevinphos applied ¹ (pounds)	Yield benefit ² (1,000 cwt.)	Control Costs	
				Current (—1,000 dollars—)	Alternative ³
Alfalfa, seed	92,800	56,801	12	1,740	928
Apple	4,740	3,740	8	134	137
Broccoli	61,892	71,764	689	1,510	1,368
Brussels sprouts	3,000	1,710	28	55	30
Cabbage	46,310	63,597	2,367	606	2,537
Pea, southern	250	141	---	2	7
Carrot	3,535	1,326	14	25	32
Cauliflower	34,400	35,445	1,401	802	296
Celery	15,743	19,143	1,478	595	535
Greens	24,985	69,667	198	180	1,775
Corn, field	820	410	---	6	12
Sweet corn	12,712	7,060	145	213	106
Cucumber	10,126	3,337	---	74	199
Eggplant	401	160	---	4	8
Grape	11,419	6,509	50	210	114
Muskmelons	38,463	17,271	622	529	527
Lettuce	209,500	241,748	7,501	5,610	3,867
Onion	5,550	6,300	673	179	77
Parsley	400	1,000	9	3	11
Pea, english	500	125	---	8	3
Pepper	17,769	5,284	24	145	271
Potato	658	247	---	7	17
Sorghum	1,700	850	---	12	7
Squash	4,650	4,369	14	40	88
Strawberry	8,177	93,729	1,025	201	1,585
Tomato	6,922	3,543	83	77	99
Turnip	200	100	---	2	1
Artichoke	10,000	5,700	344	184	100
Bean, lima	23	17	<1	<1	1
Bean, snap	10,169	6,188	2	102	264
Okra	1,855	4,174	19	17	72
Spinach	13,910	16,040	187	204	134
Watermelon	13,145	9,323	26	139	283
Total	666,723	756,815		13,612	15,491

¹Total pounds active ingredient (a.i.) applied equals average number of applications times average application rate times acres treated.

²Yield benefit equals percent change in yield times production treated with mevinphos.

³Acreage treated with mevinphos times average cost of chemical and/or nonchemical control alternatives.

Table 36. U.S. crop production: Mevinphos-treated farm production and yield benefit, U.S. aggregate farm production, farm price, and yield benefit, by crop (average annual), 1982-86.

Crop	Mevinphos-treated		U. S. aggregate		
	Farm production (1,000 cwt.)	Yield benefit (percent)	Farm production (1,000 cwt.)	Farm price (dollars/cwt.)	Yield benefit (percent)
Alfalfa seed	805	1.5	1,651	97.13	0.7
Apple	630	1.3	80,827	11.44	0.0
Broccoli ¹	5,642	12.2	10,767	22.86	6.4
Brussels sprouts	285	10.0	---	---	---
Cabbage ¹	12,798	18.5	26,732	7.67	8.9
Pea, southern	8	---	---	---	---
Carrot	387	3.7	23,450	9.49	0.1
Cauliflower ¹	3,479	40.3	6,875	25.41	20.4
Celery ¹	5,419	27.3	18,443	12.89	8.0
Greens ¹	3,736	5.3	5,262	22.79	3.8
Corn, field	195	0.0	4,167,561	4.23	0.0
Sweet corn	1,357	10.7	15,131	12.89	1.0
Cucumber	1,106	0.0	---	---	---
Eggplant	81	0.0	---	---	---
Grape	832	6.0	14,279	20.15	0.0
Muskmelons ¹	5,574	11.2	19,895	13.07	3.1
Lettuce ¹	62,097	12.1	62,031	11.57	12.1
Onion	2,856	23.6	42,436	9.13	1.6
Parsley	62	15.0	---	---	---
Pea, english	15	---	---	---	---
Pepper	1,868	1.3	5,729	22.12	0.4
Potato	150	0.0	364,055	20.44	0.0
Sorghum, forage	NA	7.5	NA	NA	0.2
Sorghum, grain	20	2.5	406,537	3.94	0.0
Squash	649	2.2	---	---	---
Strawberry ¹	2,314	44.3	10,097	45.77	10.2
Tomato	3,575	2.3	28,592	24.01	0.3
Turnip	NA	---	---	---	---
Artichokes	763	45.0	---	---	---
Bean, lima	1	2.5	---	---	---
Bean, snap	372	0.4	---	---	---
Okra	63	30.0	---	---	---
Spinach ¹	1,324	14.2	4,474	25.54	14.2
Watermelon	1,497	1.8	24,499	5.69	0.1
Total	120,336	14.1			

¹See note 2 at end of text.

SOURCES: For mevinphos-treated production and yield effects--1988-89 survey of states. For U.S. aggregate production and price--U.S. Department of Commerce, 1982 Census of Agriculture, National Summary; USDA, National Agricultural Statistics Service, Crop Production, annual issues; USDA, National Agricultural Statistics Service, Crop Values, annual issues.

effects, the benefits of mevinphos include lower prices. For the other crops, with less than 2 percent yield benefit over total U.S. production, the price effects are considered negligible and the total benefits amount to a cost differential to the growers.

Yield, Price, and Crop Value Benefits

The yield benefit of mevinphos with respect to price is assumed to be a result of higher supplies and lower price to consumers. That is, considering total production, the higher yield causes price to be lower, holding other factors constant. For producers of a commodity exhibiting relatively stable (or inflexible) prices, higher production means higher total crop value. (For a recent example of similar considerations, see Barse, Ferguson, and Seem.)

The price characteristics of selected vegetables are given for 1954-1988 in Table 37. The minimum, maximum, and mean price vary widely over different crops; the coefficient of variation (the ratio of the standard deviation to the mean) is similar among crops. The price flexibilities of demand were calculated from price-dependent equations. Farm-level own-price was regressed on per capita use and the farm-level price of all other vegetables, using Ordinary Least Squares regression. The per capita use quantities were adjusted for carryover stocks and net exports, and all prices were deflated by the Gross National Product deflator (1982=100). For all items, other adjustments were made. For example, utilization differences, consumer income, and binary variables were included where necessary. (APPENDIX IV)

The average value of fresh-market vegetables, regressed on per capita use and deflated U.S. disposable personal income, exhibits price-flexible demand at the farm level (Table 37). The individual items, though, exhibited price-inflexible demand. This is a likely result for several reasons. First, individual items may substitute for one another, compared to the aggregate group which has fewer substitutes. Second, the quantities of individual items are all less than the aggregate group, which lowers the flexibility estimate, *ceteris paribus*. Third, the vegetables with the lowest flexibilities are of a type including cauliflower and broccoli that may have more substitutes than a type including lettuce which is consumed regularly in combinations with other foods.

Selected items of interest from the output generated by Economic Analysis Simulator are summarized in Table 38. The estimates are based on acreage in production, proportion of production treated with mevinphos, changes in yield, farm-level price, and price flexibility of demand. Only those crops with a yield benefit from mevinphos large enough to affect the U.S. domestic market are included. For a comparison of production and crop value with and without the availability of mevinphos, percentage changes are presented in Table 39.

Table 37. U.S. price characteristics for selected crops (average annual), 1954-88.

Crop ²	Farm-level price ¹			C.V.	Flexibility ³
	Min.	Max.	Mean		
	(—dollars/cwt.—)				
Broccoli ⁴	7.36	25.23	13.64	0.45	-0.49
Cabbage ⁴	1.56	8.63	4.12	0.52	-0.82
Cauliflower	5.90	26.16	13.85	0.54	-0.21
Celery ⁴	3.11	13.68	6.90	0.46	-0.67
Greens ⁴	---	---	---	---	-0.82
Muskmelons ⁴	3.85	14.09	7.04	0.42	-0.70
Lettuce	3.28	15.19	7.10	0.48	-0.80
Strawberry	14.17	49.41	28.76	0.39	-0.53
Spinach ⁴	5.69	27.36	12.06	0.52	-0.19
Fresh-market vegetables ⁵	4.03	15.55	8.08	0.48	-2.45

¹Min.=minimum, Max.=maximum, C.V.=coefficient of variation, flexibility of farm-level price with respect to quantity demanded.

²Fresh-market (dual-purpose uses for broccoli, cauliflower, and strawberry).

³Defined as the percentage change in price per one-percent change in quantity used. Estimated with 1982-86 average quantity and predicted price.

⁴Estimated from 1954-81 data only. Due to lack of data, greens given the spinach flexibility. Muskmelons are cantaloupe in this table.

⁵Includes broccoli, cauliflower, celery, honeydew melons, lettuce, carrots, sweet corn, onions, tomatoes.

SOURCE: Computed from USDA, Agricultural Statistics, Annual issues

Table 38. U.S. price, production, and value of production for selected crops resulting from the unavailability of mevinphos.

Crop	U.S. Farm-Level		
	Price (\$/cwt.)	Production (1,000 cwt.)	Value (1,000 dollars)
Broccoli	23.24	10,406	241,789
Cabbage	7.94	25,593	203,157
Cauliflower	25.96	6,165	160,050
Celery	13.09	18,009	235,796
Greens	23.29	5,120	119,267
Muskmelons	13.15	19,722	259,336
Lettuce	12.69	54,526	691,928
Strawberry	46.34	9,861	456,929
Spinach	25.74	4,286	110,338

SOURCE: Calculated from Economic Analysis Simulator (Ferguson)

Table 39. Summary of economic impact on U.S. crop producers resulting from the unavailability of mevinphos, as compared to 1982-86 average annual, by crop.

Crop	Farm-level		Farm value		Control costs (1,000 dollars)
	Production (percent)	Price (percent)	Absolute (1,000 dollars)	Relative (percent)	
Alfalfa, seed	-	-	-	-	-812
Apple	-	-	-	-	+3
Broccoli	-3	+2	-4,345	-2	-142
Brussels sprouts	-	-	-	-	-25
Cabbage	-4	+4	-1,877	-1	+1,931
Pea, southern	-	-	-	-	+5
Carrot	-	-	-	-	+7
Cauliflower	-10	+2	-14,644	-8	-506
Celery	-2	+2	-1,934	-1	-60
Greens	-3	+2	-654	-	+1,595
Corn, field	-	-	-	-	+6
Sweet corn	-	-	-	-	-107
Cucumber	-	-	-	-	+125
Eggplant	-	-	-	-	+4
Grape	-	-	-	-	-96
Muskmelons	-1	+1	-692	-	-2
Lettuce	-12	+10	-25,771	-4	-1,743
Onion	-	-	-	-	-102
Parsley	-	-	-	-	8
Pea, english	-	-	-	-	-5
Pepper	-	-	-	-	+126
Potato	-	-	-	-	+10
Sorghum	-	-	-	-	-5
Squash	-	-	-	-	+48
Strawberry	-2	+1	-5,211	-1	+1,384
Tomato	-	-	-	-	+22
Turnip	-	-	-	-	-1
Artichoke	-	-	-	-	-84
Bean, lima	-	-	-	-	+1
Bean, snap	-	-	-	-	+162
Okra	-	-	-	-	+55
Spinach	-4	+1	-3,928	-3	-70
Watermelon	-	-	-	-	+144
Total			-59,056	-2	+1,879

At the farm level, opposite changes in production and price are nearly offsetting in cabbage, celery, greens, muskmelons, and strawberry. Due to the estimated inflexibility of prices for broccoli, spinach, and especially cauliflower, decreases in U.S. production are not as nearly offset by increases in price. Lettuce price would increase the most because its yield benefit from mevinphos is the greatest and its demand is relatively price responsive. Therefore, the yield benefit of mevinphos is proportionately greater on these latter four crops compared to the others. The benefit of mevinphos use, based on changes in crop value, totals \$59.1 million. The largest benefit comes from use on lettuce, followed by cauliflower, strawberry, broccoli, spinach, and celery.

In addition to the benefit of mevinphos on crop value, the net change in production costs for mevinphos's alternatives amounts to \$1.9 million (Table 39, column 4). The major cost increases of substituting chemical and other means to control pests previously controlled by mevinphos would occur on cabbage (+\$1.9 million), greens (+\$1.6 million), and strawberry (\$1.3 million). Offsetting reductions in costs for substituting mevinphos's alternatives could occur in lettuce (-\$1.7 million), alfalfa seed (-\$0.8 million), and cauliflower (-\$0.5 million).

The average annual net benefit of mevinphos in U.S. crop production, adding the change in total crop value to the net change in control costs, is \$60.9 million or about \$91 per acre treated. This estimate of total benefit is sensitive to different estimates of yield change and price flexibility. For example, a lower absolute value for price flexibility produces a higher value of any yield benefit from mevinphos use. Also, the net difference in mevinphos's alternative control costs are assumed to be borne entirely by the producer and not passed on to the consumer. The availability of mevinphos is likely to be a risk reducing (and thus cost reducing) opportunity for producers, suggesting an unmeasured benefit. Finally, for producers and farm labor, mevinphos may pose other risks which are not taken into account here.

Consumer Benefits

The benefits to consumers of a product treated with mevinphos are assumed to be measured by the difference in retail price that would be paid for the product treated instead with mevinphos's alternatives. Based on the farm-level yield and price effects of mevinphos (Table 39), assuming mevinphos is unavailable and assuming the price change is passed on to the retail level of distribution, consumers benefit from mevinphos use on broccoli, cabbage, cauliflower, celery, greens, muskmelons, lettuce, strawberry, and spinach.

Annual average retail prices are available for comparison with farm-level prices for several major fresh-market vegetables (Economic Research Service). The correlation coefficients estimated from 1970-1988 data are high and statistically significant for carrot (0.98), celery (0.97), lettuce (0.97), onion (0.85), and tomato (0.98). On average, during 1982-86 a one percent change in the farm-level price resulted in a 0.96 percent change in the retail price of celery and a 1.2 percent change in the retail price of lettuce. Assuming similar relationships between farm-level and retail prices

for broccoli, cabbage, cauliflower, greens, muskmelons, strawberry, and spinach, the yield benefit of mevinphos on these crops is likely to be lower prices also. The retail price benefit in percentage terms is likely to be on the order of estimates listed in Table 39 (column 2).

The consumers' total benefit from mevinphos use results from the lower prices paid for total production caused by the lower marginal value of the increased yield. Instead of using retail prices, which are unavailable for most of the items affected, farm-level prices and quantities are used to calculate the consumer's benefit from mevinphos. The product of the change in price times the total production of crops whose yield is increased by using mevinphos equals \$94 million, comprising broccoli (\$4 million), cabbage (\$7 million), cauliflower (\$3.6 million), celery (\$3.7 million), greens (\$2.6 million), muskmelons (\$1.6 million), lettuce (\$65 million), strawberry (\$5.6 million), and spinach (\$0.9 million).

The total benefit of mevinphos use is the sum of the increased production costs expected with mevinphos's alternatives (\$1.9 million), the higher farm value of production (\$59 million), and the higher consumer value of production (\$94 million). The total benefit of mevinphos use is \$155 million--61 percent of which is received by the consumer of the farm product.

Of course, in the event that mevinphos is unavailable, higher prices will put pressure on the market to adjust by increasing supply, possibly through increasing imports or through changes in production practices. The present analysis does not attempt to estimate market adjustments; it is intended to indicate where pressure for adjustment is more likely to occur.

Macroeconomic Effects

The commodities benefitting from mevinphos use occupy a relatively minor share of U.S. gross national product and require relatively minor shares of U.S. labor and capital. Also, the price and income impacts of mevinphos unavailability would likely cause negligible changes in inflation and interest rates.

Social/Community Impacts

Employment impacts in local areas where particular crops are important might occur. These could result from one or both of the following: (1) a crop may no longer be grown in specific local areas if mevinphos is unavailable; and (2) slowdown in packing houses and with shippers in cases where the rejection rate for produce due to insect damage becomes very high or packing houses shut down due to the loss of a crop.

These are very important events to the affected individuals and communities, in the short run, but are fairly minor for individual states or nationally. For the most part, crops on which mevinphos is used (i.e. fresh produce) are produced in areas where soil and climate are favorable for a wide range of crops. Therefore, there are usually alternative uses for land and labor.

Notes

1. Regions are defined as:

Northeast: Connecticut, Delaware, Maryland, Massachusetts, Maine, New Hampshire, New Jersey, New York, Pennsylvania, Rhode Island, Vermont

North Central: Iowa, Illinois, Indiana, Michigan, Missouri, Minnesota, Ohio, Wisconsin

South: Alabama, Arkansas, Florida, Georgia, Kentucky, Louisiana, Mississippi, North Carolina, South Carolina, Tennessee, Virginia, West Virginia

South Central: Colorado, Kansas, Montana, North Dakota, Nebraska, Oklahoma, South Dakota, Texas, Wyoming

West: Arizona, California, New Mexico, Nevada, Utah

Northwest: Idaho, Oregon, Washington

Other: Alaska, Hawaii, Guam, Puerto Rico, Virgin Islands, District of Columbia

2. U.S. aggregate farm production was estimated for cabbage, greens (collards, kale, mustard greens, turnip greens), muskmelons (cantaloupe and honeydew), and spinach by multiplying the 1982 Census of Agriculture estimate of 1982 acreage times the average yield calculated from mevinphos respondents. For broccoli, cauliflower, lettuce, and strawberry, USDA-estimated 1984-86 average production was used. For celery, USDA-estimated 1982-86 average production was used.

EXPOSURE CONSIDERATIONS AND THEIR RELATIONSHIP TO RISK ANALYSIS.

In general, our knowledge of the exposure and environmental risks from the use of mevinphos are incomplete (EPA 1988). Nevertheless, we can conclude some general trends based upon our knowledge of its general toxicity and residual longevity (Matsumura 1985).

Mevinphos is a highly toxic, broad spectrum cholinesterase inhibitor. It is highly toxic to a broad range of beneficial insects, birds, fish, and mammals. Most animals in or adjacent to an area to which mevinphos is applied are at risk. The following acute toxicity dosages have been reported: sharp-tailed grouse, 1.34 mg/kg dermal; mallard duck, 4.63 mg/kg dermal, 1991 ppm oral; Japanese quail, 236 ppm oral; rainbow trout, 11.9 ppb 96-hr; bluegill sunfish, 22.5 ppb 96-hr; sheepshead minnow, 640 ppb LC₅₀; Daphnia pulex, 0.18 ppb; and Pteronarcys californica, 5.0 ppb (EPA 1988). Nevertheless, no documented field kills of wildlife have been noted (EPA 1988). This is probably because mevinphos is rapidly hydrolyzed in aqueous solution (Hayes 1982). Mevinphos is one of the preferred insecticides for use on alfalfa grown for seed. On this crop the alfalfa leafcutting bee and the alkali bee are essential pollinators and must be carefully conserved. Insecticide treatments are applied exclusively at night when the pollinators have returned to their nests. The short residual properties of mevinphos reduces to acceptable levels the hazards to the bees working the treated crop the next day (Alfalfa Seed Industry 1989). The available information on the impact of mevinphos on non-target organisms may be summarized as: its acute toxicity to exposed organisms is very high, but its residual effect appears to be slight.

The physiological effects of mevinphos on rats, human volunteers, and accounts of accidents are included in an earlier section (Pesticide Characteristics). More details are reported by Hayes (1982). Poisoning reports from California indicate that mevinphos was among the top five pesticides causing occupational poisoning in that state from 1981-85. These were primarily from dermal exposure, with some from inhalation (EPA 1988). Thus, extreme caution must be exercised when working with mevinphos or working in an area where mevinphos is being applied. On the other hand, as with non-target organisms, a relatively short pre-harvest interval and a short treated field reentry period are probably justified, based on the rapid rate of hydrolysis of mevinphos residues.

Data currently available are insufficient to characterize mevinphos's leaching potential for contamination of ground water, but they indicate that its residues are highly mobile in sandy loam, silt loam, loam, and clay loam soils (EPA 1988). This is not surprising since technical mevinphos is miscible in water (Hayes 1982). Nevertheless, its water contamination potential is likely to be small because of its susceptibility to hydrolysis.

Because of the extremely high level of toxicity of mevinphos, it is crucial that label instructions be followed and that persons using this chemical be fully instructed in the appropriate safety measures. The acute toxicity hazard is such that mevinphos should be used only when other less hazardous alternative insecticides are not available. If used properly and

cautiously, residue and toxicity problems should be minimal. The residue hazard is probably not a major factor.

RECOMMENDATIONS

The recommendations contained herein are based on the assumption that the alternatives (pesticides) listed for specific site/pest/crop use remain viable. Loss of registration or availability of these alternatives to mevinphos would require a re-examination of recommendations listed below.

Based upon the results of the benefits analysis the continued registration of mevinphos on the following sites is considered essential:

- a. Strawberry - because the short preharvest interval allows for effective control of insect pests during a long harvest period characterized by frequent harvests at short intervals.
- b. Alfalfa for seed - because the short period of residual toxicity to pollinators uniquely allows for insect pest control while protecting essential pollinators.
- c. Artichoke, Beans (pole), Broccoli, Cabbage, Cauliflower, Celery, Lettuce, Okra, Onions, Parsley, Spinach, Leafy Greens (including Collards) - because the short preharvest interval of mevinphos on these crops uniquely provides for a broad spectrum of insect control close to harvest. In these crops this is essential to avoid insect damage to and contamination of harvested produce.

The economic analyses revealed significant economic benefits associated with mevinphos use on all of the crops listed above except artichoke, beans, okra, onions and parsley. For these latter, however, the pest control requirements are such that none of the available alternatives constitute fully acceptable alternative in all production areas.

The continued availability of mevinphos on commodities other than those listed above contributes 1) to grower ability to respond to rapid pest outbreaks, especially close to harvest and 2) to the diversity of insecticides available to counteract the development of resistance.

According to questionnaire response, mevinphos has only limited usage on the following commodities:

Field Corn
Popcorn
Sorghum (grain and forage)
Peaches
Pears
Plums
Sour cherries
Oranges
Lemons
Grapefruit
Raspberries
Walnuts

Cited usage in this report and the recommendations voiced by the mevinphos assessment team indicates that further or future support for use in mevinphos should be addressed on a commodity by community basis and contingency of resistance alternative insecticide.

Mevinphos is highly toxic and may enter the body by ingestion or by dermal contact. As such, mevinphos is classified as a Category I pesticide and the risk of acute poisoning is documented. As a result it is imperative that industry representatives continue to explore safer formulation or safer means of handling and applying mevinphos. In addition, extension and pesticide specialists should continue to instruct and train growers, farm managers, applicators and field labor on the safe handling and use of mevinphos to reduce and avoid occupational poisoning. Efforts to obtain additional base data on occupational poisoning should be documented and validated as to misuse and to accidental or intentional misuse.

Mevinphos appears to have a 'niche' in its directed use on several commodities in several regions of the U.S. However, research and adoption of alternative chemical and nonchemical tactics for insect management should continue to be integrated into the production and marketing of an economic, stable and safe food supply in a clean environment.

REFERENCES

- 19 Jan 1989. Air contaminants. Fed. Regist., 54(12, Bk. 2), 2332-983.
1986. Organophosphorus pesticides in sewage sludge 1985; organophosphorus pesticides in river and drinking water, an addition, 1985. Methods Exam. Waters Assoc. Mater., (Organo-Phosphorus Pestic. Sewage).
- 23 May 1984. Tolerances for pesticide chemicals in or on raw agricultural commodities; certain pesticide chemicals. Fed. Regist., 49(101), 21710-11.
- 4 Oct 1985. Pesticide chemicals category effluent limitations guidelines, pretreatment standards, and new source performance standards. Fed. Regist., 50(193), 40672-777.
- 5 Dec 1984. Tolerances and exemptions from tolerances for pesticide chemicals in or on raw agricultural commodities; certain pesticide chemicals. Fed. Regist., 49(235), 47493-4.
- Ackermann, Heinz. 1969. Thin-layer chromatographic-enzymic detection of organophosphorus insecticides. Thin-layer chromatographic behavior of some other insecticides. J. Chromatogr., 44(2), 414-18.
- Agnello, A., J. Kovach, J. Nyrop, and H. Reissig. 1988. Simplified Insect Management Program: A Guide for Apple Sampling Procedures in New York. Cornell Cooperative Extension. Cornell University. 20 pp.
- Agrawal, S. C.; Kushwaha, K. S. 1978. Efficacy of insecticidal control schedules against aphids and jassids on tomato. Indian J. Plant Prot., 6(2), 40-3.
- Ahmed, M.; Grant, W. F. 1972. Cytological effects of the pesticides Phosdrin and Bladex on Tradescantia and Vicia faba. Can. J. Genet. Cytol., 14(1), 157-65.
- Ajjan, Iskandar. 1974. Persistence and absorption of some organophosphorus insecticides when applied to alfalfa. Use Isot. Pestic. Pest Control, Proc. Symp., 243-52.
- Alabaster, J. S. 1969. Survival of fish in 164 herbicides, insecticides, fungicides, wetting agents, and miscellaneous substances. Int. Pest Contr., 11(2), 29-35.
- Alfalfa Seed Industry. 1989. Biological and economic assessment of ODM, trichlorfon, naled and mevinphos in alfalfa seed production. draft report. 22 p.
- Ambrus, Arpad; Hargitai, Eva; Karoly, Gabriella; Fulop, Andras; Lantos, Janos. 1981. General method for determination of pesticide residues in samples of plant origin, soil, and water. II. Thin layer

- chromatographic determination. J. Assoc. Off. Anal. Chem., 64(3), 743-8.
- Andersen, Karsten Stig. 1981. Pesticide residues in Danish food (1978-1979). Publ. - Statens Levnedsmiddelinst. (Den.), 54, 75 pp.
- Anderson, Arne. 1986. Monitoring and biased sampling of pesticide residues in fruits and vegetables. Methods and results, 1981-1984. Vaar Foeda, 38(Suppl. 1), 8-55.
- Andersson, Arne; Ohlin, Birgit. 1986. A capillary gas chromatographic multiresidue method for determination of pesticides in fruits and vegetables. Vaar Foeda, 38(Suppl. 2), 79-109.
- Anon. 1986. Mevinphos. Dangerous Prop. Ind. Mater. Rep., 6(1), 97-101.
- Anon. 1978. Pesticide mixture for use in agricultural and animal health outlets. Res. Discl., 176, 7-8.
- Anon. 20 Jun 1968. Food additives. 1-Methoxycarbonyl-1-propen-2-yl dimethylphosphate and its beta isomer. Fed. Regist., 33, 9071.
- Anon. 3 Jul 1979. Methyl 3-[(dimethoxyphosphinyl)oxy]butenoate, alpha and beta isomers; tolerances for residues. Fed. Regist., 44(129), 38845.
- Antal, A. 1975. Pharmacodynamic approach to the contradictions of acute experimental organophosphate intoxication. Environ. Qual. Saf., Suppl., 3(Pesticides), 615-17.
- Antoine, O.; Mees, G. 1971. Routine method for the determination of organophosphorus insecticides by thin-layer chromatography. J. Chromatogr., 58(2), 247-56.
- Antonin, P.; Fellay, D. 1979. Effect of volume and quantity of active material per hectare on the efficiency of phytosanitary treatment in arboriculture. Rev. Suisse Vitic., Arboric., Hortic., 11(3), 143-54.
- Arato Sugar, Eva; Szabari, Mrs. Gyorgy. 1970. Determination of organophosphorus insecticides in the air. Munkavedelem, 16(1-3), 50-4.
- Arretz V., Patricio; Lamborot C., Lilianne; Araya C., Jaime. 1976. Fruit plagues. I. Population studies and control of the aphid *Pentatrichopus* sp. Invest. Agric. (Santiago), 2(2), 67-73.
- Arretz V., Patricio; Lamborot C., Lilianne; Araya C., Jaime. 1976. Fruit plagues. II. Population study and control of *Tetranychus urticae* Koch. Invest. Agric. (Santiago), 2(2), 75-9.
- Atkins, E. L.; Kellum, D. 1986. Comparative morphogenic and toxicity studies on the effect of pesticides on honeybee brood. J. Apic. Res., 25(4), 242-55.

- Attia, F. I. 1977. Insecticide resistance in Plodia interpunctella (Hubner) (Lepidoptera: Pyralidae) in New South Wales, Australia. J. Aust. Entomol. Soc., 16, Pt. 2, 149-52.
- Attia, F. I. 1976. Insecticide resistance in Cadra cautella in New South Wales, Australia. J. Econ. Entomol., 69(6), 773-4.
- Attia, F. I. 1974. Laboratory evaluation of insecticides against Nysius vinitor and N. clevelandensis. J. Aust. Entomol. Soc., 13, Pt. 3, 161-4.
- Atwal, A. S.; Chaudhary, J. P.; Sohi, B. S. 1969. Chemical control of red vegetable mite, Tetranychus cucurbitae (Tetranychidae; acarina). J. Res., Punjab Agr. Univ., 6(1)(Suppl.), 214-19.
- Audemard, H.; Beauvais, Françoise; Descoins, C. 1977. Control of the codling moth (Laspeyresia pomonella L.) with a synthetic sex pheromone by the [male disruption] method: first trial in a commercial apple orchard. Rev. Zool. Agric. Pathol. Veg., 76(1), 15-24.
- Augustin, C.; Monteoliva, M. 1985. Contamination by pesticides in arable land in the province of Granada [Spain]. An. Edafol. Agrobiol., 44(11-12), 1785-94.
- Ault, James A.; Schofield, C. Michael; Johnson, Lyle D.; Waltz, R. H. 1979. Automated gel permeation chromatographic preparation of vegetables, fruits, and crops for organophosphate residue determination utilizing flame photometric detection. J. Agric. Food Chem., 27(4), 825-8.
- Aziz, Shawkyy A. 1973. Toxicity of certain insecticide standards against the southern armyworm. J. Econ. Entomol., 66(1), 68-70.
- Babad, Harry; Herbert, Washington. 1968. Nuclear magnetic resonance studies of phosphorus(V) pesticides. I. Chemical shifts of protons as a means of identification of pesticides. Anal. Chim. Acta, 41(2), 259-68.
- Bache, Carl A.; Lisk, Donald J. 1968. Versatility of OV-17 substrate for gas chromatography of pesticides. J. Ass. Offic. Anal. Chem., 51(6), 1270-1.
- Bailey, P. T.; O'Sullivan, D. F. 1978. Insecticides against larvae of the cacao webworm Pansepta teleturga Meyrick (Lepidoptera: Xyloryctidae). Papua New Guinea Agric. J., 29(1-4), 27-31.
- Baker, R. T. 1978. Insecticide resistance in the green peach-potato aphid Myzus persicae (Sulz.) (Homoptera: Aphididae). N. Z. J. Exp. Agric., 6(1), 77-82.
- Banares-Munoz, M. A.; Flores-Gonzalez, L. V.; Perez-Bernal, M. E.; Ruano-Casero, R. J.; Sanchez-Escribano, V. 1985. Adsorption of

organophosphorus pesticides on graphitic oxide. II. Phosdrin.
Agrochimica, 29(5-6), 435-44.

Baran, Milan; Pidany, Jan. 1973. Chemical protection against the cereal aphid (Sitobion avenae) on wheat. Pol'nohospodarstvo, 19(7), 533-9.

Barbulescu, A. 1975. Testing of various insecticides against the Cirphis unipuncta Haw. larvae at Rasht, Iran. An. Inst. Cercet. Cereale Plante Teh.-Fundulea, Acad. Stiinte Agric.

Barroga, S. F.; Morallo-Rejesus, B. 1981. Mechanism of joint action of insecticides on malathion-resistant diamond-back moth (Plutella xylostella). Philipp. Entomol., 5(1), 115-38.

Barroga, Salud F. 1969. Control of Lepidopterous pests attacking bush bean (Phaseolus vulgaris) blossoms and pods. Philipp. J. Plant Ind., 34(3-4), 159-62.

Barroga, Salud F. 1968. Biological notes and control of citrus leaf miner (Phyllocnistis citrella Stainton) affecting citrus seedlings. Philipp. J. Plant. Ind., 33(1-2), 17-36.

Barse, Joseph R., W. Ferguson, and R. Seem. 1988. Economic effects of banning soil fumigants. USDA, Economic Research Service, AER-602.

Becker, Guenther. 1979. Multimethod for simultaneous detection of 75 plant treatment preparations on plant material. Dtsch. Lebensm.-Rundsch., 75(5), 148-52.

Becker, Guenther. 1971. Simultaneous gas chromatographic determination of chlorinated hydrocarbons and phosphates in plant material. Deut. Lebensm.-Rundsch., 67(4), 125-6.

Beckman, Herman; Garber, Dennis. 1969. Pesticide residues. Recovery of 65 organophosphorus pesticides from Florisil with a new solvent-elution system. J. Ass. Offic. Anal. Chem., 52(2), 286-93.

Benedek, Pal; Komlodi, Jozsef; Herczeg, Gyula; Dely, Gyorgy. 1976. Preliminary experiments towards seed lucerne pest control harmless on bees. Toxicity of field-weathered insecticide residues to honeybees. Novenytermeles, 25(2), 129-40.

Benfatto, Domenico. 1973. Results of preliminary trials of chemical control against Archips rosanus in Sicily in 1973. Inform. Fitopatol., 23(12), 11-13.

Bennett, J. W.; Kofsky, Stanton; Bulbin, Alan; Dutton, Michael. 1985. Comparison of two defined media for inhibitor and incorporation studies of aflatoxin biosynthesis. Dev. Ind. Microbiol., 26, 479-86.

Berisford, Yvette C.; Ayres, John C. 1976. Effect of insecticides on growth and zearalenone (F-2) production by the fungus, Fusarium

graminearum. Environ. Entomol., 5(4), 644-8.

- Beyermann, K.; Eckrich, W. 1973. Gas chromatographic determination of insecticide residues in air samples. Fresenius' Z. Anal. Chem., 265(1), 4-7.
- Beynon, K. I.; Hutson, D. H.; Wright, A. N. 1973. Metabolism and degradation of vinyl phosphate insecticides. Residue Rev., 47, 55-142.
- Bezin, A. A. 1981. Agar diffusion method for the detection of organophosphorus pesticides in cadaveric material. Sud.-Med. Ekspert., 24(3), 38-40.
- Bhatia, S. K.; Pradhan, S. 1970. Resistance to insecticides in Tribolium castaneum. II. Cross-resistance characteristics of the p,p'-DDT-resistant strain. Indian J. Entomol., 32(Pt. 1), 32-8.
- Blanchard, R. A. and W. A. Douglas. 1953. The corn earworm as an enemy of field corn in the Eastern States. USDA Farmers Bull. 1651 (rev.): 15 pp.
- Blank, R. H.; Bell, D. S.; Cox, N. R. 1981. Screening seed protectants against black field cricket. Proc. N. Z. Weed Pest Control Conf., 34th, 156-60.
- Blank, R. H.; Osborne, G. O. 1979. Studies on cholinesterase of the mites Sancassania berlesei (Tyroglyphidae) and Tetranychus urticae (Tetranychidae): substrate-activity relationships and organophosphorus acaricide inhibition. N. Z. J. Agric. Res., 22(3), 491-6.
- Blank, R. H. 1979. Studies on the non-specific esterase and acetylcholinesterase isozymes electrophoretically separated from the mites Sancassania berlesei (Tyroglyphidae) and Tetranychus urticae (Tetranychidae). N. Z. J. Agric. Res., 22(3), 497-506.
- Blank, R. H.; Osborne, G. O. 1978. Comparison of the toxicity of organophosphorus acaricides to the mites Sancassania berlesei (Tyroglyphidae) and Tetranychus urticae (Tetranychidae). N. Z. J. Agric. Res., 21(3), 527-31.
- Bonnemaïson, L.; Hogrel, Mrs. R.; Augendre, M. 1972. Insecticide trials in the laboratory on caterpillars and eggs of Adoxophyes reticulana (Tortricidae) and Mamestra brassicae (Noctuidae). Phytat.-Phytopharm., 21(3), 171-81.
- Boocock, Geoff; Camilleri, Patrick. 1985. beta.-Cyclodextrin inclusion complex of mevinphos. J. Agric. Food Chem., 33(6), 1032-4.
- Boutibonnes, P.; Fischer, F. C. 1978. Effect of some compounds on antimicrobial properties of aflatoxin B1. Bull. Acad. Vet. Fr., 51(1),

- Boutibonnes, P. 1977. Effect of food additives and drugs on cytotoxic activity of aflatoxin B1 on bacterial cells. IRCS Med. Sci.: Libr. Compend., 5(12), 560.
- Bowman, B. T.; Sans, W. W. 1983. Further water solubility determinations of insecticidal compounds. J. Environ. Sci. Health, Part B, B18(2), 221-7.
- Bowman, Malcolm C.; Beroza, Morton. 1970. GLC [gas-liquid chromatographic] retention times of pesticides and metabolites containing phosphorus and sulfur on four thermally stable columns. J. Ass. Offic. Anal. Chem., 53(3), 499-508.
- Boyadzhiev, S.; Manolov, K. 1975. On the mechanism of some organophosphorus compounds influence on the functional state of the rat thyroid. Agressologie, 16(Spec. C), 67-72.
- Branca, Paolo; Quaglino, Paola. 1988. Pesticide residues in imported potatoes. Ind. Aliment. (Pinerolo, Italy), 27(260), 427-31.
- Bro-Rasmussen, Finn; Orboek, K.; Dahl, Mogens H. 1971. Disappearance of five pesticides after applications on lettuce from greenhouse and field experiments. Tidsskr. Planteavl, 75(1), 136-44.
- Brodesser, Josef; Schoeler, Heinz Friedrich. 1987. Determination of pesticides in water. Vom Wasser, 69, 61-71.
- Brodesser, Josef; Schoeler, Heinz Friedrich. 1987. An improved extraction method for the quantitative analysis of pesticides in water. Zentralbl. Bakteriol., Mikrobiol. Hyg., Ser. B, 185(1-2), 183-5.
- Brown, H. R.; Sharma, R. P. 1976. Synaptosomal adenosine triphosphatase (ATPase) inhibition by organophosphates. Experientia, 32(12), 1540-2.
- Brunelli, Agostino; Pancaldi, Davide; Svampa, Giorgio. 1975. Temperature effects on the in vivo and in vitro degradation of certain insecticides. Inf. Fitopatol., 25(3), 7-12.
- Buitendag, C. H.; Bronkhorst, G. J. 1980. Phosdrin 2 E.C. for bollworm and plant louse management - the biological control situation in citrus plants. Citrus Subtrop. Fruit J., 603, 16.
- Bunt, J. A. 1975. Effect and mode of action of some systemic nematicides. Meded. Landbouwhoges. Wageningen, 75-10, 127 pp.
- Burns, R. G. 1971. Loss of phosdrin and phorate insecticides from a range of soil types. Bull. Environ. Contam. Toxicol., 6(4), 316-21.
- Busch, Kenneth L.; Bursey, Maurice M.; Hass, J. Ronald; Sovocool, G. Wayne. 1978. Comparison of five ionization methods for producing mass

- spectra of typical organophosphorus pesticides. Appl. Spectrosc., 32(4), 388-99.
- Butani, Dharmo K.; Sahni, V. M. 1971. Toxicity of insecticides applied to cotton. Indian J. Agr. Sci., 41(1), 11-16.
- Butcher, M. R.; Penman, D. R. 1983. The effects of chemicals used on strawberry crops on introduced phytoseiid mites. Proc. N. Z. Weed Pest Control Conf., 36, 67-70.
- Calafiori, Maria H.; Kussama, Amelia R.; Machado de Franco, Ana C.; Bueno, Maria A. de Lourdes; Azzini, Vera da Conceicao; Pascoli, Marney; Angeli, Amria de Lourdes Testing of insecticides for the control of kale aphids [Brevicoryne brassicae]. 1969. Testing of insecticides for the control of kale aphids [Brevicoryne brassicae]. Solo, 61(1), 61-3.
- Calderon, Jose I.; Hare, Colin J. 1986. Control of diamondback moth in Southeast Asia by profenofos. Diamondback Moth Manage., Proc. Int. Workshop, 1st, Meeting Date 1985.
- Calora, Feliciano B.; Glass, Edward H.; Pescador, Manuel L.; Barroga, Salud F. 1968. Granular soil systemic insecticides (thimet and dithiolane E. I. 47,031) and phosdrin foliar spray against Plutella maculipennis Curtis and other insect pests of cabbage. Philipp. J. Plant. Ind., 33(1-2), 67-83.
- Calumpang, S. M. F.; Tejada, A. W.; Magallona, E. D. 1984. Photodecomposition on silica gel plates of nine selected organophosphorus insecticide formulations. Philipp. Entomol., 6(2), 191-204.
- Camoni, I.; Di Muccio, A.; Bellisai, M. S.; Bersacchi, M.; Citti, P. 1988. Pesticide residues in local foods (1983-1985). Rapp. ISTISAN, ISTISAN 88/17, 88 pp.
- Carere, A.; Ortali, V. A.; Cardamone, G.; Morpurgo, G. 1978. Mutagenicity of dichlorvos and other structurally related pesticides in Salmonella and streptomyces. Chem.-Biol. Interact., 22(2-3), 297-308.
- Carricaburu, Pierre; Lacroix, Roger; Lacroix, Josyane. 1981. Electroretinographic study of the white mouse intoxicated by organophosphorus: mevinphos and malathion. Toxicol. Eur. Res., 3(2), 87-9, 91.
- Carter, M. Kathleen; Maddux, Betty. 1974. Interaction of dichlorvos and anticholinesterases on the in vitro inhibition of human blood cholinesterases. Toxicol. Appl. Pharmacol., 27(2), 456-63.
- Cate, James R., Jr.; Bottrell, D. G. 1969. Insecticidal control of the greenbug on grain sorghum in the southern high plains of Texas, 1968. Tex. Agr. Exp. Sta., Progr. Rep., No. 2698, 10 pp.

- Cerna, V.; Karmazin, M. 1981. Pesticide residues on oranges (Citrus aurantium L. ssp. aurantium). I. Determination of phosphoorganic insecticide residues by gas chromatography. *Cesk. Farm.*, 30(7), 238-40.
- Chalfant, Richard B.; Todd, J. W.; Mullinix, Ben. 1979. Cabbage looper: ovicidal activity of pesticides in the laboratory. *J. Econ. Entomol.*, 72(1), 30-2.
- Chalfant, Richard B. 1976. Chemical control of insect pests of the southern pea in Georgia. *Ga., Agric. Exp. Stn., Res. Bull.*, 179, 31 pp.
- Chalfant, Richard B. 1973. Cabbage looper. Effect of temperature on toxicity of insecticides in the laboratory. *J. Econ. Entomol.*, 66(2), 339-41.
- Chalfant, Richard B. 1969. Toxicity of insecticides against two looper species, Pseudoplusia includens and Trichoplusia ni, in the laboratory. *J. Econ. Entomol.*, 62(6), 1343-4.
- Chan, Kim-Chi. 1978. Effect of some insecticides on the common Trichogramma (Trichogramma evanescens Westw.). *Gradinar. Lozar. Nauka*, 15(2), 73-9.
- Chang, Liang-Chuan. 1974. Toxicity of insecticides to parasite (Apanteles plutellae) of diamond-back moth. *Nung Yeh Yen Chiu*, 23(2), 143-8.
- Chang, Lian-Chuang. 1972. Vegetable insect control by thuricide and other insecticides. *T'ai-Wan Nung Yeh Chi K'an*, 8(4), 164-9.
- Chapman, R. Bruce; Penman, David R. 1980. The toxicity of mixtures of a pyrethroid with organophosphorus insecticides to Tetranychus urticae Koch. *Pestic. Sci.*, 11(6), 600-4.
- Chaudhry, M. Ismail; Shah, Bashir H. 1974. Preliminary studies on the biology and control of shisham bark borer, Agrilus dalbergiae Thery. *Pak. J. For.*, 24(4), 383-92.
- Chen, J. S.; Sun, C. N. 1986. Resistance of diamondback moth (Lepidoptera: Plutellidae) to a combination of fenvalerate and piperonyl butoxide. *J. Econ. Entomol.*, 79(1), 22-30.
- Chen, Jeng Shong; Lo, Kang Chen. 1989. Susceptibility of two bulb mites, Rhizoglyphus robini and R. setosus (Acarina: Acaridae), to some acaricides and insecticides. *Exp. Appl. Acarol.*, 6(1), 55-66.
- Chen, Zongmao. 1985. Comparative study on photodegradative rates of 35 pesticides. *Huanjing Kexue Xuebao*, 5(1), 70-84.
- Chen, Zongmao; Zabik, Matthew J.; Leavitt, Richard A. 1984. Comparative study of thin film photodegradative rates for 36 pesticides. *Ind.*

Eng. Chem. Prod. Res. Dev., 23(1), 5-11.

- Cheng, E. Y.; Lin, D. F.; Tsai, T. c.; Kao, C. H. 1988. Insecticide resistance study in Plutella xylostella L. IX. The selective metabolism of insecticides. Chung-hua Nung Yeh Yen Chiu, 37(3), 328-39.
- Cheng, Edward Y.; Kao, C. H.; Lin, D. F.; Tsai, T. C. 1986. Insecticide resistance study in Plutella xylostella (L.). VIII. The specificity of oxidative detoxication mechanism in larval stage. Chung-hua Nung Yeh Yen Chiu, 35(3), 375-86.
- Cheng, Edward Y.; Chou, Tou Mei; Kao, Ching Hua. 1985. Insecticide resistance study in Plutella xylostella (L.). VI. An experimental analysis of organophosphorus and synthetic pyrethroid resistances. Chung-hua Nung Yeh Yen Chiu, 34(1), 96-104.
- Cheng, Edward Y.; Chou, Tao Mei; Kao, Ching Hua. 1984. Insecticide resistance study in Plutella xylostella (L.). V. The induction, cross resistance and glutathione-S-transferase in relation to mevinphos-resistance. Chung-hua Nung Yeh Yen Chiu, 33(1), 73-80.
- Cheng, Edward Y.; Chou, Tao Mei; Kao, Ching Hua. 1983. Insecticide resistance study in Plutella xylostella (L.). IV. The activities of glutathione-S-transferase in the organophosphorus-resistant strains. Chung-hua Nung Yeh Yen Chiu, 32(4), 373-8.
- Chiriac, A.; Chiriac, Veronica; Mracec, M.; Simon, Z.; Vilceanu, R. 1976. Molecular orbital calculations and structure-anticholinesterasic activity correlations for enol-phosphates with pesticidal action. Univ. Timisoara, [Prepr.], Sect. Chim., (5), 11 pp.
- Chou, Tao Mei; Kao, Ching Hua; Cheng, Edward Y. 1984. The occurrence of insecticide resistance in three lepidopterous pests on vegetables. Chung-hua Nung Yeh Yen Chiu, 33(3), 331-6.
- Chou, Tao Mei; Cheng, Edward Y. 1983. Insecticide resistance study in Plutella xylostella (L.). III. The insecticide susceptibilities and resistance response of a native susceptible strain. Chung-hua Nung Yeh Yen Chiu, 32(2), 146-54.
- Chung, C. S. 1989. Non-toxic detergent for cleaning fruit and vegetables.
- Cleveland, J. P. and J. F. Treon. 1961. Response of experimental animals to Phosdrin insecticide in their daily diets. J. Agric. Food Chem. 9:484-88.
- Colburn, Richard; Asquith, Dean. 1973. Platynota idaeusalis. Toxicity of some insecticides to larvae. J. Econ. Entomol., 66(4), 991-2.
- Conner, A. J. 1983. The comparative toxicity of vineyard pesticides to wine yeasts. Am. J. Enol. Vitic., 34(4), 278-9.

- Constantinescu, Virginia. 1965. Biology and control of the cabbage aphid (Brevicoryne brassicae). An. Sect. Prot. Plant., Inst. Cent. Cercet. Agric., Bucharest, No. 3,.
- Cornette, Jimmy C.; Agerton, Bobby M.; Wolverton, Billy C. 1971. Detoxification of Phosdrin by Bacillus megaterium enzymes. U. S. Nat. Tech. Inform. Serv., AD Rep., No. 735273, 18 pp. Avail.
- Costa, C. L.; Costa, A. S.; Nagai, H. 1977. Insecticide application for the control of tomato spotted wilt virus. III. Effect of soil application of a systemic insecticide combined with foliage sprays. Fitopatol. Bras., 2(1), 23-34.
- Coutselinis, Antony; Dimopoulos, George. 1971. Organophosphorus pesticides in liver: their identification by gas-liquid chromatography (GLC). J. Forensic Med., 18(1), 11-14.
- Cowart, Rubel P.; Bonner, Frances L.; Epps, E. A., Jr. 1971. Rate of hydrolysis of seven organophosphate pesticides. Bull. Environ. Contam. Toxicol., 6(3), 231-4.
- Cremlyn, R. 1978. Pesticides, preparation and mode of action. John Wiley & Sons. New York. 240 p.
- Cunningham, I. C. 1971. Tobacco insecticide screening trials, 1961-1966. Queensl. J. Agr. Anim. Sci., 28(1), 13-18.
- Dabrowski, Zbigniew T. 1969. Laboratory studies on the toxicity of pesticides to Typhlodromus finlandicus and Phytoseiulus macropilis. Roczn. Nauk Roln., Ser. A, 95(3), 337-69.
- Daldrup, T.; Michalke, P.; Boehme, W. 1982. A screening test for pharmaceuticals, drugs and insecticides with reversed-phase liquid chromatography - retention data of 560 compounds. Chromatogr. Newsl., 10(1), 1-7.
- Daldrup, T.; Susanto, F.; Michalke, P. 1981. Combination of TLC, GLC (OV 1 and OV 17) and HPLC (RP 18) for a rapid detection of drugs, intoxicants and related compounds. Fresenius' Z. Anal. Chem., 308(5), 413-27.
- Damico, Joseph N. 1966. Mass spectra of some organophosphorus pesticide compounds. J. Assoc. Off. Anal. Chem., 49(5), 1027-45.
- Dang Quoc Quan, E. Fournier; Sonnier, M.; Lecorsier, A.; Bernadou, J. 1975. Reactivation by 1-hydroxyimino-4-methyl-2-pentanone of dog plasma cholinesterases inhibited by different organophosphates. Eur. J. Toxicol. Environ. Hyg., 8(4), 220-5.
- Danko, Jan. 1975. Effectiveness of several insecticides for aphids on tobacco. Agrochimica, 15(8), 241-3.

- Darakchieva, L. 1982. Resistance of the spider mite (Tetranychus urticae Koch.) to some acaricides in greenhouses. Gradinar. Lozar. Nauka, 19(7), 57-65.
- Davies, John E.; Enos, Henry F.; Barquet, Ana; Morgade, Carmen; Danauskas, Joseph X. 1979. Pesticide monitoring studies. The epidemiologic and toxicologic potential of urinary metabolites. Dev. Toxicol. Environ. Sci., Volume Date 1978, 4(Toxicol. Occup.
- De Batista, G. C.; Amaral, E.; Neto, A. Passarella. 1975. Toxicity of some insecticides and acaricides to hybrid workers of Apis mellifera ligustica L. and Apis mellifera adansonii L. (Hymenoptera, Apidae). An. Soc. Entomol. Bras., 4(1), 73-7.
- De Giovanni, Giovanni; Brunelli, Agostino. 1978. Control of aphids. Ital. Agric., 115(3), 105-10.
- DeBarr, Gary L.; Nord, John C. 1978. Contact toxicity of 34 insecticides to second-stage nymphs of Leptoglossus corculus (hemiptera:coreidae). Can. Entomol., 110(9), 901-6.
- Deang, Ricardo T.; Caburubias, Romeo B.; Quero, Elisea M. 1969. Insecticide test for the control of abaca corm weevil. Philipp. J. Plant Ind., 34(1-2), 79-87.
- Devillers, J.; Meunier, T.; Chambon, P. 1985. Usefulness of the dosage-effect-time relation in ecotoxicology for determination of different chemical classes of toxicants. Tech. Sci. Munic., (7-8), 329-34.
- Dick, G. L.; Heenan, M. P.; Love, J. L.; Udy, P. B.; Davidson, Flora. 1978. Survey of trace elements and pesticide residues in the New Zealand diet. 2. Organochlorine and organophosphorus pesticide residue content. N. Z. J. Sci., 21(1), 71-8.
- Dietzschold, Dirk; Weichbrodt, Karl Heinz; Weber, Frank. 1988. Two-wire arc spraying using flux-cored wires. Part 2. Schweissttechnik (Berlin), 38(12), 561-3.
- Dimova, P.; Boyadzhiev, S. 1982. Cytological and cytochemical changes in reactive connective tissue cells following exposure to Phosdrin. Folia Med. (Plovdiv), 24(4), 23-31.
- Dishovski, Kh. 1983. Studies of neuromuscular junctions under conditions of neuromuscular blockade. Acta Morphol. (Sofia), 4, 18-22.
- Dishovski, Kh.; Kavrakirova, S.; Popov, P. 1979. Effect of phosdrin on the cyclic adenosine 3':5'-monophosphate content in striated muscle. Dokl. Bolg. Akad. Nauk, 32(11), 1585-7.
- Dolan, John W.; Seiber, James N. 1977. Chlorine-selective detection for liquid chromatography with a Coulson electrolytic conductivity detector. Anal. Chem., 49(2), 326-31.

- Domeracki, Stefan; Studzinski, Andrzej; Jaruzelski, Marcin. 1976. Dynamics of the disappearance of pesticides in medicinal plants. Part V. Studies of organophosphorus insecticide residues in medicinal plants. *Herba Pol.*, 22(2), 117-31.
- Doull, J. 1976. The treatment of pesticide poisoning. pp 649-67 in C. F. Wilkinson (ed), *Insecticide Biochemistry and Physiology*. Plenum Press. New York. 768 p.
- Drake, John L.; Crowder, Larry A.; Ware, George W. 1971. Effects of x-irradiation on the toxicity of organophosphate insecticides to the house fly. *Pestic. Biochem. Physiol.*, 1(3-4), 373-80.
- Draughon, Frances A.; Ayres, John C. 1978. Effect of selected pesticides on growth and citrinin production by Penicillium citrinum. *J. Food Sci.*, 43(2), 576-8, 583.
- Dubois, Kenneth P. 1969. Combined effects of pesticides. *Can. Med. Ass. J.*, 100(4), 173-9.
- Dumitrescu, H.; Barduta, Zenovia; Dumitrescu, Domnica. 1976. Determination of organophosphorus insecticides by thin-layer chromatography and enzymic detection. *Rev. Ig., Bacteriol., Virusol., Parazitol., Epidemiol.*,.
- Duran, P.; Nicaud, J. M.; Mallevalle, J. 1984. Detection of organophosphorous pesticides with an immobilized cholinesterase electrode. *J. Anal. Toxicol.*, 8(3), 112-17.
- Dutton, M. F.; Anderson, M. S. 1980. Inhibition of aflatoxin biosynthesis by organophosphorus compounds. *J. Food Prot.*, 43(5), 381-4.
- Dyte, C. E.; Blackman, Dorothy G. 1972. Laboratory evaluation of organophosphorus insecticides against susceptible and malathion-resistant strains of Tribolium castaneum (Coleoptera, Tenebrionidae). *J. Stored Prod. Res.*, 8(2), 103-9.
- E.P.A. 1988. Mevinphos, Pesticide Fact Sheet No. 156. U. S. Environmental Protection Agency. Washington, DC.
- Ebing, W. 1985. Multimethod for the determination of pesticide residues in dead honeybees. I. Chlorine and phosphorus insecticides. *Fresenius' Z. Anal. Chem.*, 321(1), 45-8.
- Edland, Torgeir. 1976. Effectiveness of insecticides at different concentrations on aphids and natural enemies. *Forsk. Fors. Landbruket*, 27(6), 683-99.
- Eisler, Ronald. 1970. Latent effects of insecticide intoxication to marine mollusks. *Hydrobiologia*, 36(3-4), 345-52.

- Eisler, Ronald. 1970. Acute toxicities of organochlorine and organophosphorus insecticides to estuarine fishes. Tech. Pap. Bur. Sport Fish. Wildl. (US), No. 45-46, No. 46, 12 pp.
- Eisler, Ronald. 1970. Factors affecting pesticide-induced toxicity in an estuarine fish. Tech. Pap. Bur. Sport Fish. Wildl. (US), No. 45-46, No. 45, 20 pp.
- Engelhard, Arthur W. 1973. Chemical sprays. Concentration, drying rate, and temperature effects on chrysanthemum flower quality. Proc. Fla. State Hort. Soc., Volume Date 1972, 85, 370-4.
- Environmental Protection Agency. 1988. Pesticide Fact Sheet No. 156, U.S. EPA Office of Pesticide Programs, Washington, D.C.
- Ernst, G. F.; Schuring, F. 1970. Modified enzymic detection method for thin-layer chromatograms of pesticides. J. Chromatogr., 49(2), 325-8.
- Eto, M. 1974. Organophosphorus Pesticides: Organic and Biological Chemistry. CRC Press. Cleveland, OH. 387 p.
- Faust, Samuel D.; Gomaa, Hassan M. 1972. Chemical hydrolysis of some organic phosphorus and carbamate pesticides in aquatic environments. Environ. Lett., 3(3), 171-201.
- Fehringer, Norbert V.; Walters, Stephen M. 1984. Evaluation of capillary gas chromatography for pesticide and industrial chemical residue analysis. I. Comparison of retention ratios obtained on methyl silicone-coated capillary columns with published values for packed columns. J. Assoc. Off. Anal. Chem., 67(1), 91-5.
- Feng, Guolei; Shi, Junyue; Zhang, Guilin; Geng, Kunyuan. 1985. Synergism of SVI with insecticides and effects of their control on cotton aphids. Kunchong Zhishi, 22(4), 167-72.
- Feng, H. T. 1984. Joint action of insecticide-synergist mixtures on the diamondback moth. Chih Wu Pao Hu Hsueh Hui Hui K'an, 26(4), 401-12.
- Ferguson, Walter. 1989. Economic analysis simulator. Computer program obtained through personal communication.
- Fida, Peter. 1979. Results of pesticide residue studies on vegetable cultures under glass. Pflanzenarzt, 32(1), 9-11.
- Filip, I.; Alexandri, A. A. 1977. Studies on the biology, and chemical and biological control of the grape moth (*Lobesia botrana* Schiff.) in the Murfatlar vineyards. An. Inst. Cercet. Prot. Plant., Acad. Stiinte Agric. Silvice, 13,.
- Fjelddalen, Jac.; Renvall, Siv. 1974. Pesticide residues in field crops in the Nordic countries. Investigations on insecticides 1968-70. Acta Agr. Scand., 24(1), 17-32.

- Flaherty, D. L., M. A. Hoy, C. D. Lynn, and W. L. Peacock. 1981. Spider mites. in: Flaherty, D. L., F. L. Jensen, A. N. Kassimatis, H. Kido, and W. J. Moller eds. Grape Pest Management. Publication 4105. Cooperative Extension University of California Division of Agriculture and Natural Resources. 312 pp.
- Flockhart, Ian R.; Casida, John E. 1972. Relation of the acylation of membrane esterases and proteins to the teratogenic action of organophosphorus insecticides and eserine in developing hen eggs. *Biochem. Pharmacol.*, 21(19), 2591-603.
- Floru, Stefania; Goga, Elena; Tucaliuc, Mariana; Suta, Victoria; Isac, Mariana. 1977. Pesticide residues after the application of treatments in commercial orchards. *An. Inst. Cercet. Prot. Plant., Acad. Stiinte Agric. Silvice.*
- Fodor-Csorba, K.; Dutka, F. 1986. Selectivity and sensitivity of some thin-layer chromatographic detection systems. *J. Chromatogr.*, 365, 309-14.
- Foschi, S.; Cesari, A.; Ponti, I.; Bentivogli, P. G.; Bencivelli, A. 1970. Degradation and vertical movement of pesticides in the soil. *Notiz. Mal. Piante*, No. 82-83, 37-49.
- Fournier, E.; Sonnier, M.; Lecorsier, A. 1975. Distribution of organophosphate pesticides in two artificial plasmas, FSB and FSP. *Eur. J. Toxicol. Environ. Hyg.*, 8(4), 205-11.
- Frank, R.; Braun, H. E.; Holdrinet, M.; Stonefield, K. I.; Elliot, J. M.; Zilkey, B.; Vickery, L.; Cheng, H. H. 1977. Metal contents and insecticide residues in tobacco soils and cured tobacco leaves collected in southern Ontario. *Tob. Sci.*, 21, 74-80.
- Freed, V. H.; Schmedding, D.; Kohnert, R.; Haque, R. 1979. Physical chemical properties of several organophosphates: some implication in environmental and biological behavior. *Pestic. Biochem. Physiol.*, 10(2), 203-11.
- Fu, Lijie; Li, Li; Tang, Xiane; Gao, Xiaoqi; Wang, Shuqin; Qiao, Sibin. 1986. Toxicological studies on mevinphos. *Zhonghua Yufangyixue Zazhi*, 20(1), 12-14.
- Fukuto, T. R., E. O. Hornig, R. L. Metcalf, and M. Y. Winton. 1961. Configuration of the alpha- and beta-isomers of methyl 3-(dimethoxyphosphinyloxy)-crotonate (Phosdrin). *J. Org. Chem.* 26:4620-23.
- Furedi, Janos; Balint, Andor; Badawi, M. 1975. Study of the effects and after effects of pesticides on peas, by factorial analysis. *Agrartud. Kozl.*, 34(4), 719-37.

- Furedi, J.; Badawi, Mohamed E. 1974. Investigations into the physiological, cumulative and post-effect of pesticides in the seed production of peas. *Agrartud. Egy. Kozl.* 239-55.
- Furness, G. O. 1977. Chemical and integrated control of the long-tailed mealybug, Pseudococcus longispinus (Targioni-Tozzetti) (Hemiptera:Coccidae) in the riverland of South Australia. *Aust. J. Agric. Res.*, 28(2), 319-32.
- Fytizas-Danielidou, Raymondi. 1974. Effect of three solvents on the cutaneous toxicity of mevinphos on white rats. *Meded. Fac. Landbouwwet., Rijksuniv. Gent*, 39(2, Pt. 1), 857-63.
- Gabor, Miklos; Szalay, Laszlo; Szikszay, Margit. 1975. Effect of pesticides on the capillary resistance of the conjunctiva. Rabbit experiments. *Kiserl. Orvostud.*, 27(5), 496-9.
- Gabor, Miklos; Szikszay, Margit. 1975. Capillary toxicity of pesticides. *Kiserl. Orvostud.*, 27(5), 491-5.
- Gaj, Tibor; Karaman, Jozsef; Kiss, Laszlo. 1981. Effectiveness of pesticides used against apion in red clover seed production, nontoxic to bees. *Novenyvedelem (Budapest)*, 17(3), 131-2.
- Gargav, Vishnu P. 1967. Selective toxicity of some organophosphorus insecticides for grubs (Aphis gossypii and Alphis craccivora) and their entomophage (Coccinella septempunctata). *Tr. Tashkent. Sel'skokhoz. Inst.*, No. 21, 27-35.
- Garten, Charles T., Jr.; Trabalka, John R. 1983. Evaluation of models for predicting terrestrial food chain behavior of xenobiotics. *Environ. Sci. Technol.*, 17(10), 590-5.
- Genkov, D.; Zhelev, Zh.; Boyadzhiev, S. 1983. The effect of phosdrin on the acid resistance of erythrocytes in white rats. *Folia Med. (Plovdiv)*, 25(4), 54-7.
- Gera, Ramesh; Gupta, Dalip S. 1978. Potentiation of malathion by other insecticides against Musca domestica. *Pestic. Sci.*, 9(2), 151-4.
- Ghadiri, Modjtaba. 1967. Malformation and toxic effects of two organic phosphates and a carbamate on the chicken embryo and the adult chicken. 131 pp. Avail. 68-13,773.
- Ghyczy, Miklos; Wendel, Armin; Etschenberg, Eugen. 1984. Pesticides.
- Glofke, Edith. 1976. The results of bee poisoning with 22 insecticides. *Land- Forstwirtschaft. Forsch. Oesterr.*, 7, 125-9.
- Gomaa, El-Adarosy A. 1978. Effect of organophosphorus insecticides on dry beans. *Actes Congr. Int. Composes Phosphores*, 1st, Meeting Date 1977, 671-6.

- Goncalves, W.; De Batista, G. C.; Giacomelli, E. J. 1974. Susceptibility of *Parisoschoenus* species to insecticides and acaricides under laboratory conditions. *Cientifica*, 2(2), 258-62.
- Goodyer, G. J. 1978. Toxicity of insecticides to the green vegetable bug, *Nezara viridula* L. (Hemiptera: Pentatomidae). *J. Entomol. Soc. Aust.* (N.S.W.), 10, 50-2.
- Goodyer, G. J.; Brown, G. R. 1976. Laboratory screening of insecticides against larvae of the tobacco looper, *Chrysodeixis argentifera* (Guenee) (Lepidoptera: Noctuidae). *J. Entomol. Soc. Aust.* (N.S.W.), 9, 22-3.
- Goulart, B., G. Hamilton, J. Meade, S. Race, and J. Springer. 1988. New Jersey Commercial Strawberry Production Manual. Rutgers Cooperative Extension, Cook College, Rutgers, The State University of New Jersey. 27 pp.
- Goza, S. W. 1972. Infrared analysis of pesticide formulations. *J. Ass. Off. Anal. Chem.*, 55(5), 913-17.
- Gras, G.; Cisse, B. 1986. Toxicity of some organophosphate and carbamate compounds for "mange mil" (*Quelea quelea* L.). *J. Toxicol. Clin. Exp.*, 6(3), 175-87.
- Grigorov, St. P. 1969. Biology and control of aphids (Homoptera, Aphidodea) on berry crops. *Nauch. Tr., Vissh Selskostop. Inst.*, Sofia, Agron. Fak.
- Gruen, Gerhard; Sadek, Heinrich; Clausing, Peter. 1982. Evaluation of acute toxicity of plant protection agents in birds in relation to possible side effects in the field. *Nachrichtenbl. Pflanzenschutz DDR*, 36(6), 127-30.
- Gudehn, Ann; Kolmodin-Hedman, Birgitta. 1987. Sampling and determination of fenitrothion, dimethoate, mevinphos, linuron, metoxuron and trifluralin from air. *J. Chromatogr.*, 387, 420-7.
- Guignard, E.; Antonin, P.; Baillod, M. 1984. Efficacy of the secondary effects of insecticides used to control grape cluster worms. *Rev. Suisse Vitic., Arboric., Hortic.*, 16(6), 338-46.
- Gunderson, Ellen C.; Anderson, C. Clarine. 1980. Development and validation of methods for sampling and analysis of workplace toxic substances. DHHS (NIOSH) Publ. (U. S.), 80-133, 59 pp.
- Gunther, F. A.; Iwata, Y.; Papadopoulou, E.; Berck, B.; Smith, C. A. 1980. Rapid field method for estimating organophosphorus insecticide residues on crop foliage. *Bull. Environ. Contam. Toxicol.*, 24(6), 903-9.

- Gupta, S. P.; Sharma, S. S.; Singh, P. 1977. Electronic structure and anthelmintic activity of some organophosphorus anthelmintics. Indian J. Chem., Sect. B, 15(8), 731-3.
- Guthrie, Frank E. 1968. Nature and significance of pesticide residues on tobacco and in tobacco smoke. Beitr. Tabakforsch., 4(6), 229-46.
- Haider, Shamim; Upadhyaya, Netaji. 1986. Effect of commercial formulation of four organophosphorus insecticides on the LH-induced germinal vesicle breakdown in the oocytes of a freshwater teleost, Mystus vittatus (Bloch) - a preliminary in vitro study. Ecotoxicol. Environ. Saf., 12(2), 161-5.
- Haider, Shamim; Upadhyaya, Netaji. 1985. Effect of commercial formulation of four organophosphorus insecticides on the ovaries of a freshwater teleost, Mystus vittatus (Bloch) - a histological and histochemical study. J. Environ. Sci. Health, Part B, B20(3), 321-40.
- Hamilton, J. T.; Attia, F. I. 1976. The susceptibility of the parasite Apanteles glomeratus (L.) (Hym.: Braconidae) to insecticides. J. Entomol. Soc. Aust. (N.S.W.), 9, 24-5.
- Hammock, Bruce D.; Sparks, Thomas C.; Mumby, Susanne M. 1977. Selective inhibition of JH esterases from cockroach hemolymph. Pestic. Biochem. Physiol., 7(6), 517-30.
- Hansen, Torkild; Noehr Rasmussen, A.; Schadegg, E. 1972. Fungicides and insecticides in fruit and garden crops, 1971. Tidsskr. Planteavl, 76(5), 682-706.
- Hansen, Torkil; Noehr Rasmussen, A.; Schadegg, E. 1972. Experiments with fungicides and insecticides in fruit and gardening crops. Tidsskr. Planteavl, 76(1), 77-104.
- Hara, Arnold H.; Kaya, Harry K. 1983. Development of the entomogenous nematode, Neoaplectana carpocapsae (Rhabditida: Steinernematidae), in insecticide-killed beet armyworm (Lepidoptera: Noctuidae). J. Econ. Entomol., 76(3), 423-6.
- Hara, Arnold H.; Kaya, Harry K. 1982. Effects of selected insecticides and nematocides on the in vitro development of the entomogenous nematode Neoaplectana carpocapsae. J. Nematol., 14(4), 486-91.
- Harris, C. R.; Turnbull, S. A.; McLeod, D. G. R. 1985. Contact toxicity of twenty-one insecticides to adults of the carrot rust fly (Diptera: Psilidae). Can. Entomol., 117(8), 1025-7.
- Harris, C. R.; Svec, H. J.; Tolman, J. H.; Chapman, R. A. 1983. Laboratory and field studies on the effectiveness and persistence of some insecticides used to control the tomato hornworm. Tob. Int., 185(15), 178-82.

- Harris, C. R.; Turnbull, S. A.; Whistlecraft, J. W.; Surgeoner, G. A. 1982. Multiple resistance shown by field strains of house fly, Musca domestica (Diptera: Muscidae), to organochlorine, organophosphorus, carbamate, and pyrethroid insecticides. *Can. Entomol.*, 114(5), 447-54.
- Harris, Charles R.; Svec, Harry J. 1970. Laboratory studies on the contact toxicity of some insecticides to honeybees. *Proc. Entomol. Soc. Ont.*, Volume Date 1969, 100, 165-7.
- Harris, Charles R.; Gore, F. 1970. Laboratory tests on the contact toxicity of some insecticides to adults of the cabbage flea beetle, Phyllotreta cruciferae. *J. Econ. Entomol.*, 63(3), 1025-7.
- Harris, Charles R.; Svec, H. J. 1970. Laboratory studies of the toxicity of technical formulations of some insecticides to cabbage loopers. *J. Econ. Entomol.*, 63(2), 666-7.
- Harris, Charles R.; Svec, H. J. 1970. Laboratory tests on the toxicity of some insecticides to the cabbage aphid. *J. Econ. Entomol.*, 63(2), 665-6.
- Harvan, Donald J.; Hass, J. Ronald; Busch, Kenneth L.; Bursey, Maurice M.; Ramirez, Fausto; Meyerson, Seymour. 1979. Direct observation of the monomeric metaphosphate anion. *J. Am. Chem. Soc.*, 101(24), 7409-10.
- Hassan, Sherif A. 1983. Results of laboratory testing of a series of pesticides on egg parasites of the genus Trichogramma (Hymenoptera, Trichogrammatidae). *Nachrichtenbl. Dtsch. Pflanzenschutzdienstes* (Braunschweig), 35(2),.
- Hauri, P.; Freuler, J.; Bertuchoz, P.; Hugi, H. 1973. Preliminary control of lettuce root aphid, Pemphigus bursarius in cultivated composites. *Rev. Suisse Viticult., Arboricult. Hort.*, 5(6), 179-82.
- Hayes, W. L. Jr. 1982. *Pesticides studied in man*. Williams & Wilkins. Baltimore, MD. 672 p.
- Hebling-Beraldo, Maria Jose A.; Vicelli-Zanao, Regina C. 1983. Effects of organophosphorus and carbamate insecticides on the oxygen consumption of Atta laevigata (F. Smith, 1858) and Atta sexdens rubropilosa, Forel, 1908 (Hymenoptera-Formicidae). *Cienc. Cult. (Sao Paulo)*, 35(2), 211-14.
- Henriet, J.; Martijn, A. 1986. Design and conduct of collaborative studies: CIPAC collaborative experience in pesticide formulations. *J. Assoc. Off. Anal. Chem.*, 69(3), 408-9.
- Hill, Elwood F.; Heath, Robert G.; Spann, James W.; Williams, Joseph D. 1975. Lethal dietary toxicities of environmental pollutants to birds. *U. S., Fish Wildl. Serv., Spec. Sci. Rep.: Wildl.*, 191, 61 pp.

- Hirka, Gabor. 1980. Biochemical changes parallel to toxic functional modifications in heart tissue cultures. *Nehezvegyip. Kut. Intez. Kozl.*, 10, 223-9.
- Ho, H. S.; Liu, T. S. 1969. Comparison of insecticides against the diamond back moth. *T'ai-Wan Nung Yeh Chi K'an*, 5(2), 90-3.
- Holmes, Joseph H.; Starr, Herbert G., Jr.; Hanisch, Robert C.; Von Kaulla, Kurt N. 1974. Short-term toxicity of mevinphos in man. *Arch. Environ. Health*, 29(2), 84-9.
- Holmstead, Roy L.; Casida, John E. 1974. Chemical ionization mass spectrometry of organophosphorus insecticides. *J. Assoc. Off. Anal. Chem.*, 57(5), 1050-5.
- Hoogcarspel, A. P.; Jobsen, J. A. 1984. Laboratory method for testing side-effects of pesticides on Encarsia formosa Gahan (Hym., Aphelinidae). Results with pesticides used in glasshouse tomato crops in the Netherlands. *Z. Angew. Entomol.*, 97(3), 268-78.
- Hopper, Marvin L. 1982. Automated gel permeation system for rapid separation of industrial chemicals and organophosphate and chlorinated pesticides from fats. *J. Agric. Food Chem.*, 30(6), 1038-41.
- Horng, Wei-Huai. 1979. Safety regulations in use of pesticides in Taiwan. *FFTC Book Ser.*, 14(Sensible Use Pestic.), 189-97.
- Hrdy, Ivan; Kremheller, Hedwig Th.; Kuldova, Jelena; Luders, Wolfgang; Sula, Jan. 1986. Resistance to insecticides of the hop aphid, Phorodon humuli, in Bohemian Bavarian and Baden-Wurtembergian hop growing areas. *Acta Entomol. Bohemoslov.*, 83(1), 1-9.
- Hrdy, Ivan; Skrobal, Dimitrij. 1976. Effect of two juvenoids on honey bee colonies (Apis mellifera L.). *Apidologie*, 7(1), 39-47.
- Hrdy, Ivan; Zeleny, Jiri. 1973. Effects of juvenoids on the population density of Phorodon humuli in a cage experiment (Homoptera, Aphididae). *Acta Entomol. Bohemoslov.*, 70(6), 386-9.
- Hrlec, G. 1984. Pesticide mobility in fine calcareous loam. *Zemljiste Biljka*, 33(2), 189-93.
- Hsieh, Dennis P. H. 1973. Inhibition of aflatoxin biosynthesis of dichlorvos. *J. Agr. Food Chem.*, 21(3), 468-70.
- Hu, Chia-Chien; Wang, Lian-Chuan. 1969. *Thosea sinensis* Walker. *T'ai-Wan Nung Yeh Chi K'an*, 5(4), 107-15.
- Hudson, Rick H.; Haegeler, M. A.; Tucker, Richard K. 1979. Acute oral and percutaneous toxicity of pesticides to mallards: correlations with mammalian toxicity data. *Toxicol. Appl. Pharmacol.*, 47(3), 451-60.

- Hummel, Susan V.; Yost, Richard A. 1986. Tandem mass spectrometry of organophosphate and carbamate pesticides. *Org. Mass Spectrom.*, 21(12), 785-91.
- Hurkova, Johana; Weyda, Frantisek. 1982. Response to pesticides of diapausing female Tetranychus urticae (Acarina, Tetranychidae). *Vestn. Cesk. Spol. Zool.*, 46(2), 92-9.
- Hurkova, Johana; Horak, Antonin. 1979. Toxicity of the organophosphate and carbamate aphicides to the pea aphid, Acyrtosiphon pisum (Homoptera, Aphididae). *Acta Entomol. Bohemoslov.*, 76(2), 137-9.
- Hurkova, Johana. 1978. Toxicity of 30 pesticides for a susceptible strain of the two-spotted spider mite, Tetranychus urticae. *Acta Entomol. Bohemoslov.*, 75(2), 109-18.
- Hurkova, Johana. 1973. Multiple resistance to insecticides in greenhouse samples of Myzus persicae (Homoptera, Aphidoidea). *Acta Entomol. Bohemoslov.*, 70(1), 13-19.
- Hyde, K. M.; Crandall, J. C.; Kortman, K. E.; McCoy, W. K. 1978. EEG, ECG, and respiratory response to acute insecticide exposure. *Bull. Environ. Contam. Toxicol.*, 19(1), 47-55.
- Isenhour, David J. 1985. Efficacy of insecticides against Spissistilus festinus (Say), Empoasca fabae (Harris), and Lygus lineolaris (Palisot de Beauvois) in alfalfa in Georgia. *J. Entomol. Sci.*, 20(1), 121-8.
- Ishikura, Shunji; Onodera, Sukeo; Sumiyashiki, Shunji; Kasahara, Teruhisa; Nakayama, Motoko; Watanabe, Shozo. 1984. Evaporation and thermal decomposition of organophosphorus pesticides during cooking of rice. *Shokuhin Eiseigaku Zasshi*, 25(2), 203-8.
- Ito, Kan. 1977. Water quality standards. 66. Nippon Suisan Shigen Hogo Kyokai Geppo, 151, 16-19.
- Janda, Jiri. 1975. Toxicity of the insecticide Phosdrin 24 EC for small game. *Sci. Agric. Bohemoslov.*, 7(2), 127-33.
- Jendzelovsky, Jozef. 1975. Biological effectiveness of aphicides under field conditions. *Agrochemia*, 15(7), 207-11.
- Jensen, F. L., and D. L. Flaherty. 1981a. Grape leafhopper. in: Flaherty, D. L., F. L. Jensen, A. N. Kassimatis, H. Kido, and W. J. Moller eds. *Grape Pest Management*. Publication 4105. Cooperative Extension University of California Division of Agriculture and Natural Resources. 312 pp.
- Jensen, F. L., and D. L. Flaherty. 1981b. Grape leafroller. in: Flaherty, D. L., F. L. Jensen, A. N. Kassimatis, H. Kido, and W. J. Moller eds. *Grape Pest Management*. Publication 4105. Cooperative Extension University of California Division of Agriculture and Natural

Resources. 312 pp.

- Jensen, F. L., and H. Kido. 1981. Aphids. in: Flaherty, D. L., F. L. Jensen, A. N. Kassimatis, H. Kido, and W. J. Moller eds. Grape Pest Management. Publication 4105. Cooperative Extension University of California Division of Agriculture and Natural Resources. 312 pp.
- Johansen, Carl A.; Mayer, Daniel F.; Eves, Jack D.; Kious, Christopher W. 1983. Pesticides and bees. Environ. Entomol., 12(5), 1513-18.
- Juarez, Americo Osvaldo; Guzman, Jorge A. 1988. Acute toxicity of mevinphos, parathion and phosphamidon to toad Bufo arenarum (H). Comun. Biol., 6(3), 259-63.
- Kaempe, Bent. 1974. GLC [gas-liquid chromatographic] system for the identification of an unknown drug in forensic chemistry. Arch. Pharm. Chemi, Sci. Ed., 2(5), 145-52.
- Kahlon, Prem S.; Banerjee, Mukul R. 1979. Evaluation of bravo, phosdrin and telvar as possible environmental mutagens. Bull. Environ. Contam. Toxicol., 22(3), 365-70.
- Kalinova, G.; Izmirova, N.; Kaloyanova, F. 1987. Minimum period (after plant spraying) for safe work with Phosdrin and Vydate in greenhouses. Khig. Zdraveopaz., 30(5), 39-43.
- Kanda, Mizuho; Takahama, Keiichi; Kiyofuzi, Takashi; Watanabe, Chiyuki; Miyazaki, Yoshimitsu; Masuda, Tetsuro; Noguchi, Kenichi; Miki, Motoi Analysis of chlorinated and organic phosphorus pesticides by gas chromatography. 1969. Analysis of chlorinated and organic phosphorus pesticides by gas chromatography. Nippon Hoigaku Zasshi, 23(6), 477-86.
- Kanda, Mizuho; Takahama, Keiichi; Tsubaki, Masaki; Yoshizaki, Hisayuki; Watanabe, Chiyuki; Kiyofuzi, Takashi. 1968. Qualitative analyses of synthetic organic pesticides by spot test. Nippon Hoigaku Zasshi, 22(3), 229-51.
- Kao, Ching Hua; Cheng, Edward Y.; Lin, Doun Fang. 1986. Insecticide resistance study in Plutella xylostella (L.) VII. Action patterns of insecticides and the resistance impacts. Chung-hua Nung Yeh Yen Chiu, 35(1), 107-17.
- Kaplan, J.; Lenormand, C.; Comba, D. 1972. Protection of date bunches against bird attacks. Fruits, 27(6), 439-44.
- Kaspar, Petr; Leblova, Sylva. 1982. Contribution to the understanding of the effect of some herbicides on maize (Zea mays L.). Biologia (Bratislava), 37(12), 1169-73.
- Keith, Lawrence H.; Alford, Ann L. 1969. Supplementary interpretations of the N.M.R. spectra of phosphorus pesticides. Anal. Chim. Acta, 44(2),

- Khalil, S. K.; Shah, M. A.; Naeem, M. 1985. Laboratory studies on the compatibility of the entomopathogenic fungus Verticillium lecanii with certain pesticides. *Agric., Ecosyst. Environ.*, 13(3-4), 329-34.
- Khan, D. U.; Ahmed, K. A. 1967. Field trials on the comparative effectiveness of some insecticides for the control of mustard aphid (Aphis brassicae). *Agr. Pak.*, 18(3), 299-303.
- Khattak, Sana Ullah Khan. 1984. Laboratory evaluation of some insecticides for the control of painted bug, Bagrada hilaris (Burm.) (Heteroptera: Pentatomidae). *Pak. J. Sci. Ind. Res.*, 27(4), 240-2.
- Khera, K. S.; Lyon, D. A. 1968. Chick and duck embryos in the evaluation of pesticide toxicity. *Toxicol. Appl. Pharmacol.*, 13(1), 1-15.
- Khristova, E. 1976. Resistance of the peach aphid (Myzodes persicae Sulz.) to Wofatox and phosdrin. *Gradinar. Lozar. Nauka*, 13(6), 55-9.
- Kido, H.; Stafford, E. M.; McCalley, N. F. 1971. Orange tortrix on grapes in Salinas Valley. *Calif. Agr.*, 25(7), 10-11.
- Kim, Yong Hwa; Woodrow, James E.; Seiber, James N. 1984. Evaluation of a gas chromatographic method for calculating vapor pressures with organophosphorus pesticides. *J. Chromatogr.*, 314, 37-53.
- Kinoshita, G. B.; Harris, C. R.; Svec, H. J.; McEwen, F. L. 1978. Laboratory and field studies on the chemical control of the crucifer flea beetle, Phyllotreta cruciferae (Coleoptera: Chrysomelidae), on cruciferous crops in Ontario. *Can. Entomol.*, 110(8), 795-803.
- Klunker, Roland. 1974. Problems of resistance in mites, with particular reference to resistance to dimethoate. *Tagungsber., Akad. Landwirtschaftswiss. D. D. R.*, 126, 133-50.
- Knaak, J. B.; Jackson, T.; Fredrickson, A. S.; Rivera, L.; Maddy, K. T.; Akesson, N. B. 1980. Safety effectiveness of closed-transfer, mixing-loading, and application equipment in preventing exposure to pesticides. *Arch. Environ. Contam. Toxicol.*, 9(2), 231-45.
- Knaak, J. B.; Jackson, T.; Fredrickson, A. S.; Maddy, K. T.; Akesson, N. B. 1980. Safety effectiveness of pesticide mixing-loading and application equipment used in California in 1976. *Arch. Environ. Contam. Toxicol.*, 9(2), 217-29.
- Kodama, J. K., M. S. Horse, H. H. Anderson, M. K. Dunlap, and C. H. Hine. 1954. Comparative toxicity of two vinyl-substituted phosphates. *Arch. Ind. Hyg. Occup. Med.* 9:45-61.
- Koellner, Volkard. 1973. Resistance to organophosphorus compounds of spider mites from Chrysanthemum cultures. *Nachrichtenbl. Deut.*

Pflanzenschutzdienstes (Brunswick), 25(1), 7-11.

Koenig, K.; Hassan, S. A. 1986. Resistance and cross-resistance of the predatory mite Phytoseiulus persimilis (Athias-Henriot) to insecticidal phosphate esters. J. Appl. Entomol., 101(3), 206-15.

Korsos, Istvan, Mrs.; Lantos, Janos. 1984. Thin-layer chromatographic identification of pesticides. Novenyvedelem (Budapest), 20(1), 30-4.

Kotev, G.; Krustanov, L. 1978. Changes in electroencephalographic and electromyographic activity after acute phosdrin intoxication and prophylaxis with PA-2 (ulkarol). Acta Med. Bulg., 5(1), 84-7.

Kotev, G.; Atsev, E.; Kolchev, Kh.; Mashkova, D. 1976. Some neurophysiological mechanisms in poisonings with organophosphorus pesticides. Nevrol., Psikhiatr. Nevrokhir., 15(1), 35-42.

Kotev, G.; Kilimov, N.; Machkova, D.; Filipova, D. 1975. Electromyographic and kymographic studies of phosdrin poisoning and its treatment with the preparation AV-72. Dokl. Bolg. Akad. Nauk, 28(9), 1277-80.

Koula, Vaclay; Oliberius, Jiri. 1982. Study of the effect of low-volume applications on the biological activity of insecticides. Rostl. Vyroba, 28(7), 689-94.

Kowalska, Tatiana; Szczepanska, Krystyna; Pruszyński, Stefan; Bartkowiak, Antonina. 1980. Toxicity of pesticides to parasitic and predatory insects. Entomol. Intensyf. Roln., [Mater. Ses. Nauk.], Meeting Date 1977,.

Kowalska, Tatiana; Pruszyński, Stefan. 1969. Toxicity of some insecticides to the common lace-wing fly, Chrysopa carnea (Neuroptera, Chrysopidae). Biul. Inst. Ochr. Rosl., No. 45, 99-107.

Krause, Richard T.; August, E. Michael. 1983. Applicability of a carbamate insecticide multiresidue method for determining additional types of pesticides in fruits and vegetables. J. Assoc. Off. Anal. Chem., 66(2), 234-40.

Krijgsman, Willem; Van de Kamp, Cornelis G. 1976. Analysis of organophosphorus pesticides by capillary gas chromatography with flame photometric detection. J. Chromatogr., 117(1), 201-5.

Krizi, Josef. 1975. Insecticide resistance of the hop aphid. Soripar, 22(4), 147-9.

Larson, L. L., E. E. Kenaga and R. W. Morgan. 1985. Commercial and experimental organic insecticides. Entomol. Soc. Amer. Misc. Pub. 105 p.

Laska, Pavel. 1981. Low pirimicarb-susceptibility of buckthorn-potato

aphid, Aphis nasturtii (Homoptera, Aphididae). Acta Entomol. Bohemoslov., 78(1), 61-2.

Laska, Pavel. 1980. Residues of Phosdrin 24 EC in vegetables and fruits. Agrochemia (Bratislava), 20(11), 326-8.

Leary, John S., Jr. 1974. Controlling mycotoxin production using certain dialkyl enol phosphates.

Leblova, Sylva; Cerovska, Noemi; Cerna, Vera. 1981. The effect of herbicides on the activity of pea alcohol dehydrogenase. Biochem. Physiol. Pflanz., 176(3), 283-5.

Lee, Sung-Lien; Lee, Wen-Tai. 1979. Studies on the resistance of diamondback moth, Plutella xylostella to commonly used insecticides. Chung-hua Nung Yeh Yen Chiu, 28(4), 225-36.

Lehotzky, Kornelia; Magda, Katalin. 1977. Effect of spironolactone and atropine on the experimental poisoning by organophosphates. Munkavedelem, 23(1-3, Suppl.), 17-20.

Leoni, V.; Puccetti, G. 1971. Thin-layer chromatographic analysis of organophosphate pesticides in the study of environmental contamination (Italian surface waters). Farmaco, Ed. Prat., 26(7), 383-93.

Leoni, Valerio; Puccetti, Giovanna. 1969. Gas-liquid chromatography of pesticides on OV-17 stationary phase. J. Chromatogr., 43(3), 388-91.

Lewin, J. F.; Love, J. L. 1974. Death caused by the ingestion of mevinphos. Forensic Sci., 4(3), 253-5.

Lewis, Mark F.; Mertens, Henry W.; Steen, Jo Ann. 1973. Behavioral changes from chronic exposure to pesticides used in aerial application. Effects of Phosdrin on the performance of monkeys and pigeons on variable interval reinforcement schedules. Aerosp. Med., 44(3), 290-3.

Lewis, Mark F.; Mertens, Henry W.; Steen, Jo Ann. 1972. Behavioral changes from chronic exposure to pesticides used in aerial applications. Effects of phosdrin on the performance of monkeys and pigeons on variable interval reinforcement schedules. U. S. Nat. Tech. Inform. Serv., AD Rep., No. 749893, 6 pp. Avail.

Li, Gwo Chen; Wong, Sue Sun; Jiang, Chi Ling. 1981. Establishment of maximum pesticide tolerances in food crops. K'o Hsueh Nung Yeh (Taipei), 29(7-8), 211-21.

Li, Gwo-Chen; Wong, Sue-Sun; Tang, Ching-Chyuan. 1979. Residue analysis of several pesticides in asparagus. Chih Wu Pao Hu Hsueh Hui Hui K'an, 21(3), 277-84.

- Li, Hsi-Shan; Wu, Kuo-Chia; Liu, Tu-Hsin; Wen, Hung-Tse. 1979. Evaluation of the insecticides for control of the diamond back moth and imported cabbageworm. T'ai-wan Nung Yeh Shuang Yueh K'an, 15(6), 28-36.
- Lienk, S. E. 1980. European Red Mite. Tree Fruit IPM, Insect Identification Sheet No. 10. New York State Agricultural Experiment Station, Geneva, NY.
- Lin, Tsung. 1986. Studies on control of insect pests of stored grain with insecticide smoke's. Chung-hua Nung Yeh Yen Chiu, 35(3), 360-74.
- Lo, Kang-Chen; Chao, Rou-Su. 1976. Preliminary studies on the resistance of three spider mites to chemicals in Taiwan. Chung Hua Nung Yeh Yen Chiu, 25(1), 23-36.
- Lo, Paul Kang-Chen; Chao, Susan Rou-Su. 1975. Preliminary study on the toxicity of common pesticides to natural enemies of red spider mites. Chung-Hua Nung Hsueh Hui Pao, 92, 81-6.
- Lopez-Lopez, M. C.; Hermoso, R.; Monteoliva, M. 1982. Cholinesterase isoenzymes and their inhibition by organophosphorus pesticides in Moniezia expansa, Fasciola hepatica, and Dicrocoelium dendriticum. Rev. Iber. Parasitol., Extraordinario, 291-300.
- Love, J. L.; McGrath, H. J. W.; Donnelly, R. G. C. 1975. Mevinphos residues on strawberries. N. Z. J. Exp. Agric., 3(2), 187-9.
- Luke, Milton A.; Masumoto, Herbert T.; Cairns, Thomas; Hundley, Harvey K. 1988. Levels and incidences of pesticide residues in various foods and animal feeds analyzed by the Luke multiresidue methodology for fiscal years 1982-1986. J. - Assoc. Off. Anal. Chem., 71(2), 415-33.
- Luke, Milton A.; Froberg, Jerry E.; Masumoto, Herbert T. 1975. Extraction and cleanup of organochlorine, organophosphate, organonitrogen, and hydrocarbon pesticides in produce for determination by gas-liquid chromatography. J. Assoc. Off. Anal. Chem., 58(5), 1020-6.
- Magallona, E. D.; Callejas, E. E. 1979. Degradation of triazophos and mevinphos in cabbage and soil. Philipp. Agric., 62(4), 276-84.
- Magallona, E. D.; Tejada, A. W.; Varca, L. M.; Mallari, V. C., Jr. 1977. Insecticide residues in vegetables. I. Methyl parathion, mevinphos and triazophos residues in cabbage. Philipp. Entomol., 3(5-6), 261-71.
- Magallona, E. D.; Callejas, E. E. 1977. Loss of triazophos and mevinphos in soil. Philipp. Entomol., 3(5-6), 339-52.
- Maier, J. 1988. Hop quality and plant protection. Monogr. - Eur. Brew. Conv., 13, 56-62.

- Malhotra, C. P.; Katiyar, R. N. 1977. Control of *Eublemma amabilis* Moore - a serious predator of the lac insect, *Kerria lacca* (Kerr). I. Screening of insecticides for their safety to lac insect. Indian J. Entomol., Volume Date 1975, 37(4), 385-96.
- Manem, J.; Mallevialle, J.; Durant, P.; Chabert, E. 1983. [Environmental] determination of organophosphorus pesticides and carbamates by a butyrylcholinesterase-containing enzymic electrode. Eau, Ind., Nuisances, 74, 31-4.
- Mansingh, Ajai; Rhodes, Llewellyn F. 1985. Residual toxicity of various insecticidal formulations to the coffee berry borer, *Hypothenemus hampei* Ferrari (Scolytidae: Coleoptera). Insect Sci. Its Appl., 6(2), 209-12.
- Mansingh, Ajai; Rhodes, Llewellyn F. 1983. Bioassay of various formulations of insecticides on the egg and larval stages of the coffee berry borer *Hypothenemus hampei* Ferrari (Scolytidae: Coleoptera). Insect Sci. Its Appl., 4(3), 223-6.
- Martel, Pierre; Boivin, Guy; Belcourt, Jean. 1986. Efficacy and persistence of different insecticides against the tarnished plant bug, *Lygus lineolaris* (Heteroptera: Miridae), on a season-long host plant, *Coronilla varia*. J. Econ. Entomol., 79(3), 721-5.
- Martin, A. D.; Norman, G.; Stanley, P. I.; Westlake, G. E. 1981. Use of reactivation techniques for the differential diagnosis of organophosphorus and carbamate pesticide poisoning in birds. Bull. Environ. Contam. Toxicol., 26(6), 775-80.
- Martin, Henry; Drabek, Jozef. 1970. Pesticidal O,O-dialkyl phosphates.
- Martincova, J.; Citkova, M. 1969. Metathion and Phosdrin in plant material. Souhrn Ref. Diskusnich Prispevku Prednesenych Ved. Semin., "Prakt."
- Mashkova, D. 1979. Antidote for treating the acute intoxication of animals and poultry by organophosphorus-containing pesticides.
- Mashkova, D.; Kotev, G.; Kolchev, Kh. 1978. Changes in electromyographic potentials in cats after poisoning with superlethal doses of Phosdrin and treatment with a preparation containing dolin. Eksp. Med. Morfol., 17(1), 32-6.
- Mashkova, D.; Kolchev, Kh.; Kotev, G. 1977. Electrophysiological changes in the functional antagonism between organophosphorus pesticides and their antidotes. Eksp. Med. Morfol., 16(4), 216-22.
- Mashkova, D. 1976. Effect of arpenal on EMG-potentials of the anterior tibial muscle of cats after acute poisoning with Phosdrin. Eksp. Med. Morfol., 15(4), 237-41.

- Mass, J. L., ed. 1984. Compendium of Strawberry diseases. The American Phytopathological Society. 138 pp.
- Matsuani, Shigenobu. 1969. Toxicity of pesticides to eggs and adults of Tetranychus telarius. Noyaku Kensasho Hokoku, No. 9, 14-18.
- Matsumura, F. 1985. Toxicology of Insecticides. 2d ed. Plenum Press. New York. 598 p.
- McCann, J. A.; Teeters, W.; Urban, D. J.; Cook, N. 1981. A short-term dietary toxicity test on small mammals. ASTM Spec. Tech. Publ., 757(Avian Mamm. Wildl. Toxicol.), 132-42.
- McClanahan, R. J.; Founk, J. 1983. Toxicity of insecticides to the green peach aphid (Homoptera: Aphididae) in laboratory and field tests, 1971-1982. J. Econ. Entomol., 76(4), 899-905.
- McClanahan, Robert J. 1976. Tests of insecticides for asparagus beetle control and residues on the crop. Proc. Entomol. Soc. Ont., Volume Date 1975, 106, 52-5.
- McClanahan, Robert J. 1975. Insecticides for control of the Colorado potato beetle (Coleoptera, Chrysomelidae). Can. Entomol., 107(6), 561-5.
- McEwen, Freeman L.; Ritcey, Gwen; Braun, Heinz; Frank, Richard; Ripley, Brian D. 1980. Foliar pesticide residues in relation to worker re-entry. Pestic. Sci., 11(6), 643-50.
- McLeod, Paul. 1987. Influence of temperature on contact and volatile toxicities of aphicides against the green peach aphid, Myzus persicae (Sulzer) (Homoptera: Aphididae). J. Entomol. Sci., 22(4), 362-6.
- Meier, Robert W. 1978. A field portable mass spectrometer for monitoring organic vapors. J., Am. Ind. Hyg. Assoc., 39(3), 233-9.
- Meisch, M. V.; Randolph, N. M.; Teetes, George L. 1970. Phytotoxicity and effectiveness of insecticides applied to corn. Tex. Agr. Exp. Sta., Progr. Rep., PR-2708, 9 pp.
- Meister, C. W.; Meister, J. S.; Bardalaye, P. C. 1984. Labeling insecticides for use in fields of Chinese cabbage and endive. Proc. Fla. State Hortic. Soc., Volume Date 1983, 96, 113-14.
- Melkebeke, G.; Verstraeten, R.; Steurbaut, W.; Dejonckheere, W. 1984. Uptake and phytotoxicity of some insecticides-acaricides in hydroponic culture. Meded. Fac. Landbouwwet., Rijksuniv. Gent, 49(3b), 1203-9.
- Mertens, Henry W.; Steen, Jo Ann; Lewis, Mark F. 1976. Some behavioral effects of pesticides: the interaction of mevinphos and atropine in pigeons. Aviat., Space Environ. Med., 47(2), 137-41.

- Mertens, Henry W.; Steen, Jo Ann; Lewis, Mark F. 1975. Effects of mevinphos on appetitive operant behavior in the gerbil. *Psychopharmacologia*, 41(1), 47-52.
- Mertens, Henry W.; Lewis, Mark F.; Steen, Jo A. 1974. Behavioral effects of pesticides. Phosdrin and free operant escape-avoidance behavior in gerbils. *Aerosp. Med.*, 45(10), 1171-6.
- Mestres, R.; Chevallier, C.; Espinoza, C.; Cornet, R. 1977. Application of coupled gas chromatography-mass spectrometry in methods for the study and determination of pesticide residues and organic micropollutants in environmental and food materials. *Ann. Falsif. Expert. Chim.*, 70(751), 177-88.
- Metcalf, C. L., W. P. Flint, and R. L. Metcalf. 1962. Destructive and Useful Insects. Their Habits and Control. 4th ed. McGraw-Hill Book Company. 1087 pp.
- Meyerson, Seymour; Harvan, Donald J.; Hass, J. Ronald; Ramirez, Fausto; Marecek, James F. 1984. The monomeric metaphosphate anion in negative-ion chemical-ionization mass spectra of phosphotriesters. *J. Am. Chem. Soc.*, 106(23), 6877-83.
- Michalek, H.; Nemesio, R.; Meneguz, A.; Bisso, G. M. 1983. Interactions between anticholinesterase agents and neuroleptics in terms of cholinesterase inhibition in brain and other tissues of rats. *Arch. Toxicol., Suppl.*, 6, 386-90.
- Mierzwinski, Adam; Witkiewicz, Zygfryd. 1987. Piezoelectric detectors coated with liquid crystal materials. *Talanta*, 34(10), 865-71.
- Miles, James W.; Dale, William E. 1978. Gas chromatographic determination of organophosphorus pesticides by in-block methylation. *J. Agric. Food Chem.*, 26(2), 480-2.
- Miller, Hope; Cramer, Paul; Drinkwine, Arbor; Shan, Alice; Trischan, Glenn; Going, John. 1981. Development of methods for pesticides in wastewater: applicability of a general approach. *Adv. Identif. Anal. Org. Pollut. Water*, Volume 1, 115-38.
- Minarik, Erich; Ragala, Peter. 1980. Effect of insecticides and acaricides on the yeast flora of grapes and musts. *Kvasny Prum.*, 26(4), 89-94.
- Molnar, Jenő; Szeberenyi, Judit; Varga Aniko, P.; Nagy, Margit. 1975. Pupil dilation activity of locally applied choline esterase inhibitors in mice. *Kiserl. Orvostud.*, 27(1), 78-82.
- Molnar, Annamaria M.; Tuske, Marton; Molnar, Jenő; Szeberenyi, Judit M.; Paksy, Katalin A. 1974. Aerosol-inhalation of pesticides in rats. Acute inhalation toxicity of mevinphos. *Egeszsegtudomány*, 18(3), 254-7.

- Monteoliv A, M.; Augustin, C. 1985. Contamination by pesticides in surface waters in the province of Granada [Spain]. *An. Edafol. Agrobiol.*, 44(11-12), 1767-83.
- Morallo-Rejesus, B.; Baldos, E. P.; Tejada, A. M. 1981. Evaluation of insecticides against coffee berry borer and its residues in processed coffee. *Philipp. Entomol.*, 4(5), 415-33.
- Morallo-Rejesus B.; Eroles, D. 1976. Potentiation of paired insecticide combinations on Plutella xylostella (Linn.) and other cabbage insects. *Philipp. Entomol.*, 3(3-4), 225-40.
- Morallo-Rejesus, B.; Chow, Hue Tze; Sanchez, F. F.; Cadapan, E. P. 1972. Malathion and phosdrin residues in petsai. *Philipp. Entomol.*, 2(4), 291-5.
- Morello, A., A. Vardanis and E. Y. Spencer. 1968. Mechanism of detoxication of some organophosphorus compounds: the role of glutathione-dependent demethylation. *Can. J. Biochem.* 46:885.
- Morpurgo, Giorgio; Aulicino, Francesca; Bignami, Margherita; Conti, Luigi; Velcich, Anna. 1977. Relationship between structure and mutagenicity of dichlorvos and other pesticides. *Atti Accad. Naz. Lincei, Cl. Sci. Fis., Mat. Nat., Rend.*, 62(5),.
- Moscioni, A. David; Engel, Judith L.; Casida, John E. 1977. Kynurenine formamidase inhibition as a possible mechanism for certain teratogenic effects of organophosphorus and methylcarbamate insecticides in chicken embryos. *Biochem. Pharmacol.*, 26(23), 2251-8.
- Mueller, B. 1973. Determination of pesticide residues in bee honey. I. Semiquantitative thin-layer chromatographic determination of insecticide residues in bee honey. *Nahrung*, 17(3), 381-6.
- Myhr, Brian C. 1973. Screen for pesticide toxicity to protein and RNA synthesis in HeLa cells. *J. Agr. Food Chem.*, 21(3), 362-7.
- Nagano, Masayoshi; Sakai, Michihiko. 1974. Ovicidal and larvicidal activities of cartap (Padan) against the diamondback moth, Plutella xylostella. *Takeda Kenkyusho Ho*, 33(4), 252-9.
- Nagy, Ferenc; Szalay-Marzso, Laszlo. 1969. Control of Leptinotarsa decemlineata in *Solanum laciniatum* crops. *Herba Hung.*, 8(3), 93-9.
- Naneva, E.; Georgiev, B. 1976. Study on the extent of the biologic effect of organophosphorus anticholinesterase agents and some possibilities of their quantitative biological analysis. *Vet.-Med. Nauki*, 13(8), 36-42.
- Naqvi, S. M. Z. 1973. Toxicity of twenty-three insecticides to a tubificid worm Branchiura sowerbyi from the Mississippi delta. *J. Econ. Entomol.*, 66(1), 70-4.

- Naqvi, Syed M.; Ferguson, Denzel E. 1969. Pesticide tolerances of selected freshwater invertebrates. J. Miss. Acad. Sci., Volume Date 1968, 14, 121-7.
- Natoff, I. L. 1970. Influence of the route of exposure on the acute toxicity of cholinesterase inhibitors. Arh. Hig. Rada Toksikol., 21(4), 347-52.
- Natoff, Ian L. 1970. Influence of the route of exposure on the acute toxicity of cholinesterase inhibitors. J. Eur. Toxicol., 3(6), 363-7.
- Natoff, I. L. 1967. Influence of the route of administration on the toxicity of some cholinesterase inhibitors. J. Pharm. Pharmacol., 19(9), 612-16.
- Natskova, V. 1982. Control of hothouse whitefly on flower plants. Gradinarstvo, 63(8), 42-4.
- Nehez, M.; Boros, P.; Ferke, A.; Mohos, J.; Palotas, M.; Vetro, G.; Zimanyi, M.; Desi, I. 1988. Cytogenetic examination of people working with agrochemicals in the southern region of Hungary. Regul. Toxicol. Pharmacol., 8(1), 37-44.
- Nicholas, Mark L.; Powell, David L.; Williams, Theodore R.; Huff, Sarah R. 1976. Reference Raman spectra of ten phosphorus-containing pesticides. J. Assoc. Off. Anal. Chem., 59(5), 1071-80.
- Noeddegaard, E.; Hansen, Torkil; Noehr Rasmussen, A. 1970. Testing of fungicides and insecticides in 1969. Tidsskr. Planteavl, 74(5), 618-61.
- Nottoli, Raffaello; Taponeco, Giuseppe. 1975. Consumption of pesticides in the province of Pisa, and the use of infrared spectroscopy for their characterization. Boll. Chim. Unione Ital. Lab. Prov., 1(3), 138-49.
- Obersteiner, E. J.; Sharma, R. P. 1978. Evaluation of cytotoxic responses caused by selected organophosphorus esters in chick sympathetic ganglia cultures. Can. J. Comp. Med., 42(1), 80-8.
- Oehmichen, M.; Pedal, I.; Besserer, K.; Gencic, M. 1984. Inhibition of non-specific leukocyte esterase activity. Absence of monocyte esterase activity due to phosphoric and thiophosphoric acid ester intoxication. Forensic Sci. Int., 25(3), 181-9.
- Oomen, P. A. 1986. A sequential scheme for evaluating the hazard of pesticides to bees, Apis mellifera. Meded. Fac. Landbouwwet., Rijksuniv. Gent, 51(3B), 1205-13.
- Orbaek, Kirsten. 1987. Pesticide residues in Danish food 1984 to 1985. Publ. - Levnedsmiddelstyr. (Den.), 147, 135 pp.

- Orbaek, Kirsten. 1985. Pesticide residues in Danish food from 1982 to 1983. Publ. - Statens Levnedsmiddelinst. (Den.), 107, 140 pp.
- Ordog, Vince. 1979. The effect of organophosphate insecticides on green algae. Hidrol. Kozl., 59(1), 36-42.
- Osselton, Michael David; Snelling, Robert Duncan. 1986. Chromatographic identification of pesticides. J. Chromatogr., 368(2), 265-71.
- Pablo, F. E. 1979. Pesticide residues in leafy vegetables. Nat. Appl. Sci. Bull., Volume Date 1977, 29(2), 51-68.
- Pallade, Sulamith; Gabrielescu, Eliana; London, Micaela; Simionovici, Rodica. 1970. Toxicity of some organophosphoric insecticides. Igiena, 19(1), 43-52.
- Pancaldi, Davide; Vismara, Clelia; Lugaresi, Claudio. 1975. Safety times of some pesticides used to protect strawberries. Inf. Fitopatol., 25(7), 15-20.
- Pandey, U. K.; Pandey, Alok; Lekha, Chandra. 1982. Insecticidal control of garden pest-earwig. Z. Angew. Zool., 69(2), 133-7.
- Pavkov, Sava; Sovljanski, Radmila; Soldatovic, Danilo; Stefanovic, Olga. 1981. Isolation of some organophosphorus insecticides from water by adsorption. II. Adsorption of some organophosphorus insecticides from water by talc and activated carbon. Arh. Farm., 31(5-6), 297-303.
- Pavkov, Sava; Sovljanski, Radmila; Soldatovic, Danilo; Stefanovic, Olga. 1981. Isolation of some organophosphorus insecticides from water by adsorption. I. Adsorption of some organophosphorus insecticides from water by Florisil and kieselguhr. Arh. Farm., 31(5-6), 289-95.
- Peairs, F. B. 1986. Alfalfa webworm control Johnstown, Co, 1985. Insect. and Acar. Tests 11: 216.
- Perng, F. S.; Yao, M. C.; Hung, C. F.; Sun, C. N. 1988. Teflubenzuron resistance in diamondback moth (Lepidoptera: Plutellidae). J. Econ. Entomol., 81(5), 1277-82.
- Pflugmacher, J.; Ebing, W. 1974. Purification of phosphoric acid insecticide residues in vegetable extracts by gel chromatography on Sephadex LH-20. J. Chromatogr., 93(2), 457-63.
- Pickering, R. G.; Pickering, C. E. 1974. Methods for the estimation of acetylcholinesterase activity in the erythrocytes of laboratory animals given carbamates or organophosphorus compounds. Arch. Toxicol., 31(3), 197-216.
- Pickering, C. E.; Pickering, R. G. 1971. Estimation of acetylcholinesterase activity in the plasma and brain of laboratory animals given carbamates or organophosphorus compounds. Arch.

Toxikol., 27(3), 292-310.

- Popov, Veselina. 1968. Chemical control experiments against the cotton bollworm (Chloridea obsoleta) in tobacco. Bulg. Tyutyun, 13(4), 24-5.
- Pradhan, S.; Jotwani, M. G.; Sarup, Prakash. 1968. Bioassay of different insecticides on the important insect pests and predators of agricultural importance. Pestic., Symp., Meeting Date 1964, 92-103. Edited by: Majumder,.
- Pressley, Thomas A.; Longbottom, James E. 1982. The determination of organophosphorus pesticides in industrial and municipal wastewater: method 622. Report, EPA-600/4-82-008; Order No. PB82-156027, 25 pp. Avail. NTIS.
- Prinsloo, Susan M.; De Beer, Pieter R. 1987. Gas chromatographic relative retention data for pesticides on nine packed columns: II. Organophosphorus and organochlorine pesticides, using electron-capture detection. J. Assoc. Off. Anal. Chem., 70(5), 878-88.
- Prinsloo, Susan M.; De Beer, Pieter R. 1985. Gas chromatographic relative retention data for pesticides on nine packed columns: I. Organophosphorus pesticides, using flame photometric detection. J. Assoc. Off. Anal. Chem., 68(6), 1100-8.
- Proctor, Nick H.; Moscioni, A. David; Casida, John E. 1976. Chicken embryo NAD levels lowered by teratogenic organophosphorus and methylcarbamate insecticides. Biochem. Pharmacol., 25(7), 757-62.
- Pumala, Kanjana; Viriyanondha, Somchit. 1987. Improper labeled concentration of pesticide formulation, a cause of an environmental health problem. J. Natl. Res. Council. Thailand, 19(1), 45-59.
- Qian, ChuanFan; Sanders, Paul F.; Seiber, James N. 1985. Accelerated degradation of organophosphorus pesticides with sodium perborate. Bull. Environ. Contam. Toxicol., 35(5), 682-8.
- Quinot, E.; Moncelon, B.; Millard, M. 1978. Toxic substances in the air. Determination method for calculation of exposure limit value. Cah. Notes Doc., 93, 547-65.
- Rabloczky, G.; Mader, Maria R. 1977. Effects on blood pressure of the anticholinesterases physostigmine and mevinphos in course of development in chickens. Acta Physiol. Acad. Sci. Hung., 50(4), 437-46.
- Rabloczky, Gyorgy; Magda, Katalin. 1977. Occurrence and drug effects on lung edema in acute experimental intoxication by cholinesterase inhibitors. Kiserl. Orvostud., 29(5), 530-8.
- Rabloczky, G.; Jancso, T.; Mader, Maria R. 1976. Alterations of the function of the sympathetic nervous system after anticholinesterase

- treatment. Congr. Hung. Pharmacol. Soc., [Proc.], Volume Date 1974, 2(6, Symp.)
- Rabloczky, Gyorgy; Jancso, Tamas. 1973. Analysis of the pressor activity of the cholinesterase inhibitor Phosdrin, in the lesser circulation. Kiserl. Orvostud., 25(2), 161-9.
- Reagan, Ronald. 1989. Economic report of the President. U.S. Government Printing Office, Washington, D.C.,.
- Reiff, Bernard. 1976. Reactivation of phosphorylated brain cholinesterase in vivo. Med. Prot. Chem.-Warf. Agents, [Pap. Symp.], Meeting Date 1974, 82-7.
- Reiff, B.; Lambert, S. M.; Natoff, I. L. 1971. Inhibition of brain cholinesterase by organophosphorus compounds in rats. Arch. Int. Pharmacodyn. Ther., 192(1), 48-60.
- Reissig, W. H. 1980. Obliquebanded leafroller. Tree Fruit IPM, Insect Identification Sheet No. 5. New York State Agricultural Experiment Station, Geneva, NY.
- Reith, Jan F.; Schrijver, C. A.; Drost, R. H. 1971. Demonstration of toxic substances with Drosophila melanogaster. Pharm. Weekbl., 106(10), 77-90.
- Renvall, Siv; Clementz, Eva; Ekstrom, Georg. 1976. Residues of organophosphorus pesticides in fruits, berries, vegetables, juices, syrups, wines and rice. Var Foeda, 28, Suppl. 2, 58-84.
- Renvall, Siv. 1975. Examination of phosphorus pesticides in fruits and vegetables. Var Foeda, 27(8-9), 310-12.
- Revzin, Alvin M. 1980. Effects of organophosphate pesticides and alcohol on visual mechanisms. Neurotoxic. Visual Syst., [Rochester Int. Conf. Environ. Toxic.],.
- Revzin, Alvin M. 1976. Effects of mevinphos (Phosdrin) on unit discharge patterns in avian hippocampus. Aviat., Space Environ. Med., 47(6), 608-11.
- Revzin, Alvin M. 1976. Effects of organophosphate pesticides and other drugs on subcortical mechanisms of visual integration. Aviat., Space Environ. Med., 47(6), 627-9.
- Revzin, Alvin M. 1973. Transient blindness due to the combined effects of mevinphos and atrophine. U. S. Nat. Tech. Inform. Serv., AD Rep., No. 763555, 5 pp. Avail.
- Revzin, Alvin M. 1973. Subtle changes in brain functions produced by single doses of mevinphos (Phosdrin). U. S. Nat. Tech. Inform. Serv., AD Rep., No. 763509, 9 pp. Avail.

- Rhodes, Llewellyn F.; Mansingh, Ajai. 1985. Bioefficacy of various insecticidal formulations on the coffee berry borer Hypothenemus hampei ferrari in dry coffee berries. J. Coffee Res., 15(3-4), 82-9.
- Rhodes, Llewellyn F.; Mansingh, Ajai. 1981. Susceptibility of the coffee berry borer Hypothenemus hampei Ferrari to various insecticidal formulations. Insect Sci. Its Appl., 2(4), 227-31.
- Rider, D. A.; Berger, R. S. 1985. Toxicity and electrophysiological and behavioral effects of insecticide-pheromone-related chemicals on cabbage looper moths. Environ. Entomol., 14(4), 427-32.
- Rider, J. Alfred; Puletti, Ernesto J.; Swader, Joyce I. 1975. Minimal oral toxicity level for mevinphos in man. Toxicol. Appl. Pharmacol., 32(1), 97-100.
- Rider, J. A., J. I. Swader and E. J. Puletti. 1972. Anticholinesterase toxicity studies with Guthion, Phosdrin, Di-Syston, and Trithion in human subjects. Fed. Proc. 31:520.
- Rider, J. A., E. J. Puletti, and J. I. Swader. 1975. The minimal oral toxicity level for mevinphos in man. Toxicol. Appl. Pharmacol. 32:97-100.
- Ripa S., Renato. 1981. Advances in the control of the tomato fruit moth Scrobipalpula absoluta (Meyr.). II. Chemical control trials. Agric. Tec. (Santiago), 41(3), 113-19.
- Ripley, Brian D.; Braun, Heinz E. 1983. Retention time data for organochlorine, organophosphorus, and organonitrogen pesticides on SE-30 capillary column and application of capillary gas chromatography to pesticide residue analysis. J. Assoc. Off. Anal. Chem., 66(5), 1084-95.
- Roach, John A. G.; Andrzejewski, Denis. 1987. Analysis for pesticide residues by collision-induced fragmentation. Chem. Anal. (N. Y.), 91(Appl. New Mass Spectrom. Tech. Pestic.
- Roberts, D. V.; Wilson, Andrew. 1973. Monitoring biological effects of anticholinesterase pesticides. Pestic. Environ.: Continuing Controversy, Pap. Inter-Am. Conf.
- Roche, L., E. LeJeune and R. J. M. Madonna. 1961. Four poisonings with Phosdrin. Arch. Mal. Prof. Med. Trav. Secur. Soc. 22:52-55.
- Roger, Jean C.; Upshall, David G.; Casida, John E. 1969. Structure-activity and metabolism studies on organophosphate teratogens and their alleviating agents in developing hen eggs with special emphasis on Bidrin. Biochem. Pharmacol., 18(2), 373-92.
- Roos, A. H.; Van Munsteren, A. J.; Nab, F. M.; Tuinstra, L. G. M. Th. 1987. Universal extraction/cleanup procedure for screening of

- pesticides by extraction with ethyl acetate and size exclusion chromatography. *Anal. Chim. Acta*, 196, 95-102.
- Roseboom, H.; Groenemeijer, G. S. 1981. Residues of organophosphorous pesticides as determined by capillary gas chromatography. *Meded. Fac. Landbouwwet., Rijksuniv. Gent*, 46(1), 325-30.
- Ross, Ralph T.; Biros, Francis J. 1970. Correlations between phosphorus-31 NMR chemical shifts and structures of some organophosphorus pesticides. *Anal. Chim. Acta*, 52(1), 139-41.
- Sagredos, A. N.; Eckert, W. R. 1976. Methods for the determination of phytopharmaceuticals in tobacco and tobacco products. Part I. Simultaneous determination of water-soluble organophosphorus pesticides. *Beitr. Tabakforsch.*, 8(7), 438-46.
- Saini, M. L.; Sharma, P. D. 1970. Relative efficacy of insecticides against lemon butterfly, Papilio demoleus (Papilionidae; Lepidoptera). *J. Res., Punjab Agr. Univ.*, 7(2), 203-7.
- Salame, M. 1968. Use of gas chromatography with electron capture in the analysis of organophosphate pesticides. *Ann. Biol. Clin. (Paris)*, 26(7-9), 1011-21.
- Salvisberg, W.; Becker, G.; Voss, G. 1980. Insecticides and electrophysiological symptoms: a comparative study on a tarsal motor nerve of Locusta migratoria. *Pestic. Biochem. Physiol.*, 13(1), 63-70.
- Sanchez Camazano, M.; Sanchez Martin, M. J. 1983. Factors influencing interactions of organophosphorus pesticides with montmorillonite. *Geoderma*, 29(2), 107-18.
- Sanchez Martin, M. J.; Sanchez Camazano, M. 1980. Interaction of phosdrin with montmorillonite. *Clay Miner.*, 15(1), 15-23.
- Sanchez, F. F.; Esguerra, N. M.; Calora, F. B.; Barroga, S. F. 1969. Field evaluation of the granular soil systemic insecticide Cyolane and foliar sprays of Ambithion, Accothion and Phosdrin against *Plutella xylostella* and other insect pests of cabbage. *Philipp. J. Plant Ind.*, 34(3-4), 131-41.
- Sanchez-Camazano, M.; Sanchez-Martin, M. J. 1987. Interaction of some organophosphorus pesticides with vermiculite. *Appl. Clay Sci.*, 2(2), 155-65.
- Sanchez-Camazano, M.; Sanchez Martin, M. J. 1977. Interaction of organophosphorus pesticides with montmorillonite. III. Sumithion and phosdrin. *Proc. - Eur. Clay Conf.*, 3rd, 165-6. Edited by: Rosenqvist, I. Th.
- Sanders, Paul F.; Seiber, James N. 1984. Organophosphorus pesticide volatilization. Model soil pits and evaporation ponds. *ACS Symp.*

Ser., 259(Treat. Disposal Pestic. Wastes), 279-95.

Sanders, Paul F.; Seiber, James N. 1983. A chamber for measuring volatilization of pesticides from model soil and water disposal systems. *Chemosphere*, 12(7-8), 999-1012.

Sarup, Prakash; Singh, D. S.; Sircar, Prasad; Amarpuri, Swaran; Lal, Rattan; Saxena, V. S.; Srivastava, V. S. 1972. Relative toxicity of some important pesticides to the adults of Bagrada cruciferarum (Pentatomidae, Hemiptera). *Indian J. Entomol.*, Volume Date 1971, 33, Pt. 4, 452-6.

Satpathy, J. M.; Padhi, G. K.; Dutta, D. N. 1968. Toxicity of eight insecticides to the coccinellid predator Chilomenes sexmaculata. *Indian J. Entomol.*, 30(Pt. 2), 130-2.

Saxton, Wilbur L. 1987. Emergence temperature indexes and relative retention times of pesticides and industrial chemicals determined by linear programmed temperature gas chromatography. *J. Chromatogr.*, 393(2), 175-94.

Schaeffers, Falk Immo. 1982. Preparation of capillary columns. Application examples with self-prepared capillary columns. *Lebensmittelchem. Gerichtl. Chem.*, 36(4), 83-7.

Schafer, E. W., Jr.; Bowles, W. A., Jr.; Hurlbut, J. 1983. The acute oral toxicity, repellency, and hazard potential of 998 chemicals to one or more species of wild and domestic birds. *Arch. Environ. Contam. Toxicol.*, 12(3), 355-82.

Schafer, Edward W. 1972. Acute oral toxicity of 369 pesticidal, pharmaceutical, and other chemicals to wild birds. *Toxicol. Appl. Pharmacol.*, 21(3), 315-30.

Schoeler, H. F.; Brodesser, J. 1988. An improved extraction method for the quantitative analysis of pesticides in water. *Comm. Eur. Communities*, [Rep.] EUR, EUR 11350, Org. Micropollut.

Schulten, H. R.; Sun, Sien. 1982. Field desorption mass spectrometry and identification of standard organophosphorus pesticides in wastewater. *Huanjing Huaxue*, 1(2), 123-32.

Schulten, G. G. M.; Van de Klashorst, G.; Russell, V. M. 1976. Resistance of Phytoseiulus persimilis A. H. (Acari: Phytoseiidae) to some insecticides. *Z. Angew. Entomol.*, 80(4), 337-41.

Schutz, Nandor; Katona, Antal; Mezo, Gabor; Petter, Laszlo. 1988. Experiments for controlling aphids and Noctuids in green pea stands. *Novenyvedelem (Budapest)*, 24(11), 507-11.

Seiber, J. N.; Markle, J. C. 1972. Hydrolysis and solvent partition of phosdrin and related organophosphates. *Bull. Environ. Contam.*

Toxicol., 7(2/3), 72-4.

- Seifert, Josef; Casida, John E. 1978. Relation of yolk sac membrane kynurenine formamidase inhibition to certain teratogenic effects of organophosphorus insecticides and of carbaryl and eserine in chicken embryos. *Biochem. Pharmacol.*, 27(22), 2611-15.
- Sengalevich, Georgi. 1967. Cossus cossus and agents for combatting it. *Gorskostop. Nauka*, 4(3), 43-57.
- Sergeeva, Dafina. 1971. Thin layer chromatographic demonstration of organophosphorus pesticides with pulverized and built in developers. *Khranit. Prom.*, 20(10), 30-3.
- Serghiou, Constantinos S. 1975. Control of Syringopais temperatella in Cyprus. *J. Econ. Entomol.*, 68(4), 491-4.
- Shafik, Mohammed T.; Bradway, Diane; Enos, Henry F. 1971. Method for confirmation of organophosphorus compounds at the residue level. *Bull. Environ. Contam. Toxicol.*, 6(1), 55-66.
- Sharma, R. P.; Obersteiner, E. J. 1981. Cytotoxic responses of selected insecticides in chick ganglia cultures. *Can. J. Comp. Med.*, 45(1), 60-9.
- Sharma, R. P.; Shupe, J. L.; Potter, Julie R. 1973. Tissue cholinesterase inhibition by 2-carbomethoxy-1-methylvinyl dimethyl phosphate (mevinphos). *Toxicol. Appl. Pharmacol.*, 24(4), 645-52.
- Sharma, R. P., J. L. Shupe and J. R. Potter. 1973. Tissue cholinesterase inhibition by 2-carbomethoxyl-1-methylvinyl dimethyl phosphate (mevinphos). *Toxicol. Appl. Pharmacol.* 24:645-52.
- Sharom, M. S.; Miles, J. R. W.; Harris, C. R.; McEwen, F. L. 1980. Behavior of 12 insecticides in soil and aqueous suspensions of soil and sediment. *Water Res.*, 14(8), 1095-100.
- Sharom, M. S.; Miles, J. R. W.; Harris, C. R.; McEwen, F. L. 1980. Persistence of 12 insecticides in water. *Water Res.*, 14(8), 1089-93.
- Sherman, Martin; Sanchez, Fernando F. 1968. Toxicity of insecticides and acaricides to the papaya. *Hawaii, Agr. Exp. Sta., Tech. Bull.*, 74, 63 pp.
- Shinde, V. K. R.; Verma, J. P.; Sharma, S. K. 1970. Laboratory evaluation of some insecticides against adults of Mylokerus undecimpustulatus maculosus. *Indian J. Entomol.*, 32(Pt. 3), 283-5.
- Shinde, V. K. R.; Verma, J. P.; Sharma, S. K. 1970. Testing of insecticides against Epilachna dodecastigma. *Indian J. Entomol.*, 32(Pt. 3), 218-20.

- Shinde, V. K. R.; Sharma, S. K. 1970. Laboratory evaluation of insecticides against adults of Lachnosterna (Holotrichia) consanguinea blanch (Coleoptera: Scarabaeidae). Madras Agr. J., 57(7), 360-3.
- Shirck, F. H.; Landis, B. J. 1970. Rates of absorption and translocation of some systemic insecticides in broccoli determined by mortality of feeding green peach aphid. J. Econ. Entomol., 63(3), 781-2.
- Silveira Neto, S.; Carvalho, R. P. L.; Barbin, D. 1968. Insecticides for controlling tomato borers. Solo, 60(1), 39-43.
- Simal, J.; Creus, J. M.; Torres, J. M. 1985. Residues of pesticides and polychlorobiphenyls in waters used for supplying Galicia. Tec. Lab., 9(121), 103-10.
- Simpson, G. R.; Penney, D. J. 1974. Pesticide poisonings in the Namoi and Macquarie valleys, 1973. Med. J. Aust., 1(8), 258-60.
- Simpson, G. R. 1973. Aerial spraying of organic phosphate pesticides. Lowered blood cholinesterase levels of aerial spray operators at Wee Waa. Med. J. Aust., 1(15), 735-6.
- Singh, Ashok K.; Hewetson, David W.; Jordon, Kathryn C.; Ashraf, Muhammed. 1986. Analysis of organophosphorus insecticides in biological samples by selective ion monitoring gas chromatography-mass spectrometry. J. Chromatogr., 369(1), 83-96.
- Singh, Harcharan; Singh, Malkiat. 1971. Susceptibility of different larval instars and stages of Euproctis lunata to some contact insecticides. J. Res., Punjab Agr. Univ., 8(2), 217-26.
- Sissons, D. J.; Telling, G. M. 1970. Routine determination of organophosphorus insecticide residues in vegetables. II. Screening procedure for water-soluble insecticides. J. Chromatogr., 48(3), 468-77.
- Skinner, Clint S.; Kilgore, Wendell W. 1982. Application of a dermal self-exposure model to worker reentry. J. Toxicol. Environ. Health, 9(3), 461-81.
- Skinner, Clint S.; Kilgore, Wendell W. 1982. Acute dermal toxicities of various organophosphate insecticides in mice. J. Toxicol. Environ. Health, 9(3), 491-7.
- Slama, K.; Miller, T. A. 1987. Insecticide poisoning: disruption of a possible autonomic function in pupae of Tenebrio molitor. Pestic. Biochem. Physiol., 29(1), 25-34.
- Smilowitz, Zane; Dewey, James E. 1969. Adjuvants prolonging the residual activity of dichlorvos and mevinphos in the laboratory. J. Econ. Entomol., 62(6), 1347-51.

- Sobti, R. C.; Krishan, A.; Pfaffenberger, C. D. 1982. Cytokinetic and cytogenetic effects of some agricultural chemicals on human lymphoid cells in vitro: organophosphates. *Mutat. Res.*, 102(1), 89-102.
- Sonnet, Philip E.; Lye, Thomas L.; Sackett, Richard R. 1978. Effects of selected herbicides on the toxicity of several insecticides to honey bees. *Environ. Entomol.*, 7(2), 254-6.
- Sorensen, K. A.; Kennedy, G. G.; Walgenbach, J. F. 1989. North Carolina Agricultural Chemicals Manual. North Carolina State University. Insect Control for Commercial Vegetable Producers Section, pp 81-106.
- Spencer, E. Y. 1972. Biochemistry and structure of organophosphorus pesticides. in M. A. Kahn and W. O. Haufe (eds), *Toxicology, Biodegradation and Efficacy of Livestock Pesticides*. Swets & Zeitlinger, Amsterdam.
- Spingarn, Neil E.; Northington, D. J.; Pressely, Tom. 1982. Analysis of nonvolatile organic hazardous substances by GC/MS. *J. Chromatogr. Sci.*, 20(12), 571-4.
- Spiro, Sandro; Trevisani, Gian R. 1974. Method for the simultaneous determination of chlorinated and phosphorus-containing pesticides in the air. 2. Organophosphorus derivatives and data on pesticide concentrations in the atmosphere of agricultural zones. *Boll. Lab. Chim. Prov.*, 25(9), 157-64.
- Srivastava, A. S.; Masoodi, M. A. 1985. Relative toxicity of some insecticides to the second instar larvae of Lymantria obfuscata Walker. *Entomon*, 10(3), 241-4.
- Stacherska, Bogumila. 1969. Aphid control on dill grown for green harvest. *Biul. Inst. Ochr. Rosl.*, No. 45, 195-9.
- Stacherska, Bogumila. 1968. Efficiency and the initial action of insecticides against aphids and selectivity for Coccinella septempunctata. *Pr. Nauk. Inst. Ochr. Rosl.*, 10(1), 161-71.
- Staeubli, A.; Haechler, M.; Antonin, P.; Mittaz, C. 1984. Tests on the toxicity of various pesticides towards the natural enemies of the principal [arthropod] pests of pear orchards in Switzerland. *Rev. Suisse Vitic., Arboric., Hortic.*, 16(5), 279-86.
- Stan, H. J.; Christall, B. 1988. Residue analysis of organophosphorus pesticides with capillary gas chromatography and flame photometric detection. Injection of large sample volumes with the cold splitless technique with a combination of programmed temperature vaporization solvent elimination and retention gap. *Dtsch. Lebensm.-Rundsch.*, 84(12), 375-81.
- Stan, H. J.; Mrowetz, D. 1983. Residue analysis of organophosphorus pesticides in food with two-dimensional gas chromatography using

capillary columns and flame photometric detection. HRC CC, J. High Resolut. Chromatogr. Chromatogr. Commun., 6(5),.

- Stan, H. J.; Kellner, G. 1982. Negative chemical ionization mass spectrometry of organophosphorus pesticides. Biomed. Mass Spectrom., 9(11), 483-92.
- Stan, H. J.; Kellner, G. 1982. Analysis of organophosphoric pesticide residues in food by GC/MS using positive and negative chemical ionization. Recent Dev. Food Anal., Proc. Eur. Conf. Food Chem., 1st, Meeting.
- Stan, H. J.; Abraham, B.; Jung, J.; Kellert, M.; Steinland, K. 1977. Detection of organophosphorus insecticides by gas chromatography/mass spectrometry. Fresenius' Z. Anal. Chem., 287(4-5), 271-85.
- Stan, Hans Juergen. 1989. Application of capillary gas chromatography with mass selective detection to pesticide residue analysis. J. Chromatogr., 467(1), 85-98.
- Stan, Hans Juergen; Mrowetz, Dieter. 1983. Residue analysis of pesticides in food by two-dimensional gas chromatography with capillary columns and parallel detection with flame-photometric and electron-capture detection. J. Chromatogr., 279, 173-87.
- Stan, Hans-Juergen; Goebel, Heiderose. 1983. Automated capillary gas chromatographic analysis of pesticide residues in food. J. Chromatogr., 268(1), 55-69.
- Stark, Frank L., Jr. 1967. Pesticide compositions.
- Steen, Jo A.; Hanneman, Gerald D.; Nelson, Peter L.; Folk, Earl D. 1976. Acute toxicity of mevinphos to gerbils. Toxicol. Appl. Pharmacol., 36(1), 195-8.
- Steinwandter, H. 1988. Contributions to the application of gel chromatography in residue analysis. II. A new gel chromatographic system using acetone for the separation of pesticide residues and industrial chemicals. Fresenius' Z. Anal. Chem., 331(5), 499-502.
- Stenseth, Christian. 1979. Effect of fungicides and insecticides on an organophosphorus-compound-resistant strain of the predator mite (*Phytoseiulus persimilis*) Athias-Henriot (Acarina: Phytoseiidae). Forsk. Fors. Landbruket, 30(1), 77-83.
- Stephenson, G. R.; Phatak, S. C.; Makowski, R. I.; Bouw, W. J. 1980. Phytotoxic interactions involving metribuzin and other pesticides in tomatoes. Can. J. Plant Sci., 60(1), 167-75.
- Stevenson, A. B.; Pree, D. J. 1985. Toxicity of insecticides to the aster leafhopper, *Macrostelus fascifrons* (Homoptera: Cicadellidae) in the laboratory and field. Proc. Entomol. Soc. Ont., Volume Date 1984,

115, 89-92.

- Stevenson, J. H. 1978. The acute toxicity of unformulated pesticides to worker honey bees (Apis mellifera L.). Plant Pathol., 27(1), 38-40.
- Stevenson, J. H. 1970. Laboratory and field assessment of pesticide poisoning of honeybees (Apis mellifera). Proc. Brit. Insectic. Fungic. Conf., 5th, Meeting Date 1969, Volume.
- Stijve, T.; Cardinale, E. 1971. Esterase inhibition technique for the detection of organophosphorus pesticides on thin-layer chromatograms. Mitt. Geb. Lebensmittelunters. Hyg., 62(1), 25-31.
- Stoeva, R. 1979. Study on cotton moth (Heliothis armigera Hb /Lep., Noctuidae) biology and assessment of suitable chemicals for its control. Rastenievud. Nauki, 16(8), 92-104.
- Stoltz, R. L. and C. R. Baird. 1986. Control of the spotted alfalfa aphid and pea aphid with foliar sprays in Idaho, 1985. Insect. and Acar. Tests 11: 218.
- Straka, F. 1983. Toxicity of some insecticides to Mamestra brassicae L. and M. oleracea L. Gradinar. Lozar. Nauka, 20(5), 72-81.
- Studzinski, Andrzej; Mikolajewicz, Maria. 1976. Study of the use of preparations Actellic 50 EC, Phosdrin 24 EC and Polfos in the control of Lygus species on fennel (Foeniculum vulgare Mill). Herba Pol., 22(3-4), 307-11.
- Su, Mei-Quey; Kinoshita, Florence K.; Frawley, John P.; DuBois, Kenneth P. 1971. Comparative inhibition of aliesterases and cholinesterase in rats fed eighteen organophosphorus insecticides. Toxicol. Appl. Pharmacol., 20(2), 241-9.
- Suber, E. F.; Chalfant, Richard B.; Canerday, T. Don. 1971. Toxicity of insecticides to the cowpea curculio in the laboratory. J. Econ. Entomol., 64(5), 1080-1.
- Sun, C. N.; Wu, T. K.; Chen, J. S.; Lee, W. T. 1986. Insecticide resistance in diamondback moth. Diamondback Moth Manage., Proc. Int. Workshop, 1st, Meeting Date 1985.
- Sun, Yun-Pei; Horng, Wei-Huai; Hsieh, Peng-Sheng; Wang, You-Chong; Lee, Sung-Lien. 1977. A simple method of establishing tolerances for pesticide residues in different countries. Arch. Environ. Contam. Toxicol., 5(3), 279-93.
- Sun, Yun-Pei. 1971. Speed of action of insecticides and its correlation with accumulation in fat and excretion in milk. J. Econ. Entomol., 64(3), 624-30.

- Svampa, Giorgio; Brunelli, Agostino; Mariani Tosatti, Enrico. 1974. Phytopharmaceutical mixtures. Effects of insecticides on fungicides. *Inf. Fitopatol.*, 24(12), 5-10, 43-5.
- Svobodova, Zdenka. 1980. Acute toxicity of pesticides to fish. *Agrochemia (Bratislava)*, 20(11), 328-32.
- Swift, John E. 1976. Organophosphate exposure from agricultural usage. U. S. Environ. Prot. Agency, Off. Res. Dev., [Rep.] EPA,.
- Szalontai, Gabor. 1976. High-performance liquid chromatography of organophosphorus insecticides. *J. Chromatogr.*, 124(1), 9-16.
- Szczepanska, Krystyna; Kowalska, Tatiana. 1969. Control of Agrotinae (Lepidoptera, Noctuidae, Agrotinae). II. Effectiveness of phosphoorganic and carbaminated insecticides against Agrotinae caterpillars. *Pr. Nauk. Inst. Ochr. Rosl.*, 11(2), 141-50.
- Szeberenyi, J.; Varga, A.; Pongracz, M. F.; Molnar, J. 1972. Centrally mediated contraction of the lower eyelid elicited by anticholinesterases in anesthetized rats. *J. Pharm., Pharmacol.*, 24(1), 85-6.
- Tabashnik, Bruce E.; Cushing, Nancy L.; Johnson, Marshall W. 1987. Diamondback moth (Lepidoptera: Plutellidae) resistance to insecticides in Hawaii: intra-island variation and cross-resistance. *J. Econ. Entomol.*, 80(6), 1091-9.
- Takats, Gabor. 1974. Three-year study of phosphoric ester poisoning. *Orv. Hetil.*, 115(38), 2240-2.
- Taylor, David G. 1980. NIOSH manual of analytical methods. DHHS (NIOSH) Publ. (U. S.), 80-125, 264 pp.
- Teetes, George L.; Randolph, Neal M. 1969. Chemical and cultural control of the sunflower moth in Texas. *J. Econ. Entomol.*, 62(6), 1444-7.
- Tejada, A. W.; Magallona, E. D.; Lakan-Ilaw, E. B. 1983. Insecticide residues in vegetables: application of the modified approach to organophosphate insecticide residues in string bean (Vigna sesquipedalis Fruw.). *Philipp. Agric.*, 66(4), 405-16.
- Tejada, A. W.; Varca, L. M.; Magallona, E. D. 1977. Insecticide residues in vegetables. II. Effect of dosage rates and preharvest intervals on residues in and on green beans (*Phaseolus vulgaris* L.). *Philipp. Entomol.*, 3(5-6), 273-80.
- Teotia, T. P. S.; Lal, Beche. 1973. Relative toxicity of some insecticides to the larvae of til leaf and pod borer, Antigastra catalaunalis. *Indian J. Entomol.*, Volume Date 1972, 33, Pt. 3, 257-9.

- Teotia, T. P. S.; Tiwari, G. C. 1972. Toxicity of some important insecticides to the coccinellid predator, Coccinella septempunctata. Labdev, Part B, 10(1), 17-18.
- Teotia, T. P. S.; Nigam, P. M. 1967. Relative toxicity of some insecticides as contact poisons to the larvae of Earias fabia. Indian J. Entomol., 29(Pt. 4), 391-2.
- Thompson, John F.; Mann, Jon B.; Apodaca, Alfred O.; Kantor, Edward J. 1975. Relative retention ratios of ninety-five pesticides and metabolites on nine gas-liquid chromatographic columns over a temperature range of 170 to 204.deg. in two detection modes. J. Assoc. Off. Anal. Chem., 58(5), 1037-50.
- Thomson, W. T. 1977. Agricultural Chemicals. Book I, Insecticides. Thomson Publications. Fresno, CA.
- Tieman, Charles H.; Pollard, Glenn E.; Stoutamire, Donald W.; Silverman, Milton. 1969. Dialkyl phosphinyloxy-3-butenates isoenolphosphates.
- Timar, Miklos. 1976. Questions of hygiene concerning the utilization of plant pesticides. Agrartud. Kozl., 35(1-3), 9-23.
- Torchio, Philip F. 1973. Relative toxicity of insecticides to the honey bee, alkali bee, and alfalfa leafcutting bee (Hymenoptera, Apidae, Halictidae, Megachilidae). J. Kans. Entomol. Soc., 46(4), 446-53.
- Tsvetkova, Ts.; Khristodorova, V.; Sergeeva, D. 1974. Enzymic thin-layer chromatographic method for the identification of some organophosphorus pesticides. Vet.-Med. Nauki, 11(6), 50-3.
- Twine, P. H. 1971. Insecticide trial against corn aphid (Rhopalosiphum maidis) on sorghum. Queensl. J. Agr. Anim. Sci., 28(1), 19-22.
- Ungaro, Maria Thereza S.; Guindani, Cleusa M. A.; Ferreira, Marilene da S.; Pigati, P.; Takematsu, A. P.; Kastrup, L. F. C.; Ishizaki, T. 1980. Residues of phosphorated and chlorinated insecticides in fruits and vegetables. Biologico, 46(7), 129-34.
- United States Department of Agriculture. 1965. Injurious crop insects. pp 40-54 in losses in agriculture. USDA Agr. Handb. 291:120 pp.
- United States Department of Agriculture. 1988. Vegetables and specialties, situation and outlook yearbook. Economic Research Service, TVS-246,
- United States Department of Agriculture. National Agricultural Statistics Service. Crop Production. Annual issues.
- United States Department of Agriculture. National Agricultural Statistics Service. Crop Values. Annual issues.

- United States Department of Commerce. 1982. Census of Agriculture. National Summary.
- Urfine Fogarasi, Eva; Ambrus, Arpad. 1980. Rate of metabolism and study of pesticide residues in fennel and mustard treated with lindane, mevinphos and trifluraline. *Novenyvedelem* (Budapest), 16(11), 556-62.
- Valaskova, Eva. 1979. Complex plant protection measures against diseases and pests of gerberas. *Gradinarstvo*, 60(12), 35-7.
- Vanurova, Eliska. 1970. Activity of some preparations against Malacosoma neustria caterpillars. *Agrochemia*, 10(3), 88-91.
- Vanwetswinkel, G.; Seutin, E. 1978. Study of some photostable pyrethroids in fruit arboriculture. *Phytiatr.-Phytopharm.*, 27(1), 15-26.
- Varnagy, Laszlo. 1981. Teratological examination of agricultural pesticides on Japanese quail (Coturnix coturnix Japonica) eggs. *Acta Vet. Acad. Sci. Hung.*, 29(1), 77-83.
- Varnai, Laszlo. 1968. Intoxication by organic phosphoric acid esters. *Wiss. Beitr., Martin-Luther-Univ. Halle-Wittenberg*, No. 12, 149-51.
- Venant, A.; Borrel, S.; Mallet, J.; Van Neste, E. 1989. Gel permeation chromatography as a method for the simultaneous clean up of organochlorine, organophosphate and polychlorinated biphenyl residues in fat extracts. *Analysis*, 117(1-2), 64-6.
- Verberk, M. M.; Salle, H. J. A. 1979. Neurological effects and cholinesterase inhibition in man by subacute administration of mevinphos. *Act. Nerv. Super.*, Volume Date 1977, 19(Suppl. 2), 333-4.
- Verberk, M. M. 1977. Incipient cholinesterase inhibition in volunteers ingesting monocrotophos or mevinphos for one month. *Toxicol. Appl. Pharmacol.*, 42(2), 345-50.
- Verberk, M. M.; Salle, H. J. A. 1977. Effects of nervous function in volunteers ingesting mevinphos for one month. *Toxicol. Appl. Pharmacol.*, 42(2), 351-8.
- Verma, A. N.; Verma, N. D.; Khurana, A. D. 1973. Chemical control of the diamond-back moth, Plutella xylostella, infesting cauliflower in Haryana. *Indian J. Entomol.*, Volume Date 1972, 34, Pt. 3, 206-12.
- Verma, A. N.; Sharma, P. D.; Saramma, P. U. 1969. Relative toxicity of some contact insecticides against cabbage caterpillar, Pieris brassicae (Pieridae; Lepidoptera). *J. Res., Punjab Agr. Univ.*, 6(1)(Suppl.), 197-9.
- Verma, A. N.; Saramma, P. U.; Ramzan, M. 1969. Relative toxicity of different insecticides to the adults of cotton grey weevil, Myliocerus undecimpustulatus maculosus (Curculionidae: coleoptera). *J. Res.*,

Punjab Agr. Univ., 6(1)(Suppl.), 170-3.

Verma, M. B.; Sharma, J. C. 1972. Relative toxicity of certain insecticides to the adult grasshopper, Chloeobora crassa. (Orthoptera: Acridiidae). Labdev, Part B, 10(2), 93.

Vernon, J. D. R. 1971. Biology and control of tortricid larvae on strawberries. Plant Pathol., 20(2), 73-80.

Vicelli, Regina C.; Beraldo, Maria J. A. H. 1981. Laboratory determination of the toxicity of organophosphorus and carbamate insecticides to Atta laevigata (F. Smith, 1858) and Atta sexdens rubropilosa Forel, 1908 (Hymenoptera: Formicidae). Arq. Inst. Biol., Sao Paulo, 48(1-4), 1-4.

Vighi, M.; Calamari, D. 1987. A triparametric equation to describe QSARs for heterogeneous chemical substances. Chemosphere, 16(5), 1043-51.

Vilceanu, R.; Chiriac, A.; Szabadai, Z.; Simon, Z. 1977. Structure-activity correlations for the toxicity of organic phosphorus compounds of the Schrader type. An. Univ. Timisoara, Ser. Stiinte Fiz.-Chim., 15(1), 61-77.

Vilceanu, R.; Chiriac, A.; Szabaday, Z.; Simon, Z. 1962. Multiple correlations in the anticholinesterasic activity of organic phosphorus compounds. Rev. Roum. Biochim., 10(3), 239-52.

Vilceanu, R.; Szabadai, Z.; Chiriac, A.; Simon, Z. 1972. Multiple structure-toxicity correlation of organic phosphorus compounds. Stud. Biophys., 34(1), 1-6.

Villeneuve, David C.; Butterfield, A. G.; Grant, D. L.; McCully, Keith A. 1970. Detection, separation, and quantitative recovery of thirteen organophosphorus pesticides on Silica Gel GF254 thin-layer chromatograms. J. Chromatogr., 48(3), 567-71.

Virginia Agricultural Statistics Service. 1988. Virginia Crops and Livestock. 60(24): 2.

Visi, Gyorgy, Mrs. 1984. Study of the concentration decrease of pesticides in aquatic model systems. Novenyvedelem (Budapest), 20(8), 358-63.

Vora, Kishore Nyalchand; Campbell, Donald Noble; Davis, Robert Clinton, Jr.; Spangler, Glenn Edward; Reategui, Julio Antonio. 1986. Selective ionization of gas constituents using electrolytic reactions.

Voss, G. 1981. Taxonomy-related cholinesterase inhibition patterns in insects. J. Econ. Entomol., 74(5), 555-7.

Vrochinskii, K. K. 1975. Herbicides as a potential factor of the pollution in reservoirs. Mekh. Deistviya Gerbits. Sint. Regul. Rosta

Rast. Ikh Sud'ba.

- Wachendorff, Ulrike; Klingauf, F. 1978. Esterase assay for the diagnosis of insecticide resistance in aphids. *Z. Pflanzenkr. Pflanzenschutz*, 85(3-4), 218-27.
- Walgenbach, J. F., ed. 1988. Agricultural Chemicals for North Carolina Apples. AG-37. The North Carolina Agricultural Extension Service. North Carolina State University. 52 pp.
- Walker, Norman E. 1968. Use of yolk-chemical mixtures to replace hen egg yolk in toxicity and teratogenicity studies. *Toxicol. Appl. Pharmacol.*, 12(1), 94-104.
- Wang, Qinsun; Chen, Duanmei. 1987. Optimization of gas chromatographic separation of organophosphorus pesticides. *Sepu*, 5(3), 171-4.
- Wang, Ren Tse; Chou, Shuin Shiou. 1969. Simple and rapid reversed phase thin-layer chromatography of pesticides. *Hua Hsueh*, (4), 80-5.
- Wang, Ren Tse; Chou, Shuin Shiou. 1968. Detection of pesticides by reversed phase paper chromatography. *Chung Kuo Nung Yeh Hua Hsueh Hui Chih*, 6(3-4), 63-8.
- Wang, T. C.; Feng, H. T. 1986. Diamondback moth resistance and cross resistance to four commonly used insecticides in Taiwan. *Bull. Inst. Zool., Acad. Sin.*, 25(1), 99-104.
- Watanabe, P. G.; Sharma, R. P. 1972. Inhibition of subcellular cholinesterases by 2-carbomethoxy-1-methylvinyl dimethyl phosphate (mevinphos) in vitro and in vivo. *Toxicol. Appl. Pharmacol.*, 23(4), 692-700.
- Wegman, R. C. C.; Van den Broek, H. H.; Hofstee, A. W. M.; Marsman, J. A. 1984. Determination of triazines, organophosphorus containing pesticides and aromatic amines in soil samples. *Meded. Fac. Landbouwwet., Rijksuniv. Gent*, 49(3b), 1231-9.
- White, Matthew M.; Mane, Suresh D.; Richmond, Rollin C. 1988. Studies of esterase 6 in Drosophila melanogaster. XVIII. Biochemical differences between the slow and fast allozymes. *Mol. Biol. Evol.*, 5(1), 41-62.
- Wiackowski, S. K.; Herman, E. 1971. Laboratory studies of the effect of chemical plant protectants on Encarsia formosa (Hymenoptera, Aphelinidae) brought from Canada for biological control of Trialeurodes vaporariorum (Homoptera, Aleurodidae). *Pr. Inst. Sadow. Skierniewicach*, 15, 181-3.
- Wiackowski, Stanislaw K.; Herman, Ewa. 1968. Effect of insecticides on adults of primary and secondary aphid parasites. *Pol. Pismo Entomol.*, 38(3), 593-600.

- Wiackowski, Stanislaw K.; Nowacka, Barbara. 1968. Laboratory investigations on the effect of insecticides on the larvae and adults of Coccinella septempunctata. Pol. Pismo Entomologiczne. Bull. Entomol. Pol., 38(2), 441-52.
- Wieres, R., and J. Leeper. 1980. Rosy Apple Aphid. Tree Fruit IPM, Insect Identification Sheet No. 11. New York State Agricultural Experiment Station, Geneva, NY.
- Wilkinson, Robert E.; Glaze, Norman C.; Dowler, Clyde C.; Young, Clyde T. 1979. Turnip green, cucumber, snapbean, and southern pea response to pesticides in intensive-cropping sequences. J. Agric. Food Chem., 27(4), 900-5.
- Williams, Phletus P.; Stolzenberg, Rosa L. 1972. Ruminal bacterial degradation of benzo(b)thien-4-yl methylcarbamate (Mobam) and effect of Mobam on ruminal bacteria. Appl. Microbiol., 23(4), 745-9.
- Winterlin, Wray; Mourer, Charles; Beckman, Herman. 1970. Analysis of phosdrin in vegetables using gas-liquid chromatography. J. Agr. Food Chem., 18(3), 401-4.
- Winterlin, Wray; Walker, Glenn; Frank, Hubert. 1968. Detection of cholinesterase-inhibiting pesticides following separation on thin-layer chromatograms. J. Agr. Food Chem., 16(5), 808-12.
- Wiswesser, W. J. (ed.). Pesticide index. Entomological Society of America. College Park, MD.
- Wolverton, B. C.; Wallace, W. J.; Young, A. L.; Harrison, D. D. 1969. Systemic uptake of toxic phosphorus esters by plants: morphological effects of foliar applications of the organophosphate insecticides mevinphos and methyl demeton on selected plant species. U.S. Clearinghouse Fed. Sci. Tech. Inform., AD, No. 700052, 28 pp.
- Wood, E. A., Jr. 1971. Insecticidal control of the greenbug. J. Econ. Entomol., 64(3), 704-7.
- Woodrow, James E.; Wong, Jennifer M.; Seiber, James N. 1989. Pesticide residues in spray aircraft tank rinses and aircraft exterior washes. Bull. Environ. Contam. Toxicol., 42(1), 22-9.
- Woodrow, James E.; Seiber, James N. 1978. Portable device with XAD-4 resin trap for sampling airborne residues of some organophosphorus pesticides. Anal. Chem., 50(8), 1229-31.
- Workman, R. B. 1978. Pesticides toxic to striped earwig, an important insect predator. Proc. Fla. State Hortic. Soc., Volume Date 1977, 90, 401-2.
- Workman, Ralph B.; Greene, Gerald L. 1970. Control of aphids on rutabaga at two locations in Florida. Fla. Entomol., 53(3), 135-7.

- Wuethrich, C.; Mueller, F.; Blaser, O.; Marek, B. 1985. Pesticides and other chemical residues in Swiss diet samples. Mitt. Geb. Lebensmittelunters. Hyg., 76(2), 260-76.
- Yassin, A. M. 1975. Epidemics and chemical control of leaf curl virus disease of tomato in the Sudan. Exp. Agric., 11(3), 161-5.
- Yawalkar, K. S.; Chitriv, A. J. 1969. Singhara beetle (Galerucella birmania) and its control. Pesticides, 3(10), 26-8.
- Young, J. R.; Harrell, E. A.; Hare, W. W. 1972. Mortality of adult corn earworms treated with insecticidal formulations in sweet corn in fields and in the laboratory. J. Econ. Entomol., 65(3), 786-9.
- Yu, Shyi Jian; Morrison, Frank Orville. 1969. Mevinphos and phosphamidon: influence of plant nutrition on uptake and effectiveness. J. Econ. Entomol., 62(6), 1296-304.
- Zeff, Jack D.; Leitis, Eriks; Harris, John A. 1984. Chemistry and application of ozone and ultraviolet light for water reuse - pilot plant demonstration. Proc. Ind. Waste Conf., Volume Date 1983, 38th, 105-16.
- Zeleny, Jiri; Hrdy, Ivan. 1969. Over populations of hop aphid, Phorodon humuli, in 1967; population dynamics in 1968. Effect of insecticides on hop aphids and entomocenosis of hop yards. Agrochemia, 9(8), 211-15.
- Zeleny, Jiri; Hrdy, Ivan. 1969. Population explosion of Phorodon humuli in 1967, population dynamics in 1968, effect of insecticides on Phorodon humuli, and entomological picture of hops. II. Agrochemia, 9(9-10), 257-61.
- Zhang, Qiao; Li, Weige; Huang, Shizhong; Zhai, Yanlu; Li, Zhixiang. 1988. GC method for simultaneous determination of organophosphorus pesticides. Huanjing Kexue Xuebao, 8(3), 367-72, 1 plate.
- Zhang, Qiao; Li, Weige; Xie, Faming; Han, Hauqiong; He, Youjie. 1987. Gas-chromatographic method for multi-residue determination of nine organophosphorus insecticides. Zhongguo Nongye Kexue (Beijing), 20(1), 75-80.
- Zotov, E. P.; Altymyshev, A. A. 1979. Recognition of patterns. Use of discriminant analysis for predicting the biological activity of organophosphorus compounds. Izv. Akad. Nauk Kirg. SSR, (6), 38-42.
- Zoun, P. E. F.; Spierenburg, Th. J. 1989. Determination of cholinesterase-inhibiting pesticides and some of their metabolites in cases of animal poisoning using thin-layer chromatography. J. Chromatogr., 462, 448-53.

APPENDIX I

Mevinphos (Phosdrin) Questionnaire

(Please provide a separate questionnaire for each registered crop and/or use in your State). Reporting State: _____

USE EXTRA SHEETS AS NEEDED FOR ANY QUESTION.

1. Crop/site: _____
 - 1a. Please provide the following information regarding this crop:
Total acres of crop (Average 1982-1986) _____ Acres
Total production (Average 1982-1986), specify units _____
Total crop value (Average 1982-1986) \$ _____
 2. Target Pest(s) _____

 3. Average number of acres (or please state other appropriate units) treated with mevinphos per year. Average for 1982-1986 _____.
 4. Average number of treatments per acre (or please state other appropriate units) for all pests on this crop. Average for 1982-1986 _____.
 5. Rate, active ingredient per acre per application (or please state other appropriate unit). Average for 1982-1986 _____.
 6. Cost per acre (or please state other appropriate unit) per application
 - a. Material \$ _____
 - b. Application (include labor plus machinery costs or custom rate)
\$ _____
 7. Is mevinphos applied as part of an operation that would take place regardless of whether or not mevinphos is applied (e.g. applied in conjunction with regular fungicide application, etc.)?
(Yes or No): _____.
- If yes, please describe the operation: _____

9. a. Please give the limitations or disadvantages of each alternative (e.g. longer post harvest interval, controls fewer pests and name pests controlled by each alternative, resistance, hazard to user and/or non-target organisms, environmental concerns, etc.): _____

- b. What is the degree of each limitation (e.g. severe, mild, insignificant): _____

- c. Advantages, if any, of each alternative relative to mevinphos:

10. Benefits of mevinphos vs. registered alternatives:

- a. Given the use of alternative practices above, estimate the percent change in yield or marketable product per acre on acreage currently treated with mevinphos. Indicate yield loss or gain:

- b. Quality effects, type and degree: _____

c. Other (e.g. short Post Harvest Interval, Resistance Management, etc.)

(Please provide references and/or data on an additional sheet or sheets where available for question 10. Best guesses should be provided if that is all that is available).

11. Please provide any available economic data on use and benefits of mevinphos and/or alternatives (or indicate other data sources you are aware of).

12. Does your state recommend mevinphos or alternatives (please specify which ones) and enclose a copy of your recommendations.

13. Do you perceive a need for (specific needs for pest) registration of mevinphos on this crop (site)? Please indicate extent of need (e.g. essential, important, limited, not needed). State your reasons.

14. Your Name _____
Address _____

Telephone _____

If you have questions contact Bobby Pass at 606-257-7450.

Please return completed questionnaire (s) by August 15, 1988 to:

Bobby C. Pass
Department of Entomology
University of Kentucky
Lexington, Kentucky 40546-0091

APPENDIX II

Survey responses by commodity: States reporting, acreage and acreage treated with mevinphos, mevinphos applied, mevinphos's yield benefit, and control costs with and without mevinphos available, 1982-86

U.S. production by commodity: Mevinphos-treated farm production and yield benefit, U.S. aggregate farm production, farm price, and yield benefit, by crop, 1982-86

Survey responses by commodity: Acreage, production, value, 1982-1986

Appendix Table II.1. Survey responses for vegetables: Acreage and acreage treated with mevinphos, mevinphos applied, mevinphos's yield benefit, and control costs with and without mevinphos available (average annual), 1982-86.

Crop and State	Acreage Reported		Mevinphos applied ¹ (pounds a.i.)	Yield Benefit ² (percent)	Control Costs (dollars/acre)	
	Total	Treated			Current	Alt ³
Broccoli	112,622	61,892	71,764	12.2	25.53	23.06
Cabbage	83,158	46,310	63,597	18.5	13.09	54.78
Muskmelons	55,230	38,463	17,271	11.2	13.77	13.71
Carrot	15,060	3,535	1,326	3.7	7.08	9.01
Cauliflower	51,337	34,400	35,445	40.3	23.31	8.60
Celery	30,973	15,743	19,143	27.3	37.77	33.97
Greens	33,142	24,985	69,667	5.3	7.20	71.06
Lettuce	233,496	209,500	241,748	12.1	26.78	18.46
Onion	10,110	5,550	6,300	23.6	32.17	13.82
Pepper	38,100	17,769	5,284	1.3	8.14	15.24
Potato	32,900	658	247	0.0	9.90	25.66
Spinach	22,299	13,910	16,040	14.2	14.64	9.62
Sweet corn	67,020	12,712	7,060	10.7	16.75	8.36
Tomato	58,370	6,922	3,543	2.3	11.10	14.37
Watermelon	109,610	13,145	9,323	1.8	10.56	21.56
Subtotal	953,427	505,493	567,756	13.8	21.40	23.59
Other	311,275	41,574	27,021	12.2	11.79	18.85
Total	1,264,702	547,067	594,777	13.7	20.67	23.23

¹Total pounds active ingredient (a.i.) applied equals average number of applications times average application rate times acres treated.

²Yield benefit equals the percent change in yield or marketable product per acre on acreage currently treated with mevinphos (versus registered alternatives).

³Average cost of chemical and/or nonchemical control alternatives that would be used on acreage currently treated with mevinphos.

Appendix Table II.2. Survey responses for field crops: States reporting, acreage and acreage treated with mevinphos, mevinphos applied, mevinphos's yield benefit, and control costs with and without mevinphos available (average annual), 1982-86.

Crop and State	Acreage Reported		Mevinphos applied ¹ (pounds a.i.)	Yield Benefit ² (percent)	Control Costs (dollars/acre)	
	Total	Treated			Current	Alt ³
Alfalfa, seed						
California	100,000	84,300	48,051	1.0	19.13	10.00
Oregon	8,862	2,500	1,250	1.0	12.75	10.00
Washington	25,000	6,000	7,500	12.5	60.00	10.00
Total	133,862	92,800	56,801	1.5	18.75	10.00
Corn, field						
Colorado	660,000	320	160	---	8.49	30.00
Florida	46,200	500	250	---	7.00	4.79
Total	706,200	820	410	---	7.58	14.63
Sorghum, forage						
Florida	25,000	1,250	625	7.5	7.00	4.17
Sorghum, grain						
Florida	20,045	450	225	2.5	7.00	3.91
Total	45,045	1,700	850	7.2	7.00	4.10
Total	885,107	95,320	58,061	2.9	18.44	9.93

¹Total pounds active ingredient (a.i.) applied equals average number of applications times average application rate times acres treated.

²Yield benefit equals the percent change in yield or marketable product per acre on acreage currently treated with mevinphos (versus registered alternatives).

³Average cost of chemical and/or nonchemical control alternatives that would be used on acreage currently treated with mevinphos.

Appendix Table II.3. Survey responses for fruit crops: States reporting, acreage and acreage treated with mevinphos, mevinphos applied, mevinphos's yield benefit, and control costs with and without mevinphos available (average annual), 1982-86.

Crop and State	Acreage Reported		Mevinphos applied ¹ (pounds a.i.)	Yield Benefit ² (percent)	Control Costs (dollars/acre)	
	Total	Treated			Current	Alt ³
Apple						
Illinois	14,000	1,500	450	0.0	22.50	15.00
New Jersey	5,280	40	90	0.0	37.50	10.00
New York	64,400	3,200	3,200	1.5	31.00	35.50
Total	83,680	4,740	3,740	1.3	28.36	28.80
Grape						
California	676,700	11,419	6,509	6.0	18.38	10.00
Strawberry						
California	13,342	3,000	3,420	60.0	41.25	194.24
Florida	5,160	5,160	90,300	15.0	261.98	194.24
New Jersey	600	17	9	0.0	10.00	10.00
Total	19,102	8,177	93,729	44.6	24.60	193.86

¹Total pounds active ingredient (a.i.) applied equals average number of applications times average application rate times acres treated.

²Yield benefit equals the percent change in yield or marketable product per acre on acreage currently treated with mevinphos (versus registered alternatives).

³Average cost of chemical and/or nonchemical control alternatives that would be used on acreage currently treated with mevinphos.

Appendix Table II.4. Survey responses for broccoli: States reporting, acreage and acreage treated with mevinphos, mevinphos applied, mevinphos's yield benefit, and control costs with and without mevinphos available (average annual), 1982-86.

Crop and State	Acreage Reported		Mevinphos applied ¹ (pounds a.i.)	Yield Benefit ² (percent)	Control Costs (dollars/acre)	
	Total	Treated			Current	Alt ³
New Jersey	400	65	16	---	7.50	10.00
Alabama	600	300	750	25.0	35.00	37.00
Florida	236	71	27	7.5	6.92	14.33
Texas	7,500	7,500	15,000	---	14.50	14.33
Arizona	10,874	12,427	24,854	---	33.50	52.40
California	90,700	40,372	30,683	17.5	24.00	14.33
Oregon	2,312	1,156	434	---	24.00	15.70
Total	112,622	61,892	71,764	12.2	24.39	22.10

¹Total pounds active ingredient (a.i.) applied equals average number of applications times average application rate times acres treated.

²Yield benefit equals the percent change in yield or marketable product per acre on acreage currently treated with mevinphos (versus registered alternatives).

³Average cost of chemical and/or nonchemical control alternatives that would be used on acreage currently treated with mevinphos.

Appendix Table II.5. Survey responses for cabbage: States reporting, acreage and acreage treated with mevinphos, mevinphos applied, mevinphos's yield benefit, and control costs with and without mevinphos available (average annual), 1982-86.

Crop and State	Acreage Reported		Mevinphos applied ¹ (pounds a.i.)	Yield Benefit ² (percent)	Control Costs (dollars/acre)	
	Total	Treated			Current	Alt ³
Delaware	850	500	438	---	31.50	10.00
New Jersey	3,000	484	121	---	7.50	10.00
New York	12,000	4,000	2,000	20.0	15.00	10.00
Illinois	3,500	800	800	---	10.50	10.00
Wisconsin	5,500	1,200	1,125	4.0	30.00	10.00
Alabama	700	100	250	25.0	35.00	36.80
Florida	14,190	10,461	11,769	30.0	18.75	52.40
Georgia	7,000	5,000	5,700	55.0	20.10	72.00
North Carolina	12,000	500	3,000	10.0	45.78	47.52
South Carolina	500	475	1,425	50.0	42.00	75.75
Tennessee	1,500	1,000	1,124	---	18.96	47.52
Texas	16,040	16,040	32,080	---	14.50	72.00
California	6,178	5,550	3,691	40.0	23.63	52.40
Utah	200	200	75	25.0	12.00	10.00
Total	83,158	46,310	63,597	18.5	13.09	54.78

¹Total pounds active ingredient (a.i.) applied equals average number of applications times average application rate times acres treated.

²Yield benefit equals the percent change in yield or marketable product per acre on acreage currently treated with mevinphos (versus registered alternatives).

³Average cost of chemical and/or nonchemical control alternatives that would be used on acreage currently treated with mevinphos.

Appendix Table II.6. Survey responses for carrot: States reporting, acreage and acreage treated with mevinphos, mevinphos applied, mevinphos's yield benefit, and control costs with and without mevinphos available (average annual), 1982-86.

Crop and State	Acreage Reported		Mevinphos applied ¹ (pounds a.i.)	Yield Benefit ² (percent)	Control Costs (dollars/acre)	
	Total	Treated			Current	Alt ³
Wisconsin	4,300	250	94	---	18.00	10.00
Florida	10,760	3,285	1,232	5.0	6.25	8.94
Total	15,060	3,535	1,326	3.7	7.08	9.01

¹Total pounds active ingredient (a.i.) applied equals average number of applications times average application rate times acres treated.

²Yield benefit equals the percent change in yield or marketable product per acre on acreage currently treated with mevinphos (versus registered alternatives).

³Average cost of chemical and/or nonchemical control alternatives that would be used on acreage currently treated with mevinphos.

Appendix Table II.7. Survey responses for cauliflower: States reporting, acreage and acreage treated with mevinphos, mevinphos applied, mevinphos's yield benefit, and control costs with and without mevinphos available (average annual), 1982-86.

Crop and State	Acreage Reported		Mevinphos applied ¹ (pounds a.i.)	Yield Benefit ² (percent)	Control Costs (dollars/acre)	
	Total	Treated			Current	Alt ³
New Jersey	350	57	14	---	7.50	10.00
Florida	1,200	285	214	---	11.00	8.38
Texas	1,800	1,800	3,600	---	14.50	8.38
California	44,567	31,232	31,232	45.0	24.50	8.38
Oregon	3,420	1,026	385	---	24.00	15.70
Total	51,337	34,400	35,445	40.3	23.31	8.60

¹Total pounds active ingredient (a.i.) applied equals average number of applications times average application rate times acres treated.

²Yield benefit equals the percent change in yield or marketable product per acre on acreage currently treated with mevinphos (versus registered alternatives).

³Average cost of chemical and/or nonchemical control alternatives that would be used on acreage currently treated with mevinphos.

Appendix Table II.8. Survey responses for celery: States reporting, acreage and acreage treated with mevinphos, mevinphos applied, mevinphos's yield benefit, and control costs with and without mevinphos available (average annual), 1982-86.

Crop and State	Acreage Reported		Mevinphos applied ¹ (pounds a.i.)	Yield Benefit ² (percent)	Control Costs (dollars/acre)	
	Total	Treated			Current	Alt ³
Michigan	3,100	3,100	1,163	2.0	10.75	9.70
Florida	9,040	1,080	405	2.5	5.75	39.92
California	18,833	11,563	17,576	37.5	48.00	39.92
Total	30,973	15,743	19,143	27.3	37.77	33.97

¹Total pounds active ingredient (a.i.) applied equals average number of applications times average application rate times acres treated.

²Yield benefit equals the percent change in yield or marketable product per acre on acreage currently treated with mevinphos (versus registered alternatives).

³Average cost of chemical and/or nonchemical control alternatives that would be used on acreage currently treated with mevinphos.

Appendix Table II.9. Survey responses for greens (collards, turnips, kale, mustard): States reporting, acreage and acreage treated with mevinphos, mevinphos applied, mevinphos's yield benefit, and control costs with and without mevinphos available (average annual), 1982-86.

Crop and State	Acreage Reported		Mevinphos applied ¹ (pounds a.i.)	Yield Benefit ² (percent)	Control Costs (dollars/acre)	
	Total	Treated			Current	Alt ³
New Jersey	500	81	20	---	7.50	10.00
Illinois	2,500	500	750	---	17.00	12.75
Alabama	1,200	400	1,000	25.0	35.00	60.40
Florida	1,269	381	191	7.5	9.10	5.91
Georgia	17,000	17,000	44,625	---	23.45	72.00
Louisiana	2,000	250	438	---	24.50	60.40
North Carolina	4,000	2,000	16,000	10.0	61.04	109.60
South Carolina						
greens	1,000	950	1,900	50.0	28.00	52.50
collards	1,000	950	3,325	50.0	49.00	85.75
Tennessee	1,200	1,200	900	---	39.96	109.60
Virginia	473	473	118	17.0	8.25	10.00
Oklahoma	1,000	800	400	10.0	10.40	20.00
Total	33,142	24,985	69,667	5.3	7.20	71.06

¹Total pounds active ingredient (a.i.) applied equals average number of applications times average application rate times acres treated.

²Yield benefit equals the percent change in yield or marketable product per acre on acreage currently treated with mevinphos (versus registered alternatives).

³Average cost of chemical and/or nonchemical control alternatives that would be used on acreage currently treated with mevinphos.

Appendix Table II.10. Survey responses for sweet corn: States reporting, acreage and acreage treated with mevinphos, mevinphos applied, mevinphos's yield benefit, and control costs with and without mevinphos available (average annual), 1982-86.

Crop and State	Acreage Reported		Mevinphos applied ¹ (pounds a.i.)	Yield Benefit ² (percent)	Control Costs (dollars/acre)	
	Total	Treated			Current	Alt ³
Florida	53,020	5,100	1,275	---	5.17	8.36
California	14,000	7,612	5,785	17.5	24.50	8.36
Total	67,020	12,712	7,060	10.7	16.75	8.36

¹Total pounds active ingredient (a.i.) applied equals average number of applications times average application rate times acres treated.

²Yield benefit equals the percent change in yield or marketable product per acre on acreage currently treated with mevinphos (versus registered alternatives).

³Average cost of chemical and/or nonchemical control alternatives that would be used on acreage currently treated with mevinphos.

Appendix Table II.11. Survey responses for muskmelons: States reporting, acreage and acreage treated with mevinphos, mevinphos applied, mevinphos's yield benefit, and control costs with and without mevinphos available (average annual), 1982-86.

Crop and State	Acreage Reported		Mevinphos applied ¹ (pounds a.i.)	Yield Benefit ² (percent)	Control Costs (dollars/acre)	
	Total	Treated			Current	Alt ³
Florida	1,500	30	11	5.0	12.46	13.71
Georgia	9,000	500	250	---	5.50	13.71
Texas						
cantaloupe	22,180	22,180	5,545	---	7.25	13.71
honeydew	5,000	5,000	1,250	---	7.25	13.71
California	17,550	10,753	10,215	32.5	30.63	13.71
Total	55,230	38,463	17,271	11.2	13.77	13.71

¹Total pounds active ingredient (a.i.) applied equals average number of applications times average application rate times acres treated.

²Yield benefit equals the percent change in yield or marketable product per acre on acreage currently treated with mevinphos (versus registered alternatives).

³Average cost of chemical and/or nonchemical control alternatives that would be used on acreage currently treated with mevinphos.

Appendix Table II.12. Survey responses for lettuce: States reporting, acreage and acreage treated with mevinphos, mevinphos applied, mevinphos's yield benefit, and control costs with and without mevinphos available (average annual), 1982-86.

Crop and State	Acreage Reported		Mevinphos applied ¹ (pounds a.i.)	Yield Benefit ² (percent)	Control Costs (dollars/acre)	
	Total	Treated			Current	Alt ³
Florida	3,000	484	121	---	7.50	10.00
New York	3,700	925	347	25.0	11.25	10.00
Michigan	1,000	250	94	5.0	10.75	8.59
Wisconsin	800	600	150	10.0	15.00	8.59
Florida	13,220	1,300	853	12.5	10.06	18.58
Colorado	500	200	100	---	8.63	8.59
Texas	2,700	270	135	---	7.25	8.59
Arizona	59,126	67,573	135,146	---	33.50	18.58
California	149,450	137,898	104,803	17.5	24.00	18.58
Total	233,496	209,500	241,748	12.1	26.78	18.46

¹Total pounds active ingredient (a.i.) applied equals average number of applications times average application rate times acres treated.

²Yield benefit equals the percent change in yield or marketable product per acre on acreage currently treated with mevinphos (versus registered alternatives).

³Average cost of chemical and/or nonchemical control alternatives that would be used on acreage currently treated with mevinphos.

Appendix Table II.13. Survey responses for onion: States reporting, acreage and acreage treated with mevinphos, mevinphos applied, mevinphos's yield benefit, and control costs with and without mevinphos available (average annual), 1982-86.

Crop and State	Acreage Reported		Mevinphos applied ¹ (pounds a.i.)	Yield Benefit ² (percent)	Control Costs (dollars/acre)	
	Total	Treated			Current	Alt ³
Wisconsin	1,580	300	300	---	32.00	21.96
Florida	830	750	2,625	15.0	64.96	13.35
California	7,700	4,500	3,375	25.0	36.00	13.35
Total	10,110	5,550	6,350	23.6	32.17	13.82

¹Total pounds active ingredient (a.i.) applied equals average number of applications times average application rate times acres treated.

²Yield benefit equals the percent change in yield or marketable product per acre on acreage currently treated with mevinphos (versus registered alternatives).

³Average cost of chemical and/or nonchemical control alternatives that would be used on acreage currently treated with mevinphos.

Appendix Table II.14. Survey responses for pepper: States reporting, acreage and acreage treated with mevinphos, mevinphos applied, mevinphos's yield benefit, and control costs with and without mevinphos available (average annual), 1982-86.

Crop and State	Acreage Reported		Mevinphos applied ¹ (pounds a.i.)	Yield Benefit ² (percent)	Control Costs (dollars/acre)	
	Total	Treated			Current	Alt ³
New Jersey	7,000	1,129	282	---	7.50	10.00
Florida	19,940	6,980	2,618	2.5	9.47	15.70
Tennessee	2,000	500	94	---	7.36	12.35
Texas	9,160	9,160	2,290	---	7.25	15.70
Total	38,100	17,769	5,284	1.3	8.14	15.24

¹Total pounds active ingredient (a.i.) applied equals average number of applications times average application rate times acres treated.

²Yield benefit equals the percent change in yield or marketable product per acre on acreage currently treated with mevinphos (versus registered alternatives).

³Average cost of chemical and/or nonchemical control alternatives that would be used on acreage currently treated with mevinphos.

Appendix Table II.15. Survey responses for tomato: States reporting, acreage and acreage treated with mevinphos, mevinphos applied, mevinphos's yield benefit, and control costs with and without mevinphos available (average annual), 1982-86.

Crop and State	Acreage Reported		Mevinphos applied ¹ (pounds a.i.)	Yield Benefit ² (percent)	Control Costs (dollars/acre)	
	Total	Treated			Current	Alt ³
New Jersey	6,500	1,049	262	---	7.50	10.00
Arkansas	2,110	1,000	1,500	---	19.00	15.15
Florida	45,860	4,500	1,688	2.5	12.61	15.15
Texas	3,900	373	93	---	7.25	15.15
Total	58,370	6,922	3,543	2.3	11.10	14.37

¹Total pounds active ingredient (a.i.) applied equals average number of applications times average application rate times acres treated.

²Yield benefit equals the percent change in yield or marketable product per acre on acreage currently treated with mevinphos (versus registered alternatives).

³Average cost of chemical and/or nonchemical control alternatives that would be used on acreage currently treated with mevinphos.

Appendix Table II.16. Survey responses for spinach: States reporting, acreage and acreage treated with mevinphos, mevinphos applied, mevinphos's yield benefit, and control costs with and without mevinphos available (average annual), 1982-86.

Crop and State	Acreage Reported		Mevinphos applied ¹ (pounds a.i.)	Yield Benefit ² (percent)	Control Costs (dollars/acre)	
	Total	Treated			Current	Alt ³
Maryland	1,850	100	113	1.5	37.13	10.00
New Jersey	3,700	597	149	---	7.50	10.00
New York	1,250	688	186	---	15.00	10.00
Florida	234	125	63	10.0	9.10	6.00
Virginia	400	400	250	55.0	17.25	61.83
Oklahoma	2,500	2,000	1,000	10.0	10.40	17.26
Texas	6,000	6,000	12,000	---	14.50	6.00
California	6,365	4,000	2,280	30.0	18.38	6.00
Total	22,299	13,910	16,040	14.2	14.64	9.62

¹Total pounds active ingredient (a.i.) applied equals average number of applications times average application rate times acres treated.

²Yield benefit equals the percent change in yield or marketable product per acre on acreage currently treated with mevinphos (versus registered alternatives).

³Average cost of chemical and/or nonchemical control alternatives that would be used on acreage currently treated with mevinphos.

Appendix Table II.17. Survey responses for watermelon: States reporting, acreage and acreage treated with mevinphos, mevinphos applied, mevinphos's yield benefit, and control costs with and without mevinphos available (average annual), 1982-86.

Crop and State	Acreage Reported		Mevinphos applied ¹ (pounds a.i.)	Yield Benefit ² (percent)	Control Costs (dollars/acre)	
	Total	Treated			Current	Alt ³
New Jersey	900	145	73	---	10.00	10.00
Florida	51,710	6,000	4,500	---	16.21	21.69
Georgia	40,000	5,000	3,750	---	8.25	21.69
North Carolina	17,000	2,000	1,000	20.0	6.31	21.69
Total	109,610	13,145	9,323	1.8	10.56	21.56

¹Total pounds active ingredient (a.i.) applied equals average number of applications times average application rate times acres treated.

²Yield benefit equals the percent change in yield or marketable product per acre on acreage currently treated with mevinphos (versus registered alternatives).

³Average cost of chemical and/or nonchemical control alternatives that would be used on acreage currently treated with mevinphos.

Appendix Table II.18. Survey responses for "other" vegetables: States reporting, acreage and acreage treated with mevinphos, mevinphos applied, mevinphos's yield benefit, and control costs with and without mevinphos available (average annual), 1982-86.

Crop and State	Acreage Reported		Mevinphos applied ¹ (pounds a.i.)	Yield Benefit ² (percent)	Control Costs (dollars/acre)	
	Total	Treated			Current	Alt ³
Brussels sprouts						
California	3,094	3,000	1,710	10.0	18.38	10.00
Peas, southern						
Tennessee	300	250	141	---	22.08	26.78
Cucumber						
Florida	15,680	784	441	---	14.72	19.90
Georgia	8,000	1,000	750	---	8.25	19.90
New Jersey	1,500	242	121	---	10.00	10.00
Texas	8,100	8,100	2,025	---	7.25	19.90
Eggplant						
Florida	2,396	240	120	---	11.15	27.82
New Jersey	1,000	161	40	---	7.50	10.00
Parsley						
Florida	910	400	1,000	15.0	35.00	27.50
Pea, english						
Wisconsin	88,140	500	125	---	15.00	6.87
Squash, summer						
Florida	14,724	1,500	563	10.0	15.34	18.91
Georgia	9,000	3,000	3,750	---	12.50	18.91
Squash, winter						
Florida	1,636	150	56	10.0	15.34	17.60
Turnip						
Florida	1,199	200	100	10.0	10.00	5.91
Artichoke						
California	10,417	10,000	5,700	45.0	18.38	10.00
Bean, lima						
Florida	242	23	17	2.5	29.62	26.39
Bean, snap						
Arkansas	3,363	1,500	2,250	---	19.00	26.70
Florida	45,374	2,269	851	2.5	14.81	26.70
North Carolina	7,000	2,000	1,000	---	6.31	26.70
Tennessee	7,000	3,500	1,750	---	16.12	26.70
Wisconsin	79,760	900	338	---	14.25	18.25
Okra						
Florida	2,440	1,855	4,174	30.0	40.50	38.70
Total	311,275	41,574	27,021	12.2	11.79	18.85

¹Total pounds active ingredient (a.i.) applied equals average number of applications times average application rate times acres treated.

²Yield benefit equals the percent change in yield or marketable product per acre on acreage currently treated with mevinphos (versus registered alternatives).

³Average cost of chemical and/or nonchemical control alternatives that would be used on acreage currently treated with mevinphos.

Appendix Table II.19. Survey responses for field crops: Acreage, production, value (average annual), 1982-86.

Crop	Acreage	Production (1,000 cwt.)	Value of Production		
			Total (1,000 dollars)	Average (dollars/ cwt.)	Average (dollars/ acre)
Alfalfa, seed	133,862	1,085	105,390	97.13	787
Corn, field	706,200	66,735	296,925	4.45	420
Sorghum, forage	25,000	7,500	7,500	1.00	300
Sorghum, grain	20,045	894	3,469	3.88	173
Total	885,107		413,284		467

Appendix Table II.20. U.S. field crops production: Mevinphos-treated farm production and yield benefit, U.S. aggregate farm production, farm price, and yield benefit, by crop (average annual), 1982-86.

Crop	Mevinphos-treated		U. S. aggregate		
	Farm production (1,000 cwt.)	Yield benefit (percent)	Farm production (1,000 cwt.)	Farm price (dollars/cwt.)	Yield benefit (percent)
Alfalfa, seed ¹	805	1.5	1,651	97.13	0.7
Corn, field	195	0.0	4,167,561	4.23	0.0
Sorghum, forage ²	NA	7.5	NA	NA	0.2
Sorghum, grain ³	20	2.5	406,537	3.94	0.0

¹For U.S. aggregate production, annual average obtained from 1982 yield and acreage (U.S. Department of Commerce) adjusted for mevinphos-survey results. U.S. aggregate price taken from mevinphos survey average value.

²Florida's 25,000 acres are 0.032 of the 1982-85 U.S.D.A. average of 784,000.

³For U.S. aggregate production and price, annual average taken from 1982 Census of Agriculture (U.S. Department of Commerce).

SOURCES: For mevinphos-treated production and yield effects--1988-89 survey of states. For U.S. aggregate production and price--U.S. Department of Commerce, 1982 Census of Agriculture, National Summary; USDA, National Agricultural Statistics Service, Crop Production, annual issues; USDA, National Agricultural Statistics Service, Crop Values, annual issues.

Appendix Table II.21. Survey responses for fruit crops: Acreage, production, value (average annual), 1982-86.

Crop	Acreage	Production (1,000 cwt.)	Value of Production		
			Total (1,000 dollars)	Average (dollars/ (dollars/ cwt.) acre)	
Apple	83,680	12,585	135,530	10.77	1,620
Grape	676,700	49,315	1,043,366	21.16	1,542
Strawberry	19,102	7,555	332,503	44.01	17,407
Total	779,482		1,511,399		1,939

Appendix Table II.22. U.S. fruit production: Mevinphos-treated farm production and yield benefit, U.S. aggregate farm production, farm price, and yield benefit, by crop (average annual), 1982-86.

Crop	Mevinphos-treated		U.S. Aggregate		
	Farm production (1,000 cwt.)	Yield benefit (percent)	Farm production (1,000 cwt.)	Farm price (dollars/ cwt.)	Yield benefit (percent)
Apple	630	<1	80,827	11.44	0
Grape ¹	832	6	14,279	20.15	0
Strawberry	2,314	44	9,611	45.77	11

¹Aggregate U.S. grape production and price for fresh-market only.

SOURCES: For mevinphos-treated production and yield effects--1988-89 survey of states. For U.S. aggregate production and price--USDA, National Agricultural Statistics Service, Vegetables, annual issues; USDA, National Agricultural Statistics Service, Noncitrus Fruits and Nuts, annual issues. Annual Price Behavior of Selected Vegetables.

Appendix III

Documentation and Assumptions for Survey Response Adjustment

Adjustments to the survey responses. Site numbers are given in parentheses. (as of 4/4/89). Generally, the pivotal states in assigning values to nonresponse cells are MD, IL, FL.

Connecticut:

Delaware:

(49) Cabbage--for price, used FL; for treatments per acre, given 3-4, used 3.5

Maine:

Maryland:

(14) Spinach--application costs, given: \$4.50/A (air)-\$6.00/A (ground), used \$5.25/A. marginal benefit, given less than 2%, used 1.5%

Massachusetts:

New Hampshire:

New Jersey:

for all NJ sites, application costs, given nothing, used \$5; for alternative treatment costs, given nothing, used \$10.

(32) Spinach--used 19#/bushel applied to 700 crates (given) per acre.

(34) Broccoli--Given: total crop value \$5,500 (too low), follow-up call got price \$0.25/lb., yield 95 cwt/A

(38) Watermelon--follow-up call got price \$0.10/lb., yield 150 cwt/A

(41) Strawberry--used \$5/A application costs, for alternative costs, used \$10

(42) Apples--calculated a price using 1982-1986 value of production \$0.111 per pound; multiplied price times production (given) to arrive at value (\$12,330,000); for application costs, used \$5/A; for alternatives, used \$10

New York:

(93) Lettuce, used 20# box; for alternative treatment costs, given no satisfactory alternative, used MD

(94) Apple--for alternative treatment costs, used \$35.50

(95) Spinach--for yield, given 400-500 boxes/A, used 450 boxes/A and 19 #/box; for acreage treated, given 50-60 percent, used 55 percent; for treatments per acre, given 1-2, used 1.5.; for a.i./acre/treatment, given 0.12-0.25, used 0.18; for cost per acre per application, given \$5-\$10, used \$10.; for loss figure, given conditional "could be rejected by processor", used zero

(133) Cabbage--for price, used 5-year average of fresh and processing; for alternative treatment costs, given nothing, used NJ

Pennsylvania:

Rhode Island:

Vermont:

Illinois:

- (27) Greens--for total production, changed 750,000# to 7,500,000# to result in a yield of 30 cwt/A; a.i. application rate, given 0.5-1, used 0.75
- (29) Apples--for production, given no units, used 42# bushels

Indiana:

Iowa:

Michigan:

- (131) Celery--for material cost/A, given \$3.75-\$7.50, used \$5.75/A; for marginal benefit, given maybe 2%, used 2%
- (132) Lettuce--a.i. application rate, given 0.25-0.5, used 0.38; marginal benefit, given no yield loss, used 5% quality loss

Minnesota:

Missouri:

Ohio:

Wisconsin:

- (51) Beets, fresh--ELIMINATE--incomplete information, minor crop
- (52) Celery--ELIMINATE--incomplete information, minor crop; for alternative treatment costs, used MI
- (54) Snap beans--for average number of acres treated, given under 1,000, used 900
- (55) Cabbage--for average number of treatments per acre, given 2-3, used 2.5; for a.i. rate of application, given 0.13-0.25 and 0.5, used 0.375; for material cost per acre, given \$1.50-\$3.00, used \$2.25; for marginal benefit, given 3-5%, used 4%; for alternative treatment costs, used IL
- (56) Lettuce--for a.i. rate of application, given 0.125 to 0.5, used 0.25; for alternative treatment costs, used MI
- (57) Carrots--for average number of acres treated, given 2-300, used 250; for a.i. rate of application, given 0.13-0.25 and 0.5, used 0.375; for alternative treatment costs, used IL cabbage

Alabama:

- (17) Cabbage--for yield and price, given nothing, used FL
- (18) Broccoli--for yield and price, given nothing, used FL
- (19) Collards--for yield and price, given nothing, used GA

Arkansas:

- (24) Snap beans--for yield, used FL; for mevinphos treatment costs, given NA, used \$5@ $(0.75/0.5)$ and \$2 application costs; for alternative treatment costs, given no alternatives, used FL
- (25) Tomatoes--for yield, given nothing, used 1983-85 average from USDA; for mevinphos treatment costs, given NA, used \$5@ $(0.75/0.5)$ and \$2 application costs; for alternative treatment costs, given no alternatives, used FL

Florida:

- (62) Corn, forage--for alternative treatment costs, used \$4.79
- (63) Sorghum, forage--marginal benefit, given 5-10% used 7.5%
- (64) Sorghum--used 56#/bu; marginal benefit, given up to 5%, used 2.5%
- (65) Lima beans--for a.i. rate of application, given 0.25-0.5, used 0.375; for material cost/A, given \$2.5-\$5.0, used \$3.75; for marginal benefit, given expected occasional 5% loss, used 2.5%
- (66) Snap beans--used 30#/bu; for marginal benefit, given expected occasional 5% loss, used 2.5%
- (67) Broccoli--for a.i. application rate, given 0.25-0.5, used 0.38; material costs, given \$2.5-\$5.0, used \$3.75; for marginal benefit, given expected occasional 15% loss, used 7.5%
- (68) Cabbage--used 50#/crate; for a.i. application rate, given 0.25-0.5, used 0.38; material costs, given \$2.5-\$5.0, used \$3.75; for marginal benefit, given expected occasional 60% loss, used 30%
- (69) Cantaloupe--marginal benefit, given no severe losses, used 5%
- (70) Carrots--marginal benefit, given occasional 10% loss, used 5%
- (71) Celery--used 60#/crate; marginal benefit, given occasional 5% loss, used 2.5%
- (72) Collards--for price, used GA greens; for marginal benefit, given 5-10%, used 7.5%
- (73) Sweet corn--used 50#/crate
- (74) Cucumbers--used 48#/crate
- (75) Eggplant--used 33#/crate
- (77) Lettuce--for a.i. application rate, given 0.25-0.5, used 0.38; material costs, given \$2.5-\$5.0, used \$3.75; for marginal benefit, given 10-15%, used 12.5%
- (79) Onions, dry--for treatments per acre, given 6-8, used 7; marginal benefit, given 10-20%, used 15%
- (80) Parsley--for treatments per acre, given 4-6, used 5; marginal benefit, given 10-20%, used 15%
- (82) Peppers--used 28#/bu; for a.i. application rate, given 0.25-0.5, used 0.38; material costs, given \$2.5-\$5.0, used \$3.75; for marginal benefit, given no change in yield, used 2.5%
- (83) Potatoes--for a.i. application rate, given 0.25-0.5, used 0.38; material costs, given \$2.5-\$5.0, used \$3.75
- (84) Spinach--for marginal benefit, given a maximum 10%, used 10%
- (85) Summer squash--used 48#/bu; for marginal benefit, given expected 20% loss, used 10%
- (86) Winter squash--used 48#/bu; for marginal benefit, given expected 20% loss, used 10%
- (87) Strawberries--used 10#/12-pt flat; for treatments per acre, given 14-21, used 17.5; marginal benefit, given 10-20%, used 15%
- (88) Tomatoes--used 60#/carton; for a.i. application rate, given 0.25-0.5, used 0.38; material costs, given \$2.5-\$5.0, used \$3.75; marginal benefit, given maximum loss of 5%, used 2.5%
- (89) Turnips--marginal benefit, given NA, used 10%; for application costs/A, given NA, used \$5; for alternative treatment costs, used FL collards
- (90) Watermelon--for a.i. application rate, given 0.25-0.5, used 0.38; material costs, given \$2.5-\$5.0, used \$3.75
- (91) Cauliflower--for a.i. application rate, given 0.5-1, used 0.75;

material costs, given \$5-\$10.0, used \$7.50

(92) Okra--for treatments per acre, given 3-6, used 4.5

Georgia:

(57) Cabbage--used 50#/crate; marginal benefit, given 10%-30% yield loss and 50%-60% quality loss, used 55%; for alternative treatment costs, given multiple alternatives, used 9 treatments@\$8 treatment cost/A=\$72

(58) Greens-- for alternative treatment costs, given multiple alternatives, used 9 treatments*\$8 treatment cost/A=\$72

(59) Watermelons-- for alternative treatment costs, given nothing, used FL;

(60) Squash--used 48#/(.555 bu); for alternative treatment costs, given nothing, used FL

(61) Cucumbers--used 48#/(1.11 bu); for alternative treatment costs, given nothing, used FL;

(134) Cantaloupe-- for alternative treatment costs, given nothing, used FL

Kentucky:

Louisiana:

(23) Greens--used 3.25#/bunch and changed units of production from bunches to dozen bunches; for acres treated, given 2-300, used 250; for no. treatments per acre, given 3-4, used 3.5;; for material and application costs, use AL collards

Mississippi:

N. Carolina:

(20) Melons (watermelons)--for yield, given nothing, used USDA 1979-1981 average; for price, used GA; for material costs per acre, used \$1.31; for alternative treatment costs, given nothing, used FL

(21) Greens--for yield and price, given nothing, used NJ; for material costs, used \$2.63

(22) Cabbage--for yield, given nothing, used USDA 1979-1981 average; for price, used SC; for material costs per acre, used \$2.63

(135) Snap beans--for yield, given nothing, used TN; for material costs, used \$1.31; for alternative treatment costs, given nothing, used FL

S. Carolina:

(15) Cabbage--for acres treated, given 95-100%, used 95%

(97) Greens--for acres treated, given 95-100%, used 95%

(98) Collards--used 20#/dozen bunches; for acres treated, given 95-100%, used 95%

(99) Apples--ELIMINATE--no mevinphos use

(100) Grapes--ELIMINATE--no mevinphos use

(101) Strawberry--ELIMINATE--no mevinphos use

(102) Peach--ELIMINATE--no mevinphos use; for production units, used 48#/bu

Tennessee:

(43) Peppers--for yield, given 1800 no units, used 1800 acres of 4

tons/A pimiento and 200 acres of 117 28#-bushels/A; for a.i. rate of application, given 1/8-1/4, used 0.1875; for material costs, given \$1.41-\$2.81, used \$2.11; for application costs, given \$4.5-\$6.0 used \$5.25; for alternative treatment costs, given nothing, used (average of NJ and FL) \$12.35

(44) Southern peas--for yield, given 250, used .25(250 40# bu/A)+.75(0.5 ton/A); for value, given \$229,500, used \$185,000; for a.i. rate of application, given 1/8-1/4, used 0.1875; for material costs, given \$1.41-\$2.81, used \$2.11; for application costs, given \$4.5-\$6.0 used \$5.25; for alternative treatment costs, given nothing, used NC beans

(45) Greens--for acres, given 1,500, used 1,200; for yield, given 1,200, used 0.90(4 tons/A)+0.10(2,000 3.25#-bunches/A); for value, given \$172,500, used \$2,300,000; for a.i. rate of application, given 1/8-1/4, used 0.1875; for application costs, given \$4.5-\$6.0 used \$5.25; for alternative treatment costs, given nothing, used NC

(46) Crucifers (renamed cabbage with crop variety code=22)--for acres, given 1,000, used 1,500; for yield, given 1,000, used 27 tons/A; for value, given \$2,200,000, used \$2,630,000; for a.i. rate of application, given 1/8-1/2, used 0.562; for material costs, given \$1.41-\$5.64, used \$4.23; for application costs, given \$4.5-\$6.0, used \$5.25; for alternative treatment costs, given nothing, used NC

(47) Snap beans--for acres, given 7,000, used 2,700 fresh and 6,300 processing, average yield, used 130 30# bu/Ac. for value, given \$3,000,000, used \$5,800,000; for application costs, given \$4.5-\$6.0, used \$5.25; for alternative treatment costs, used NC

Virginia:

(130) Greens--collards, used Ag Census of 473 total acres; for yield and price, used NJ; for a.i. application rate, used 0.25#; for marginal benefit of mevinphos, used 50%/3years=17% loss/year without mevinphos; for treatment and application costs, other chemicals in mix, and alternative treatment costs, given nothing, used MD

(141) Spinach--for alternative costs, eliminate Cygon's application costs results in \$61.83; for marginal yield benefit, given 50-60 percent, used 55 percent

W. Virginia:

Colorado:

(95) Field corn--production units not given, used 56# bushels; for alternative treatment costs, used \$30

(96) Lettuce--used 46#/box, for price and alternative treatment costs per acre, used MI

Kansas:

Montana:

Nebraska:

N. Dakota:

Oklahoma:

- (16) Spinach--for yield, used Texas
- (140) Greens--for price, used TN greens

S. Dakota:

Texas:

for all items with USDA 1982-86 data, used USDA acreage, production and value, where USDA 1982-86 data unavailable, used survey response for 1982-86, for acres treated, used the survey response percentage applied to acres harvested; for material costs, given nothing, used \$5; for application costs, given \$1.50/A gy ground, \$3.00/A by air, used \$2.25/A; for applied with other fungicides, given nothing, used no; for alternative costs, given nothing, used FL

- (103) Onions--ELIMINATE--no mevinphos use
- (106) Potato--ELIMINATE--no mevinphos use
- (112) Sweet corn--ELIMINATE--no mevinphos use
- (115) Carrot--ELIMINATE--no mevinphos use

Wyoming:

Arizona:

- (11) Lettuce--for alternative treatment costs, used FL
- (12) Cole crops--for alternative treatment costs, used FL cabbage

California:

for alternative treatment costs, given nothing, used FL for matching crop types, otherwise used \$10 (on grape, artichoke, alfalfa (seed) and brussels sprouts)

(118) Alfalfa (seed)--production changed from 900,000,000 to 90,000,000; for treatments per acre, given 1-3, used 2; a.i. application rate, given 0.25-0.5, used 0.38; material costs, given \$3.5-\$4, used \$3.75; application costs, given \$6.00-\$12.00, used \$9.00; for alternative treatment costs, given nothing, used \$10

(119) Artichoke--production changed from 795,300,000 to 79,530,000; for treatments per acre, given 1-2, used 1.5; a.i. application rate, given 0.25-0.5, used 0.38; material costs, given \$3-\$3.5, used \$3.25; application costs, given \$6.00-\$12.00, used \$9.00; marginal benefit, given 40-50%, used 45%

(120) Broccoli--for a.i. application rate, given 0.25-0.5, used 0.38; application costs, given \$6.00-\$12.00, used \$9; marginal benefit, given 15-20%, used 17.5%

(122) Brussels sprouts--for prices and yields, given nothing, used USDA 1979-81 averages of crop value/production/acreage; for treatments per acre, given 1-2, used 1.5; a.i. application rate, given 0.25-0.5, used 0.38; material costs, given \$3-\$3.5, used \$3.25; application costs, given \$6.00-\$12.00, used \$9.00

(123) Cabbage--for acreage, production, and value, Ken Kido sent new data; for acreage treated, Kido says use 100%, for acres treated, given that out of 5 counties reporting that 2 counties "don't use mevinphos much", settle on 5,500 acres treated; for treatments per acre, given 1-2, used 1.75;

a.i. application rate, given 0.25-0.5, used 0.38; material costs, given \$3-\$6, used \$4.5; application costs, given \$6.00-\$12.00, used \$9.00

(124) Cauliflower--for a.i. application rate, given 0.25-0.75, used 0.5; material costs, given \$3-\$3.5, used \$3.25; application costs, given \$6.00-\$12.00, used \$9; marginal benefit, given 40-50%, used 45%

(125) Celery--for a.i. application rate, given 0.25-0.5, used 0.38; application costs, given \$6.00-\$12.00, used \$9; marginal benefit, given 35-40%, used 37.5%

(126) Grape--for treatments per acre, given 1-2, used 1.5; a.i. application rate, given 0.25-0.5, used 0.38; material costs, given \$3-\$3.5, used \$3.25; application costs, given \$6.00-\$12.00, used \$9.00; for alternative costs, used \$10

(127) Lettuce--for a.i. application rate, given 0.25-0.5, used 0.38; application costs, given \$6.00-\$12.00, used \$9.00; marginal benefit, given 15-20%, used 17.5%

(129) Melons--for treatments per acre, given 1-4, used 2; a.i. application rate, given 0.25-0.5, used 0.38; material costs, given \$3-\$3.5, used \$3.25; application costs, given \$6.00-\$12.00, used \$9.00; marginal benefit, given 30-35%, used 32.5%

(136) Corn--for treatments per acre, given 1-3, used 2; a.i. application rate, given 0.25-0.5, used 0.38; material costs, given \$3-\$3.5, used \$3.25; application costs, given \$6.00-\$12.00, used \$9.00; marginal benefit, given 15-20%, used 17.5%

(139) Spinach--production changed from 8,050,800,000 to 80,508,000; for treatments per acre, given 1-2, used 1.5; a.i. application rate, given 0.25-0.5, used 0.38; material costs, given \$3-\$6, used \$4.5; application costs, given \$6.00-\$12.00, used \$9.00

(137) Strawberries--for acreage, production, and value, Ken Kido sent new data; for acres treated, Kido sent acre-treatments, use acre-treatments divided by 3 treatments per acre, round to 3,000 acres treated; for treatments per acre, use 3; a.i. application rate, given 0.25-0.5, used 0.38; material costs, given \$3-\$3.5, used \$3.25; application costs, given \$6.00-\$15.00, used \$10.50; for alternative costs, used FL

(138) Onion--for application costs, given \$6.00-\$12.00, used \$9

Nevada:

New Mexico:

Utah:

(9) Cabbage--for yield and price, nothing given, used WI; for alternative treatment costs, given nothing, used WI

Idaho:

(26) Potatoes--ELIMINATE--acreage treated=100/300,000 planted.

Oregon:

(1) Broccoli--a.i. application rate, given 0.25 aphids-0.5 worms, used 0.38; material costs per acre, given \$11.37 for aphids-\$22.64 for worms and aphids, used \$17.00;

(2) Cauliflower--a.i. application rate, given 0.25 aphids-0.5 worms, used 0.38; material costs per acre, given \$11.37 for aphids-\$22.64 for worms

and aphids, used \$17.00;

(3) Cucumber processing--ELIMINATE--no information beyond acreage, production, and value

(4) Cucumber fresh--ELIMINATE--no information beyond acreage, production, and value

(5) all broccoli--use site (1)

(6) Carrots--ELIMINATE--no information beyond acreage, production, and value

(7) Onions, dry--ELIMINATE--no information beyond acreage, production, and value

(8) Alfalfa seed--for marginal benefit, mevinphos costs, and alternative treatment costs, used California

Washington:

(13) Alfalfa seed--for acreage treated, given 5,000-7,000, used 6,000; for treatments per acre, given 2-3, used 2.5; for marginal benefit, given 10-15%, used 12.5%; for alternative treatment costs, given nothing, used \$10

Alaska:

other territories:

APPENDIX IV

Price-quantity Relationships

The Data and Methods

The data are obtained from public sources including USDA Agricultural Statistics, Vegetable and Fruit Outlook and Situation reports, Cold Storage reports, and trade publications. The data series for this analysis are available from the authors.

The farm-level annual average price data are the ratios of annual value of production to annual total quantity of production, both reported by the USDA National Agricultural Statistics Service or NASS. The NASS estimates are compilations of cooperating states' estimates and do not include all production (as compared to a census survey). The major states, though, are included in the NASS estimates and are likely to indicate accurate changes in domestic prices.

Annual per capita use is calculated from annual domestic production, net exports, and net carry-over stocks. The per capita use (formerly known as per capita consumption) time series is not perfectly consistent. Occasionally a cooperating state will drop out of the survey or NASS will drop a crop such as asparagus and then reinstate it. In the case of lettuce, broccoli, and cauliflower, California changed its reporting schedule from annual to seasonal and rendered 1983 data not comparable to previous years. This may be important, but its impact on the continuity of the time series of data is difficult to estimate objectively.

"Other price" is the farm-level average price of the major fresh-market vegetables, excluding the commodity being regressed. All prices are deflated by the Gross National Product deflator (1982=100), and per capita disposable personal income is measured in 1982 dollars (Reagan). The other two categories of variables are continuous and binary. The binary variables are

specified as zero or one as defined below.

"Other variables": for broccoli and cauliflower; A = January 1 stocks of frozen product divided by sum of previous year's January 1 stocks plus frozen pack. B = per capita consumption of processed product divided by per capita consumption total product. For celery, lettuce, and fresh-market vegetables, A = disposable personal income per capita (\$1982). For strawberry, B = per capita consumption of processed product divided by per capita consumption total product.

"Binary variables": for broccoli, A: 1986 = 1; for cauliflower, A: 1976-1988 = 1; for celery, A: 1956, 1959, 1961, 1963, 1974, 1987 = 1; B: 1955, 1958, 1962, 1968, 1978, 1983 = 1; for lettuce, A: 1961, 1970, 1972, 1975, 1977, 1980, 1984-86 = 1; B: 1955, 1971, 1973, 1978, 1982, 1988 = 1; for spinach, A: 1961, 1964, 1977 = 1; B: 1972 = 1; for cabbage, A: 1961, 1967, 1969, 1972, 1974, 1981 = 1; B: 1955, 1962, 1966, 1973, 1977, 1978 = 1; for strawberry, A: 1957, 1965 = 1; B: 1955, 1956 = 1; cantaloupe, A: 1958, 1962, 1978 = 1; B: trend 1954, 1955, . . . 1981; fresh-market vegetables, A: 1970 = 1; B: 1954-1957 = 1.

The signs on the estimated coefficients, the adjusted R-squared, and Durbin-Watson statistics suggest the models are reasonable and generally correct.

Appendix Table IV.1. Selected results from Ordinary Least Squares regression of annual average farm-level price of vegetables.¹

Crop	Estimated coefficients						R-sq. ²	DW ³	
	Con- stant	Per capita use	Other price	Other variables		Binary variables			
				A	B	A			B
Broccoli Std. err.	33.95 (4.48)	-2.34 (0.29)	0.45 (0.24)	-7.79 (5.98)	-15.96 (3.68)	-3.26 (1.30)	0.81	1.84	
Cauliflower Std. err.	12.18 (4.85)	-1.60 (0.61)	0.92 (0.31)	-0.87 (4.74)	1.60 (2.35)	3.62 (0.88)	0.56	1.48	
Celery Std. err.	12.82 (3.22)	-1.00 (0.26)	0.74 (0.14)	-4E-04 (8E-05)		-1.72 (0.36)	2.75 (0.35)	0.84	1.58
Lettuce Std. err.	12.26 (1.91)	-0.34 (0.11)	0.63 (0.12)	-7E-05 (1E-04)		-1.76 (0.28)	1.23 (0.32)	0.83	1.49
Spinach Std. err.	29.80 (1.21)	-1.10 (0.11)	0.60 (0.11)			-2.97 (0.81)	4.29 (1.37)	0.91	1.76
Cabbage Std. err.	15.45 (2.83)	-0.96 (0.12)	0.20 (0.14)			-1.16 (0.36)	2.33 (0.36)	0.86	1.57
Strawberry Std. err.	45.24 (13.4)	-5.45 (2.17)			58.61 (16.3)	-8.73 (2.80)	18.2 (2.94)	0.87	1.70
Cantaloupe Std. err.	100.5 (33.4)	-1.30 (0.19)	0.66 (0.17)			-2.01 (0.40)	-0.04 (0.02)	0.81	2.75
Fresh-market vegetables	34.48	-0.37 (0.04)		0.001 (2E-04)		-2.81 (0.73)	2.59 (0.50)	0.72	2.30

¹For variable definitions and source material, see the text of Appendix B.

²R-squared, the multiple coefficient of determination, is adjusted for degrees of freedom.

³Durbin-Watson statistic indicates a two-sided test for rho equal to zero is inconclusive or fails to reject at the 10 percent level of significance for all estimated models, except celery.

SOURCE: Data available from the authors.

NATIONAL AGRICULTURAL LIBRARY



1022337173

NATIONAL AGRICULTURAL LIBRARY



1022337173